$\Delta; \Phi \vdash \kappa_1 \lesssim \kappa_2$ κ_1 is a consistent subkind of κ_2

$$\begin{array}{ll} {\rm KCHoleL} & {\rm KCHoleR} \\ \hline \Delta; \Phi \vdash {\rm KHole} \lesssim \kappa & \overline{\Delta}; \Phi \vdash \kappa \lesssim {\rm KHole} \\ \hline \\ & \overline{\Delta}; \Phi \vdash \kappa_1 \equiv \kappa_2 \\ \hline \Delta; \Phi \vdash \kappa_1 \lesssim \kappa_2 \\ \hline \\ & \overline{\Delta}; \Phi \vdash \kappa_1 \lesssim \kappa_2 \\ \hline \\ & \overline{\Delta}; \Phi \vdash \kappa_1 \lesssim \kappa_2 \\ \hline \end{array}$$

t valid t is a valid type variable

t is valid if it is not a built in-type or keyword, begins with an alpha char or underscore, and only contains alpha numeric characters, underscores, and primes.

 $\Delta; \Phi \vdash \kappa \text{ kind}$ $\kappa \text{ forms a kind}$

$$\frac{\text{KFTy}}{\Delta;\Phi\vdash \text{Ty kind}} \qquad \frac{\text{KFHole}}{\Delta;\Phi\vdash \text{KHole kind}} \qquad \frac{\Delta;\Phi\vdash\tau \rightleftarrows \text{Ty}}{\Delta;\Phi\vdash \text{S}(\tau) \text{ kind}} \text{ 43.2a}$$

 $\Delta; \Phi \vdash \kappa_1 \equiv \kappa_2$ κ_1 is equivalent to κ_2

$$\begin{tabular}{lll} {\tt KERefl} & & {\tt KESymm} \\ \hline $\Delta; \Phi \vdash \kappa_1 \equiv \kappa_2$ & $\Delta; \Phi \vdash \kappa_1 \equiv \kappa_2$ & $\Delta; \Phi \vdash \kappa_1 \equiv \kappa_3$ \\ \hline $\Delta; \Phi \vdash \kappa \equiv \kappa$ & $\Delta; \Phi \vdash \kappa_2 \equiv \kappa_1$ & $\Delta; \Phi \vdash \kappa_1 \equiv \kappa_3$ \\ \hline \end{tabular}$$

KESingEquiv
$$\frac{\Delta; \Phi \vdash \tau_1 \equiv \tau_2}{\Delta; \Phi \vdash S(\tau_1) \equiv S(\tau_2)}$$

 $\Delta; \Phi \vdash \tau \Rightarrow \kappa$ τ synthesizes kind κ define labelled sing letter for khole

$$\frac{\Delta; \Phi \vdash \tau_1 \Leftarrow \mathtt{S}(\tau_1)}{\Delta; \Phi \vdash \tau_1 \oplus \tau_2 \Rightarrow \mathtt{S}(\tau_1 \oplus \tau_2)}$$

$$\frac{\Delta; \Phi \vdash \tau \Leftarrow S(\tau)}{\Delta; \Phi \vdash \text{list}(\tau) \Rightarrow S(\text{list}(\tau))} \qquad \frac{\text{KSEHole}}{u :: \kappa \in \Delta} \\
\frac{u :: \kappa \in \Delta}{\Delta; \Phi \vdash ()^u \Rightarrow \kappa}$$

$$\frac{u :: \kappa \in \Delta \qquad \Delta; \Phi \vdash \tau \Rightarrow \kappa'}{\Delta; \Phi \vdash (\!(\tau)\!)^u \Rightarrow \kappa}$$

 $\Delta; \Phi \vdash \tau \leftarrow \kappa$ τ analyzes against kind κ

$$\frac{\Phi \vdash \tau \Rightarrow \kappa' \qquad \Delta; \Phi \vdash \kappa' \lesssim \kappa}{\Delta; \Phi \vdash \tau \Leftarrow \kappa} \qquad \text{KAVar} \\ \frac{L :: K_1 \in \overline{\Phi} \qquad k_1 \leq k}{\Delta j \underline{\Phi} \vdash t \Leftarrow k}$$

 $\Delta; \Phi \vdash \tau_1 \equiv \tau_2 \mid \tau_1 \text{ is equivalent to } \tau_2$

KCESymm
$$\frac{\Delta; \Phi \vdash \tau_1 \equiv \tau_2}{\Delta; \Phi \vdash \tau_2 \equiv \tau_1}$$

$$\frac{\Delta; \Phi \vdash \tau_1 \equiv \tau_2 \qquad \Delta; \Phi \vdash \tau_2 \equiv \tau_3}{\Delta; \Phi \vdash \tau_1 \equiv \tau_3}$$

KCESingEquiv
$$\frac{\Delta; \Phi \vdash \tau_1 \iff S(\tau_2)}{\Delta; \Phi \vdash \tau_1 \equiv \tau_2}$$

KCEREF

SEREF KCEConst
$$\Delta; \Phi \vdash c \equiv c$$

$$\frac{t: k \in \Phi}{\Delta; \Phi \vdash k \equiv}$$

$$\begin{tabular}{ll} {\tt KCEVar} & {\tt KCEBinOp} \\ \hline $t: \kappa \in \Phi$ & $\Delta; \Phi \vdash \tau_1 \equiv \tau_2$ & $\Delta; \Phi \vdash \tau_3 \equiv \tau_4$ \\ \hline $\Delta; \Phi \vdash t \equiv t$ & $\Delta; \Phi \vdash \tau_1 \oplus \tau_3 \equiv \tau_2 \oplus \tau_4$ \\ \hline \end{tabular}$$

$$\frac{\Delta; \Phi \vdash \tau_1 \equiv \tau_2}{\Delta; \Phi \vdash \mathtt{list}(\tau_1) \equiv \mathtt{list}(\tau_2)} \qquad \frac{u :: \kappa \in \Delta}{\Delta; \Phi \vdash ()^u \equiv ()^u}$$

$$\frac{u :: \kappa \in \Delta}{\Delta; \Phi \vdash ()^u \equiv ()^u}$$

KCENEHole $\frac{u :: \kappa \in \Delta \qquad \Delta; \Phi \vdash \tau \underset{\longleftarrow}{\overset{*}{\smile}} \kappa'}{\Delta; \Phi \vdash (|\tau|)^u \equiv (|\tau|)^u}$

Aj更H T::K |

T is well formed at kind k

$$\frac{\text{difft:T}}{\text{difft:K}} | \text{f} \qquad \frac{\text{difft:T}}{\text{difft:K}} = \frac{\text{difft:K}_{1} \text{ diffk}, \text{k}_{2}}{\text{difft:K}} | \text{difft:K}_{1} \text{ difft:K}_{2} | \text{difft:K}_{2} | \text{difft:K}_{3} | \text{difft:K}_{4} | \text{difft:K}_{2} | \text{difft:K}_{3} | \text{difft:K}_{4} | \text{difft:K}_{4} | \text{difft:K}_{5} | \text{difft:K}_{5} | \text{difft:K}_{6} | \text{difft:K$$

DIEFT :: S. (T) replaces

 $\frac{\triangle i \, \mathbb{E} \vdash T_1 :: T_2 \qquad \triangle i \, \mathbb{E} \vdash T_2 :: T_2}{\triangle i \, \mathbb{E} \vdash T_1 \, \mathfrak{G} \, T_2 :: S \, \mathcal{L} \, T_1 \, \mathfrak{A}} \qquad \frac{\mathsf{t} \, \# \, \mathsf{dom}(\mathbb{E})}{\triangle i \, \mathbb{E} \vdash \, \mathbf{t} \, :: \, \mathsf{kHole}}$

 $\Phi \vdash \hat{\tau} \Rightarrow \kappa \leadsto \tau \dashv \Delta$ $\hat{\tau}$ synthesizes kind κ and elaborates to τ

$$\overline{\Phi \vdash c \Rightarrow S(c) \rightsquigarrow c \dashv \cdot}$$

TElabSBinOp

$$\frac{\Phi \vdash \hat{\tau}_1 \Leftarrow \mathsf{Ty} \leadsto \tau_1 \dashv \Delta_1 \qquad \Phi \vdash \hat{\tau}_2 \Leftarrow \mathsf{Ty} \leadsto \tau_2 \dashv \Delta_2}{\Phi \vdash \hat{\tau}_1 \oplus \hat{\tau}_2 \Rightarrow \mathsf{S}(\tau_1 \oplus \tau_2) \leadsto \tau_1 \oplus \tau_2 \dashv \Delta_1 \cup \Delta_2}$$

TElabSList

$$\frac{\Phi \vdash \hat{\tau} \Leftarrow \mathsf{Ty} \leadsto \tau \dashv \Delta}{\Phi \vdash \mathsf{list}(\hat{\tau}) \Rightarrow \mathsf{S}(\mathsf{list}(\tau)) \leadsto \mathsf{list}(\tau) \dashv \Delta} \qquad \frac{t : \kappa \in \Phi}{\Phi \vdash t \Rightarrow \mathsf{S}(t) \leadsto t \dashv \cdot}$$

TElabSUVar

$$\frac{t \not\in \mathsf{dom}(\Phi)}{\Phi \vdash t \Rightarrow \mathsf{KHole} \leadsto (\!\!|t|\!\!)^u \dashv u :: \mathsf{KHole}} \qquad \frac{\mathsf{TElabSHole}}{\Phi \vdash (\!\!|)^u \Rightarrow \mathsf{KHole} \leadsto (\!\!|)^u \dashv u :: \mathsf{KHole}}$$

$$\overline{\Phi \vdash ()^u \Rightarrow \mathsf{KHole} \leadsto ()^u \dashv u :: \mathsf{KHole}}$$

$$\frac{\Phi \vdash \hat{\tau} \Rightarrow \kappa \leadsto \tau \dashv \Delta}{\Phi \vdash (\!(\hat{\tau})\!)^u \Rightarrow \mathtt{KHole} \leadsto (\!(\tau)\!)^u \dashv \Delta, u :: \mathtt{KHole}}$$

TElabASubsume

$$\frac{\hat{\tau} \neq (\!\!|)^u \qquad \hat{\tau} \neq (\!\!|\hat{\tau}'|)^u \qquad \Phi \vdash \hat{\tau} \Rightarrow \kappa' \leadsto \tau \dashv \Delta \qquad \Delta; \Phi \vdash \kappa' \lesssim \kappa}{\Phi \vdash \hat{\tau} \Leftarrow \kappa \leadsto \tau \dashv \Delta}$$

$$\overline{\Phi \vdash ()^u \leftarrow \kappa \leadsto ()^u \dashv u :: \kappa}$$

$$\frac{\texttt{TElabAnehole}}{\Phi \vdash (\!|\!|)^u \Leftarrow \kappa \leadsto (\!|\!|)^u \dashv u :: \kappa} \qquad \frac{\frac{\texttt{TElabAnehole}}{\Phi \vdash \hat{\tau} \Rightarrow \kappa' \leadsto \tau \dashv \Delta}}{\Phi \vdash (\!|\!|\hat{\tau}|\!|)^u \Leftarrow \kappa \leadsto (\!|\tau|\!|)^u \dashv \Delta, u :: \kappa}$$

 $\Phi_1 \vdash \tau : \kappa \rhd \rho \dashv \Phi_2$ ρ matches against $\tau : \kappa$ extending Φ if necessary

RESVar

$$\frac{}{\Phi \vdash \tau : \kappa \rhd (\!\!\!) \dashv \Phi}$$

RESVarHole

$$\frac{\neg (t \ \forall a = 1 \ d)}{\Phi \vdash \tau : \kappa \rhd (|t|) \dashv \Phi}$$

 $\Gamma; \Phi \vdash e \Rightarrow \hat{\tau} \leadsto d \dashv \Delta$ e synthesizes type τ and elaborates to d

ESDefine

$$\begin{split} & \Phi_1 \vdash \hat{\tau} \Rightarrow \kappa \leadsto \tau \dashv \Delta_1 \\ & \Phi_1 \vdash \tau : \kappa \rhd \rho \dashv \Phi_2 \qquad \Gamma; \Phi_2 \vdash e \Rightarrow \tau_1 \leadsto d \dashv \Delta_2 \\ \hline & \Gamma; \Phi_1 \vdash \mathsf{type} \ \rho = \hat{\tau} \ \mathsf{in} \ e \Rightarrow \tau_1 \leadsto \mathsf{type} \ \rho = \tau : \kappa \ \mathsf{in} \ d \dashv \Delta_1 \cup \Delta_2 \end{split}$$

 $\Delta; \Gamma; \Phi \vdash d : \tau$ d is assigned type τ

$$\begin{array}{ll} \text{DEDefine} \\ \underline{\Phi_1 \vdash \tau_1 : \kappa \rhd \rho \dashv \Phi_2} & \Delta; \Gamma; \underline{\Phi_2 \vdash d : \tau_2} \\ \underline{\Delta; \Gamma; \Phi_1 \vdash \text{type} \ \rho = \tau_1 : \kappa \ \text{in} \ d : \tau_2} \end{array}$$

Theorem 1 (Well-Kinded Elaboration)

- (1) If $\Phi \vdash \hat{\tau} \Rightarrow \kappa \leadsto \tau \dashv \Delta \ then \ \Delta; \Phi \vdash \tau \Rightarrow \kappa$
- (2) If $\Phi \vdash \hat{\tau} \Leftarrow \kappa \leadsto \tau \dashv \Delta \ then \ \Delta; \Phi \vdash \tau \Leftarrow \kappa$

This is like the Typed Elaboration theorem in the POPL19 paper.

Theorem 2 (Elaborability)

- (1) $\exists \Delta \ s.t. \ if \ \Delta; \Phi \vdash \tau \Rightarrow \kappa \ then \ \exists \hat{\tau} \ such \ that \ \Phi \vdash \hat{\tau} \Rightarrow \kappa \leadsto \tau \dashv \Delta$
- (2) $\exists \Delta \ s.t. \ if \ \Delta; \Phi \vdash \tau \Leftarrow \kappa \ then \ \exists \hat{\tau} \ such \ that \ \Phi \vdash \hat{\tau} \Leftarrow \kappa \leadsto \tau \dashv \Delta$

This is similar but a little different from Elaborability theorem in the POPL19 paper. Choose the Δ that is emitted from elaboration and then there's an $\hat{\tau}$ that elaborates to any of the τ forms. Elaborability and Well-Kinded Elaboration implies we can just rely on the elaboration forms for the premises of any rules that demand kind synthesis/analysis.

Theorem 3 (Type Elaboration Unicity)

- (1) If $\Phi \vdash \hat{\tau} \Rightarrow \kappa_1 \leadsto \tau_1 \dashv \Delta_1$ and $\Phi \vdash \hat{\tau} \Rightarrow \kappa_2 \leadsto \tau_2 \dashv \Delta_2$ then $\kappa_1 = \kappa_2$, $\tau_1 = \tau_2$, $\Delta_1 = \Delta_2$
- (2) If $\Phi \vdash \hat{\tau} \Leftarrow \kappa \leadsto \tau_1 \dashv \Delta_1$ and $\Phi \vdash \hat{\tau} \Leftarrow \kappa \leadsto \tau_2 \dashv \Delta_2$ then $\tau_1 = \tau_2$, $\Delta_1 = \Delta_2$

This is like the Elaboration Unicity theorem in the POPL19 paper.

Theorem 4 (Kind Synthesis Precision) If Δ ; $\Phi \vdash \tau \Rightarrow \kappa_1$ and Δ ; $\Phi \vdash \tau \Leftarrow \kappa_2$ then Δ ; $\Phi \vdash \kappa_1 \lesssim \kappa_2$

Kind Synthesis Precision says that synthesis finds the most precise kappa possible for a given input type. This is somewhat trivial, but interesting to note because it means we can expect singletons wherever possible.

Theorem 5.1 (Kind Analysis Soundness)
If AjEtt \(\) K then AjEtt:: K

Theorem S.Z (Kind Analysis Completeners)
IF DJEHT:: K then AJEHT = K

5.1 Induction on complexity

Base cares

C::万 i k=5 当 75と73 司 C:: K

C + K 司 C司K' i K' を k 司 K' = S(c) コ S(c) た k 司 S(c) き k 司 C:: S(c) i s (c) ら k 司 C:: k

t + k 司 t 司 k' i k' ら k 司 k' > S(t) i t :: k''も 上 司 t :: l コ t :: l コ t :: l コ t :: k

t + k

L + S(t) コ k + S(t) コ k + S(t) コ k + S(t) コ k + S(t) コ