

# Pseudorandomness (III) & One-Way Functions

601.642/442: Modern Cryptography

Fall 2022

## Recap: One-bit stretch PRG $\implies$ Poly-bit stretch PRG

- A PRG with **one-bit stretch** can be used to construct a PRG with arbitrary **polynomial-bit stretch**.

## Recap: One-bit stretch PRG $\implies$ Poly-bit stretch PRG

- A PRG with **one-bit stretch** can be used to construct a PRG with arbitrary **polynomial-bit stretch**.

Construction of  $G_{poly} : \{0, 1\}^n \rightarrow \{0, 1\}^{\ell(n)}$

Let  $G : \{0, 1\}^n \rightarrow \{0, 1\}^{n+1}$  be a one-bit stretch PRG.

$$\begin{aligned} s &= x_0 \\ G(x_0) &= x_1 \| b_1 \\ &\vdots \\ G(x_{\ell(n)-1}) &= x_{\ell(n)} \| b_{\ell(n)} \end{aligned}$$

$$G_{poly}(s) := b_1 \dots b_{\ell(n)}$$

## Recap: Pseudorandomness of $G_{poly}$

- In order to show  $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$  we considered the following hybrid experiments:

## Recap: Pseudorandomness of $G_{poly}$

- In order to show  $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$  we considered the following hybrid experiments:

<u>Experiment <math>\mathcal{H}_1</math></u>	<u>Experiment <math>\mathcal{H}_2</math></u>	<u>Experiment <math>\mathcal{H}_{\ell(n)}</math></u>
$s = x_0$	$s = x_0$	$s = X_0$
$G(x_0) = x_1    b_1$	$s_1    u_1 = x_1    u_1$	$s_1    u_1 = x_1    u_1$
$G(x_1) = x_2    b_2$	$G(x_1) = x_2    b_2$	$s_2    u_2 = x_2    u_2$
$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$
$G(x_{\ell(n)-1}) = x_{\ell(n)}    b_{\ell(n)}$	$G(X_{\ell(n)-1}) = x_{\ell(n)}    b_{\ell(n)}$	$s_{\ell(n)}    u_{\ell(n)} = x_{\ell(n)}    u_{\ell(n)}$
Output $G(s) := b_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 u_2 \dots u_{\ell(n)}$

## Recap: Pseudorandomness of $G_{poly}$

- In order to show  $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$  we considered the following hybrid experiments:

<u>Experiment <math>\mathcal{H}_1</math></u>	<u>Experiment <math>\mathcal{H}_2</math></u>	<u>Experiment <math>\mathcal{H}_{\ell(n)}</math></u>
$s = x_0$	$s = x_0$	$s = X_0$
$G(x_0) = x_1    b_1$	$s_1    u_1 = x_1    u_1$	$s_1    u_1 = x_1    u_1$
$G(x_1) = x_2    b_2$	$G(x_1) = x_2    b_2$	$s_2    u_2 = x_2    u_2$
$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$
$G(x_{\ell(n)-1}) = x_{\ell(n)}    b_{\ell(n)}$	$G(X_{\ell(n)-1}) = x_{\ell(n)}    b_{\ell(n)}$	$s_{\ell(n)}    u_{\ell(n)} = x_{\ell(n)}    u_{\ell(n)}$
Output $G(s) := b_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 u_2 \dots u_{\ell(n)}$

- And established that  $\forall i \in [\ell(n) - 1], \mathcal{H}_i \approx_c \mathcal{H}_{i+1}$ .

# Contrapositive Point of View

- So far, we have proved statements in the forward direction.

*“If  $G$  is a one-bit stretch PRG, then  $G_{poly}$  is a poly-bit stretch PRG.”*

# Contrapositive Point of View

- So far, we have proved statements in the forward direction.

*“If  $G$  is a one-bit stretch PRG, then  $G_{poly}$  is a poly-bit stretch PRG.”*

- Let's now think about the **contrapositive** of these statements.

*“If  $G_{poly}$  is a **not** poly-bit stretch PRG, then  $G$  is **not** a one-bit stretch PRG.”*



# Contrapositive Point of View

- So far, we have proved statements in the forward direction.  
*“If  $G$  is a one-bit stretch PRG, then  $G_{poly}$  is a poly-bit stretch PRG.”*
- Let’s now think about the **contrapositive** of these statements.  
*“If  $G_{poly}$  is a **not** poly-bit stretch PRG, then  $G$  is **not** a one-bit stretch PRG.”*
- If  $G_{poly}$  is not a PRG, then there exists a n.u. PPT adversary  $\mathcal{A}$  who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage  $\mu$ .

# Contrapositive Point of View

- So far, we have proved statements in the forward direction.  
*“If  $G$  is a one-bit stretch PRG, then  $G_{poly}$  is a poly-bit stretch PRG.”*
- Let’s now think about the **contrapositive** of these statements.  
*“If  $G_{poly}$  is a **not** poly-bit stretch PRG, then  $G$  is **not** a one-bit stretch PRG.”*
- If  $G_{poly}$  is not a PRG, then there exists a n.u. PPT adversary  $\mathcal{A}$  who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage  $\mu$ .

# Contrapositive Point of View

- So far, we have proved statements in the forward direction.

*“If  $G$  is a one-bit stretch PRG, then  $G_{poly}$  is a poly-bit stretch PRG.”*

- Let’s now think about the **contrapositive** of these statements.

*“If  $G_{poly}$  is a **not** poly-bit stretch PRG, then  $G$  is **not** a one-bit stretch PRG.”*

- If  $G_{poly}$  is not a PRG, then there exists a n.u. PPT adversary  $\mathcal{A}$  who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage  $\mu$ .

Can we somehow use  $\mathcal{A}$  to also break security of  $G$ ?

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.

## Lemma (Alternate way to state Hybrid Lemma)

*Let  $X^1, \dots, X^m$  be distribution ensembles for  $m = \text{poly}(n)$ . Suppose there exists a distinguisher/adversary  $\mathcal{A}$  that distinguishes between  $X^1$  and  $X^m$  with probability  $\mu$ . Then  $\exists i \in [m-1]$ , such that  $\mathcal{A}$  distinguishes between  $X^i$  and  $X^{i+1}$  with advantage at least  $\mu/m$ .*



# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.
- In our previous proof, we relied on the security of  $G$  to argue indistinguishability of each pair of consecutive hybrids.

## Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.
- In our previous proof, we relied on the security of  $G$  to argue indistinguishability of each pair of consecutive hybrids.
- We will now use  $\mathcal{A}$  that has non-negligible advantage in distinguishing between  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ , to construct another adversary  $\mathcal{B}$  to break security of  $G$ .

# Contrapositive Point of View

- We just proved security of  $G_{poly}$  using a sequence of hybrids.
- If we assume that an adversary  $\mathcal{A}$  exists, who has non-negligible advantage in breaking the security of  $G_{poly}$ , then at least one of the steps of our previous proof must break down.
- By hybrid lemma,  $\mathcal{A}$  can distinguish between at least 1 pair of consecutive hybrids (say  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ ) with at least  $\mu/\ell(n)$  advantage.
- In our previous proof, we relied on the security of  $G$  to argue indistinguishability of each pair of consecutive hybrids.
- We will now use  $\mathcal{A}$  that has non-negligible advantage in distinguishing between  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ , to construct another adversary  $\mathcal{B}$  to break security of  $G$ .
- However, since  $G$  is a secure PRG, no such n.u. PPT  $\mathcal{A}$  should exist. This will give us a contradiction and imply that our assumption was incorrect.  $G_{poly}$  is in fact secure.

# Proof via Reduction

- How do we construct  $\mathcal{B}$ ?

# Proof via Reduction


- How do we construct  $\mathcal{B}$ ?
- We consider the game-based definition of PRG.

# Proof via Reduction

- How do we construct  $\mathcal{B}$ ?
- We consider the game-based definition of PRG.

# Proof via Reduction

- How do we construct  $\mathcal{B}$ ?
- We consider the game-based definition of PRG.



$$\left| \Pr[b' = 1 | b = 1] - \Pr[b' = 1 | b = 0] \right| \leq \nu(n)$$


# Proof by Reduction

- This proof technique is also called **proof by reduction**.



# Proof by Reduction

- This proof technique is also called **proof by reduction**.



# Proof by Reduction

- If  $y$  is pseudorandom, i.e., sampled as  $y = G(s)$ , then the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_i$ .

# Proof by Reduction

- If  $y$  is pseudorandom, i.e., sampled as  $y = G(s)$ , then the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_i$ .
- Otherwise, i.e.,  $y$  is (truly) random, and therefore the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_{i+1}$ .

# Proof by Reduction

- If  $y$  is pseudorandom, i.e., sampled as  $y = G(s)$ , then the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_i$ .
- Otherwise, i.e.,  $y$  is (truly) random, and therefore the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_{i+1}$ .
- Hence,  $\mathcal{B}$  has the **same advantage** in distinguishing between the output of  $G$  and a pseudorandom string that  $\mathcal{A}$  has in distinguishing between  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ .

# Proof by Reduction

- If  $y$  is pseudorandom, i.e., sampled as  $y = G(s)$ , then the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_i$ .
- Otherwise, i.e.,  $y$  is (truly) random, and therefore the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_{i+1}$ .
- Hence,  $\mathcal{B}$  has the **same advantage** in distinguishing between the output of  $G$  and a pseudorandom string that  $\mathcal{A}$  has in distinguishing between  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ .
- Moreover, since  $\mathcal{A}$  is n.u. PPT, so is  $\mathcal{B}$ . This is a contradiction!

# Proof by Reduction

- If  $y$  is pseudorandom, i.e., sampled as  $y = G(s)$ , then the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_i$ .
- Otherwise, i.e.,  $y$  is (truly) random, and therefore the input to  $\mathcal{A}$  is distributed identically to the output of  $\mathcal{H}_{i+1}$ .
- Hence,  $\mathcal{B}$  has the **same advantage** in distinguishing between the output of  $G$  and a pseudorandom string that  $\mathcal{A}$  has in distinguishing between  $\mathcal{H}_i$  and  $\mathcal{H}_{i+1}$ .
- Moreover, since  $\mathcal{A}$  is n.u. PPT, so is  $\mathcal{B}$ . This is a contradiction!
- Hence,  $G_{\text{poly}(n)}$  is a PRG.

# Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:

# Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
  - ① **Input Mapping:** How to map the input that “outer adversary”  $\mathcal{B}$  receives from the challenger to an input to the “internal adversary”  $\mathcal{A}$ ?



# Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
  - 1 **Input Mapping:** How to map the input that “outer adversary”  $\mathcal{B}$  receives from the challenger to an input to the “internal adversary”  $\mathcal{A}$ ?
  - 2 **Input Distribution:** Does the input mapping provide the right distribution of inputs that  $\mathcal{A}$  expects?

# Proof by Reduction: Key Points


- These are four important things that you must work through for a valid reduction:
  - ① **Input Mapping:** How to map the input that “outer adversary”  $\mathcal{B}$  receives from the challenger to an input to the “internal adversary”  $\mathcal{A}$ ?
  - ② **Input Distribution:** Does the input mapping provide the right distribution of inputs that  $\mathcal{A}$  expects?
  - ③ **Output Mapping:** How do we map the output that  $\mathcal{A}$  provides to an output for  $\mathcal{B}$ ?

# Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
  - ① **Input Mapping:** How to map the input that “outer adversary”  $\mathcal{B}$  receives from the challenger to an input to the “internal adversary”  $\mathcal{A}$ ?
  - ② **Input Distribution:** Does the input mapping provide the right distribution of inputs that  $\mathcal{A}$  expects?
  - ③ **Output Mapping:** How do we map the output that  $\mathcal{A}$  provides to an output for  $\mathcal{B}$ ?
  - ④ **Probability:** When we assume existence of  $\mathcal{A}$ , we also assume that  $\mathcal{A}$  wins with non-negligible advantage. What is the probability/advantage that  $\mathcal{B}$  wins, given the mappings above?


# One-Way Functions

# One Way Functions




- A function is one-way if it “easy to compute,” but “hard to invert”

# One Way Functions




- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)

# One Way Functions



- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).


# One Way Functions



- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).



# One Way Functions



- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).

*How to define one-way functions?*

# Defining One Way Functions: Attempt 1

**Attempt 1:** A function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$  is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm  $\mathcal{C}$  s.t.  
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

# Defining One Way Functions: Attempt 1

**Attempt 1:** A function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$  is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm  $\mathcal{C}$  s.t.  
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** for every non-uniform PPT adversary  $\mathcal{A}$ , for any input length  $n \in \mathbb{N}$

Probability of Inversion is Negligible

# Defining One Way Functions: Attempt 1

**Attempt 1:** A function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$  is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm  $\mathcal{C}$  s.t.  
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** for every non-uniform PPT adversary  $\mathcal{A}$ , for any input length  $n \in \mathbb{N}$

$$\Pr [\mathcal{A} \text{ inverts } f(x) \text{ for random } x] \leq \textit{negligible}.$$

This is called **average-case** hardness.

# One Way Functions: Definition

## Definition (One Way Function)

A function  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$  is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm  $\mathcal{C}$  s.t.  
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** there exists a negligible function  $\nu : \mathbb{N} \rightarrow \mathbb{R}$  s.t. for every non-uniform PPT adversary  $\mathcal{A}$  and  $\forall n \in \mathbb{N}$ :

$$\Pr \left[ x \stackrel{\$}{\leftarrow} \{0, 1\}^n, x' \leftarrow \mathcal{A}(1^n, f(x)) : f(x') = f(x) \right] \leq \nu(n).$$


- The above definition is also called **strong** one-way functions.

# One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.


# One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.



# One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.



We say that  $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$  is a one-way function if there exists a negligible function  $\nu : \mathbb{N} \rightarrow \mathbb{R}$  s.t. for every n.u. PPT adversary  $\mathcal{A}$  and  $\forall n \in \mathbb{N}$ :

$$\Pr[\mathcal{A} \text{ wins}] \leq \nu(n).$$



# Injective OWFs and One Way Permutations (OWP)

- **Injective or 1-1 OWFs:** each image has a unique pre-image:

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

- **One Way Permutations (OWP):** 1-1 OWF with the additional conditional that “each image has a pre-image”

(Equivalently: domain and range are of same size.)

# Existence of OWFs

Do OWFs exist?

# Existence of OWFs

Do OWFs exist?

- NOT Unconditionally — proving that  $f$  is one-way requires proving (at least)  $\mathbf{P} \neq \mathbf{NP}$ .

## Do OWFs exist?

- NOT Unconditionally — proving that  $f$  is one-way requires proving (at least)  $\mathbf{P} \neq \mathbf{NP}$ .
- However, we can construct them ASSUMING that certain problems are hard.

## Do OWFs exist?

- NOT Unconditionally — proving that  $f$  is one-way requires proving (at least)  $\mathbf{P} \neq \mathbf{NP}$ .
- However, we can construct them ASSUMING that certain problems are hard.
- Such constructions are sometimes called “candidates” because they are based on an assumption or a conjecture.

# Factoring Problem

- Consider the **multiplication** function  $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ :

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

# Factoring Problem

- Consider the **multiplication** function  $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ :

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.

# Factoring Problem

- Consider the **multiplication** function  $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ :

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is  $f_{\times}$  a OWF?



# Factoring Problem

- Consider the **multiplication** function  $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ :

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is  $f_{\times}$  a OWF?
- Clearly not!** With prob.  $1/2$ , a random number (of any fixed size) is even. I.e.,  $xy$  is even w/ prob.  $\frac{3}{4}$  for random  $(x, y)$ .

# Factoring Problem

- Consider the **multiplication** function  $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ :

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is  $f_{\times}$  a OWF?
- Clearly not!** With prob.  $1/2$ , a random number (of any fixed size) is even. I.e.,  $xy$  is even w/ prob.  $\frac{3}{4}$  for random  $(x, y)$ .
- Inversion: given number  $z$ , output  $(2, z/2)$  if  $z$  is even and  $(0, 0)$  otherwise! (succeeds 75% time)

# Factoring Problem (continued)

- Eliminate such trivial small factors.

# Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let  $\Pi_n$  be the set of all **prime** numbers  $< 2^n$ .

# Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let  $\Pi_n$  be the set of all **prime** numbers  $< 2^n$ .
- Choose numbers  $p$  and  $q$  randomly from  $\Pi_n$  and multiply.

## Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let  $\Pi_n$  be the set of all **prime** numbers  $< 2^n$ .
- Choose numbers  $p$  and  $q$  randomly from  $\Pi_n$  and multiply.
- This is unlikely to have small trivial factors.  
[Factoring Assumption] For every (non-uniform PPT) adversary  $\mathcal{A}$ , there exists a negligible function  $\nu$  such that

$$\Pr \left[ p \xleftarrow{\$} \Pi_n; q \xleftarrow{\$} \Pi_n; N = pq : \mathcal{A}(N) \in \{p, q\} \right] \leq \nu(n).$$

# Factoring Problem (continued)

- Factoring assumption is a well established conjecture.

# Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.



# Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.
- Best known algorithms for breaking Factoring Assumption:

$$2^{O(\sqrt{n \log n})} \quad (\text{provable})$$

$$2^{O(\sqrt[3]{n \log^2 n})} \quad (\text{heuristic})$$

# Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.
- Best known algorithms for breaking Factoring Assumption:

$$2^{O(\sqrt{n \log n})} \quad (\text{provable})$$

$$2^{O(\sqrt[3]{n \log^2 n})} \quad (\text{heuristic})$$

- Can we construct OWFs from the Factoring Assumption?

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen  $x$  and  $y$  happen to be primes, no PPT  $\mathcal{A}$  can invert (except with negligible probability). Call it the **GOOD** case.

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen  $x$  and  $y$  happen to be primes, no PPT  $\mathcal{A}$  can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability  $> \varepsilon$ ,  
 $\Rightarrow$  every PPT  $\mathcal{A}$  must fail to invert  $f_{\times}$  with probability at least  $\varepsilon$ .

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen  $x$  and  $y$  happen to be primes, no PPT  $\mathcal{A}$  can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability  $> \varepsilon$ ,  
 $\Rightarrow$  every PPT  $\mathcal{A}$  must fail to invert  $f_{\times}$  with probability at least  $\varepsilon$ .
- Now suppose that  $\varepsilon$  is a **noticeable function** (say e.g. an inverse polynomial, i.e.,  $\frac{1}{p(\cdot)}$ )  
 $\Rightarrow$  every  $\mathcal{A}$  must fail to invert  $f_{\times}$  with **noticeable** probability.

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen  $x$  and  $y$  happen to be primes, no PPT  $\mathcal{A}$  can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability  $> \varepsilon$ ,  
 $\Rightarrow$  every PPT  $\mathcal{A}$  must fail to invert  $f_{\times}$  with probability at least  $\varepsilon$ .
- Now suppose that  $\varepsilon$  is a **noticeable function** (say e.g. an inverse polynomial, i.e.,  $\frac{1}{p(\cdot)}$ )  
 $\Rightarrow$  every  $\mathcal{A}$  must fail to invert  $f_{\times}$  with **noticeable** probability.
- This is already useful!

# Multiplication Function

- Going back to the multiplication function  $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$ .

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen  $x$  and  $y$  happen to be primes, no PPT  $\mathcal{A}$  can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability  $> \varepsilon$ ,  
 $\Rightarrow$  every PPT  $\mathcal{A}$  must fail to invert  $f_{\times}$  with probability at least  $\varepsilon$ .
- Now suppose that  $\varepsilon$  is a **noticeable function** (say e.g. an inverse polynomial, i.e.,  $\frac{1}{p(\cdot)}$ )  
 $\Rightarrow$  every  $\mathcal{A}$  must fail to invert  $f_{\times}$  with **noticeable** probability.
- This is already useful!
- Usually called a **weak OWF**.



# Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

# Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

## Definition (Noticeable Function)

A function  $\nu(n)$  is noticeable if  $\exists c, n_0$  such that  $\forall n > n_0, \nu(n) \geq \frac{1}{n^c}$ .

# Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

## Definition (Noticeable Function)

A function  $\nu(n)$  is noticeable if  $\exists c, n_0$  such that  $\forall n > n_0, \nu(n) \geq \frac{1}{n^c}$ .

Note that a non-negligible function is not necessarily a noticeable function. Example:

$$f(n) = \begin{cases} 1 & \text{if } n \text{ is even} \\ 2^{-n} & \text{if } n \text{ is odd} \end{cases}.$$

This function is non-negligible, but not noticeable. **Why?**