

Master Research Report: Multilevel Multivariate Imputation by Chained Equations through Bayesian Additive Regression Trees

*Methodology and Statistics for the Behavioural, Biomedical and Social
Sciences*

Heleen Brügger



**Utrecht
University**

Word count:

Candidate Journal:

FETC Case Number:

Supervisors:

MSc. T. Volker

Dr. G. Vink

MSc. H. Oberman

2496

Computational Statistics & Data Analysis

23-1778

Utrecht University

Utrecht University

Utrecht University

1 Introduction

Incomplete data is a common challenge in many fields of research. Frequently used ad hoc strategies to deal with missing data, such as listwise deletion or mean imputation often lead to erroneous inferences in realistic situations. These strategies don't consider the multivariate nature of the data: the missingness can relate to observed values, which can lead to biased estimates and inaccurate variance estimates (Austin et al., 2021; Enders, 2017; Kang, 2013; van Buuren, 2018). Rubin defined three of such missing data mechanisms: Missing Completely At Random (MCAR) where the cause of the missing data is unrelated to the data, Missing At Random (MAR) where the missing data is related to the observed data, and Missing Not At Random (MNAR) where the missing data may also be related to unobserved data (Rubin, 1976).

Multiple imputation (MI; Rubin 1987) is considered a valid method for dealing with incomplete data, it allows us to separate the missing data problem from the analysis problem (Audigier et al., 2018; Austin et al., 2021; Burgette and Reiter, 2010; Enders, 2017; Grund et al., 2021; Hughes et al., 2014; Mistler and Enders, 2017; Van Buuren, 2007; van Buuren, 2018). MI is used to impute each missing value in the dataset more than once given the observed data, considering necessary variation associated with the missingness problem. The multiply imputed datasets are analyzed, and the corresponding inferences are pooled according to Rubin's rules (Austin et al., 2021; Carpenter and Kenward, 2013; Rubin, 1987; van Buuren, 2018). However, specifying the imputation models, the models used to impute the missing data, can be challenging. The concept of congeniality dictates that the imputation models should be at least as general as the analysis model and preferably all-encompassing (Bartlett et al., 2015; Enders et al., 2018a; Grund et al., 2016, 2018a; Meng, 1994). Otherwise, it will not capture every aspect of the data and the analysis model estimates may be biased. So, when the complexity of data increases, specifying the imputation models becomes more difficult (Grund et al., 2018a; van Buuren, 2018).

Congeniality-issues become more pronounced when MI is used in a multilevel data context (Audigier et al., 2018; Dong and Mitani, 2023; Enders et al., 2016, 2018a,b, 2020; Grund et al., 2016, 2018a,b, 2021; Lüdtke et al., 2017; Mistler and Enders, 2017; Quartagno and Carpenter, 2022; Resche-Rigon and White, 2018; Taljaard et al., 2008; van Buuren, 2018). Multilevel data is hierarchically structured, where, for example, students are nested within schools (Hox and Roberts, 2011; Hox et al., 2017). When analyzing multilevel data, the hierarchical structure should be considered. Ignoring the hierarchical structure will underestimate the intra-class correlation (ICC; Hox and Roberts 2011; Lüdtke et al. 2017; Taljaard et al. 2008; van Buuren 2018), which can be interpreted as the proportion of the total variance at level-2 (Gulliford et al., 2005; Hox and Roberts, 2011; Shieh, 2012). This can be done using multilevel models (MLM; Hox and Roberts 2011; Hox et al. 2017; Lüdtke et al. 2017). MLMs can contain both level-1, and level-2 variables, relating to the individual and class respectively, random intercepts, random slopes, and cross-level interactions (Hox and Roberts, 2011; Hox et al., 2017). Typically, the complexity of the multilevel analysis model is built step-wise with non-linearities, meaning the analysis model is not determined beforehand (Hox and Roberts, 2011; Hox et al., 2017). Thus, including the hierarchical structure, along with the complicated non-linearities from cross-level interactions in imputation models can be quite challenging (Burgette and Reiter, 2010; Hox and Roberts, 2011; van Buuren, 2018) and a very complex model might not converge (van Buuren, 2018).

A popular and flexible implementation of MI is fully conditional specification (FCS), otherwise known as chained equations (Audigier et al., 2018; Burgette and Reiter, 2010; Grund et al., 2018b; Van Buuren, 2007). FCS iteratively imputes each incomplete variable conditional on complete and previously imputed variables (Enders et al., 2016, 2018a,b; Grund et al., 2018b; Hughes et al., 2014; Mistler and Enders, 2017; van Buuren, 2018). In a multilevel context, FCS employs univariate linear mixed models to account for the hierarchical structure (Enders et al., 2018a; Mistler and Enders, 2017; Resche-Rigon and White, 2018). Furthermore, FCS can be used to impute non-linearities, such as cross-level interactions, by using 'passive imputation' or defining a separate imputation model for the non-linearities (Grund et al., 2018a; van Buuren, 2018). Still, imputation models including cross-level interaction or non-linear terms in FCS are very complicated (Grund et al., 2018a, 2021) and, thus, researchers' focus has predominantly been on the inclusion of random intercepts and slopes, but not of cross-level interactions (Enders et al., 2016, 2018a,b, 2020; Grund et al., 2016, 2018b).

Using non-parametric tree-based models might solve this problem because these models do not assume a specific data distribution and, thus, implicitly model non-linear relationships and interactions between the predictor variables, and handle continuous and categorical variables simultaneously (Breiman et al., 1984; Burgette and Reiter, 2010; Chipman et al., 2010; Hill et al., 2020; James et al., 2021; Lin and Luo, 2019; Salditt et al., 2023). In a single-level imputation context, the use of tree-based, non-parametric

models like regression trees, random forests, or Bayesian Additive Regression Trees (BART) simplified imputation models and performed better than parametric methods: the imputations showed better confidence interval coverage of the population parameters, lower variance and lower bias, especially in non-linear and interactive contexts (Burgette and Reiter, 2010; Silva and Gutman, 2022; Xu et al., 2016). Waljee et al. (2013) also found lower imputation error when imputing with a random forest algorithm compared to multivariate imputation by chained equations (MICE), K-nearest neighbors (KNN) and mean imputation.

BART models have been implemented in a multilevel prediction context. However, multilevel-BART models (M-BART) have predominantly been implemented with only random intercepts (Chen, 2020; Tan et al., 2016; Wagner et al., 2020; Wundervald et al., 2022). In a prediction context, Wagner et al. (2020) have found that this random intercept M-BART model provided better predictions with a lower mean squared error (MSE) compared to a parametric MLM, Tan et al. (2016) found higher area under the curve (AUC) values compared to a single-level BART model and linear random intercept model, and Chen (2020) found better predictions and better coverage of the estimates compared to parametric models and a single-level BART model. Other researchers modeled the random intercept as an extra split on each terminal node and found a lower MSE compared to a standard BART model and parametric MLMs (Wundervald et al., 2022). Dorie et al. (2022) developed a multilevel BART model that included random intercepts and random slopes by modeling the random parts with Stan (Lee et al., 2017) and the fixed parts with BART. Their results showed that their algorithm `stan4bart` showed better coverage of the population values and lower root mean squared error (RMSE) compared to BART models with varying intercept, BART models ignoring the multilevel structure, bayesian causal forests, and parametric MLMs.

Despite these promising findings, M-BART models have yet to be implemented in a multilevel multiple imputation context. Thus, my thesis research question will be: *Can multivariate imputation by chained equations through a multilevel bayesian additive regression trees model improve the bias, variance, and coverage of the estimates in a multilevel context compared to current practices?* Given the success of non-parametric models in single-level MI, I anticipate that employing M-BART models in a multilevel missing data context will reduce bias, accurately model variance, and improve estimate coverage compared to conventional implementations of multilevel MI, single-level MI, and listwise deletion in the R-package MICE (Buuren and Groothuis-Oudshoorn, 2011). However, in this research report, I will only focus on the implementation of M-BART models in a prediction context and asses their performance in terms of relative bias and MSE. The research question is: *Can M-BART models improve the relative bias and MSE of the predictions in a multilevel context compared to a single-level BART model?*

The research report’s sections will cover theoretical background, methods for evaluating M-BART models, preliminary results, and discussion of next steps.

2 Method

2.1 Theoretical background

2.1.1 Bayesian Additive Regression Trees (BART)

BART is a sum-of-trees model proposed by Chipman et al. (2010) that has regression trees as its building blocks (Chipman et al., 2010; Hill et al., 2020; James et al., 2021). Regression trees divide the data into subgroups by recursively splitting the data into binary subgroups based on the predictors minimizing variability within the subgroups (Hastie, 2017; James et al., 2021; Salditt et al., 2023). Recursive binary partitioning of the predictor space doesn’t assume a specific data form, making this a non-parametric model (Hastie, 2017; James et al., 2021; Salditt et al., 2023) and allows regression trees to model non-linearities well and automatically (Burgette and Reiter, 2010; Hill et al., 2020). Chipman et al. (2010) define the BART model as:

$$f(\mathbf{x}) = \sum_{k=1}^m g(\mathbf{x}; T_k, M_k), \quad (1)$$

where $f(\mathbf{x})$ is the overall fit of the model: the sum of m regression trees, \mathbf{x} are the predictor variables, T_k is the k^{th} tree and M_k is the collection of leaf parameters within the k^{th} tree (Chipman et al., 2010; Hill et al., 2020; James et al., 2021). The data are assumed to arise from a model with additive normally distributed errors: $Y = \sum_{k=1}^m g(\mathbf{x}; T_k, M_k) + \epsilon, \epsilon \sim \mathcal{N}(0, \sigma^2)$. Next to the sum-of-trees model, BART also includes a regularization prior that constrains the size and fit of each tree so that each contributes only a

small part to prevent overfitting (Chipman et al., 2010; Hill et al., 2020; James et al., 2021). BARTs are estimated using the Bayesian back-fitting Markov Chain Monte Carlo (MCMC) algorithm. It updates individual trees, considering the remaining trees, their associated parameters, and the residual standard deviation (σ). It fits a new tree to the partial residuals, r_i , treating them as the data, by perturbing the tree from the previous iteration. Perturbations entail either *growing*, *pruning*, or *changing* a tree. *Growing* means adding additional splits, *pruning* removes splits, and *changing* changes decision rules. The algorithm stops after the specified number of iterations. The partial residuals are defined as:

$$r_i = y_i - \sum_{k' < k} \hat{f}_{k'}^b(x_i) - \sum_{k' > k} \hat{f}_{k'}^{b-1}(x_i), \text{ with } i = 1, \dots, N \quad (2)$$

where $\hat{f}_k^b(x_i)$ is the prediction of the k^{th} tree in the b^{th} iteration for person i and sample size N .

2.1.2 Multilevel-BART (M-BART)

Chen (2020); Wagner et al. (2020) and Tan et al. (2016) define a M-BART model including a random intercept building on the work of Lin and Luo (2019). The M-BART algorithm breaks down the observed variable into fixed and random components. The fixed components are modeled by BART and the random components are modeled by a linear mixed effects model (Chen, 2020; Tan et al., 2016; Wagner et al., 2020). The BART model (1) can be extended to include a random intercept by:

$$f(\mathbf{x}) = \sum_{k=1}^m g(\mathbf{x}; T_k, M_k) + \alpha_j, \quad (3)$$

where, now, $f(\mathbf{x})$ is the overall fit of the model incorporating random intercept α_j for cluster j .

2.2 Simulation study

2.2.1 Data generating mechanism

The population data-generating mechanism will be based on the following MLM:

$$y_{ij} = \beta_{0j} + \sum_{k=1}^7 \beta_{kj} X_{kij} + \epsilon_{ij}, \quad X_{kij} \sim \mathcal{MVN}(0, \mathbf{\Sigma}_x), \quad \epsilon_{ij} \sim \mathcal{N}(0, 25), \quad (4a)$$

$$\beta_{0j} = \gamma_{00} + \sum_{q=1}^2 \gamma_{0q} Z_{qj} + v_{0j}, \quad (4b)$$

$$\beta_{kj} = \gamma_{k0} + \sum_{q=1}^2 \gamma_{kq} Z_{qj} + v_{kj}, \quad Z_{qj} \sim \mathcal{MVN}(0, \mathbf{\Sigma}_z), \quad \mathbf{v}_j \sim \mathcal{MVN}(0, \mathbf{T}), \quad (4c)$$

$$\gamma_{00} = 10, \gamma_{0q} = \begin{pmatrix} .5 \\ .5 \end{pmatrix}, \gamma_{k0} = \begin{pmatrix} \gamma_{10} \\ \gamma_{20} \\ \gamma_{30} \\ \gamma_{40} \\ \gamma_{50} \\ \gamma_{60} \\ \gamma_{70} \end{pmatrix}, \gamma_{kq} = \begin{pmatrix} .35 & 0 \\ .35 & 0 \\ 0 & .35 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad (4d)$$

$$\mathbf{\Sigma}_x = \begin{pmatrix} 6.25 & & & & & & \\ 2.25 & 9 & & & & & \\ 1.5 & 1.8 & 4 & & & & \\ 2.25 & 3.06 & 2.04 & 11.56 & & & \\ 1.5 & 1.8 & 1.2 & 2.04 & 4 & & \\ 1.125 & 1.35 & 0.9 & 1.53 & .9 & 2.25 & \\ 3.3 & 3.96 & 2.64 & 4.488 & 2.64 & 1.98 & 19.36 \end{pmatrix}, \mathbf{\Sigma}_z = \begin{pmatrix} 1 & \\ .48 & 2.56 \end{pmatrix}, \quad (4e)$$

$$\mathbf{T} = \begin{pmatrix} t_{00} & & & & & & \\ .3 & 1 & & & & & \\ .3 & .3 & 1 & & & & \\ .3 & .3 & .3 & 1 & & & \\ .3 & .3 & .3 & .3 & 1 & & \\ .3 & .3 & .3 & .3 & .3 & 1 & \\ .3 & .3 & .3 & .3 & .3 & .3 & 1 \end{pmatrix}, \quad (4f)$$

where y_{ij} is a continuous level-1 outcome variable for person i in group j and X_{kij} are k continuous level-1 variables and Z_{qj} are q continuous level-2 variables. They are multivariate normally distributed with means of 0 and variance-covariance matrix $\mathbf{\Sigma}_x$ and $\mathbf{\Sigma}_z$ respectively. Equation 4e shows the variance-covariance matrices of the level-1 and level-2 variables. The covariances between the variables were calculated as such that the correlation between the variables was .3, aligned with Cohen's (1990) medium effect size benchmark. The residuals are normally distributed with a mean of 0 and a variance of 25. The random intercept β_{0j} is determined by the overall intercept γ_{00} , the q group-level effects $\gamma_{0q}Z_{qj}$ and the group-level random residuals v_{0j} . The overall intercept γ_{00} was set to 10 and the group-level effects γ_{01} and γ_{02} to .5. The k regression coefficients β_{kj} for the continuous variables X_{kij} depend on the intercepts γ_{k0} , the cross-level interactions $\gamma_{kq}Z_{qj}$, and the random slopes v_{kj} . The k intercepts, or within-group effect sizes, γ_{k0} are varied in the simulations, the cross-level interactions γ_{11} , γ_{21} , and γ_{32} were set to .35. The random slopes are multivariate normally distributed with a mean of 0 and a variance-covariance matrix \mathbf{T} shown in equation 4f. Again, the covariances were calculated to yield a correlation of .3. The variance of v_{0j} , the group-level random residuals t_{00} , were scaled such that the specified ICC values as in table 1 was obtained. The following formula was used to calculate v_{0j} following the variance decomposition from Rights and Sterba (2019):

$$\text{ICC} = \frac{\gamma^{b'} \phi^b \gamma^b + \tau_{00}}{\gamma^{w'} \phi^w \gamma^w + \gamma^{b'} \phi^b \gamma^b + \text{tr}(\mathbf{T}\mathbf{\Sigma}) + \tau_{00} + \sigma^2}, \quad (5)$$

where γ^b and γ^w are the level-1 and level-2 fixed effects, ϕ^b and ϕ^w are the variance-covariance matrices of a vector with 1, for the intercept, and all level-2 predictors and all cluster-mean-centered level-1 predictors respectively, τ_{00} is the variance of the random intercept, \mathbf{T} is the variance-covariance matrix of the random intercept and slopes, Σ is the variance-covariance matrix of a vector containing 1, for the intercept, and the level-1 variables, and σ^2 is the residual variance. The value for τ_{00} was calculated using the function `uniroot` in R (R Core Team, 2023).

2.2.2 Simulation design

Table 1 shows the variations considered in the simulation study. They are realistic in practice and/or previously proposed (Enders et al., 2018b, 2020; Grund et al., 2018a; Gulliford et al., 1999; Hox et al., 2017; Murray and Blitstein, 2003). For each combination of varying parameters, 1000 datasets are simulated. 6 different imputation methods are compared:

1. listwise deletion,
2. conventional single-level imputation,
3. conventional multilevel imputation,
4. single-level BART imputation,
5. multilevel BART imputation accounting for random intercepts (Chen, 2020; Tan et al., 2016; Wagner et al., 2020; Wundervald et al., 2022),
6. multilevel BART imputation accounting for random effects and cross-level interactions (Dorie et al., 2022).

Table 1: Simulation design

Parameter	Values
Number of clusters (j)	30, 50
Within-cluster sample size (n_j)	5, 15, 35, 50
Intraclass Correlation (ICC)	0, .05, .3, .5
Missing data mechanism	MAR, MCAR
Amount of missingness	0%, 25%, 50%
Within-group effect size (γ)	.2, .5, .8

The second and third methods will be implemented with the R-package MICE (Buuren and Groothuis-Oudshoorn, 2011).

The fourth, single-level BART, fifth, random intercept BART and sixth method, multilevel BART methods will be implemented by writing functions in R (R Core Team, 2023) for the package MICE (Buuren and Groothuis-Oudshoorn, 2011) using the R-packages `dbarts` (Dorie, 2023a) and `stan4bart` (Dorie, 2023b). The function `bart` from the `dbarts` package is used to

2.2.3 Missing data generation

2.2.4 Evaluation

The fitted models are evaluated in terms of relative bias and Mean Squared Error (MSE) of the predictions (Morris et al., 2019):

$$\text{Bias} = \frac{1}{n_{\text{sim}}} \sum_{t=1}^{n_{\text{sim}}} (\hat{\theta}_t - \theta), \quad (6)$$

$$\text{MSE} = \frac{1}{n_{\text{sim}}} \sum_{t=1}^{n_{\text{sim}}} (\hat{\theta}_t - \theta)^2, \quad (7)$$

where $\hat{\theta}_t$ is the estimated parameter in simulation t , θ is the true value, and n_{sim} is the number of simulated datasets and smaller is better.

3 Results

4 Discussion

References

- Vincent Audigier, Ian R. White, Shahab Jolani, Thomas P. A. Debray, Matteo Quartagno, James Carpenter, Stef Van Buuren, and Matthieu Resche-Rigon. Multiple Imputation for Multilevel Data with Continuous and Binary Variables. *Statistical Science*, 33(2), May 2018. ISSN 0883-4237. doi: 10.1214/18-STS646.
- Peter C. Austin, Ian R. White, Douglas S. Lee, and Stef Van Buuren. Missing Data in Clinical Research: A Tutorial on Multiple Imputation. *Canadian Journal of Cardiology*, 37(9):1322–1331, September 2021. ISSN 0828282X. doi: 10.1016/j.cjca.2020.11.010.
- Jonathan W Bartlett, Shaun R Seaman, Ian R White, James R Carpenter, and for the Alzheimer’s Disease Neuroimaging Initiative*. Multiple imputation of covariates by fully conditional specification: Accommodating the substantive model. *Statistical Methods in Medical Research*, 24(4):462–487, August 2015. ISSN 0962-2802, 1477-0334. doi: 10.1177/0962280214521348.
- Leo Breiman, Jerome H. Friedman, Richard A. Olshen, and Charles J. Stone. *Classification And Regression Trees*. Routledge, 1 edition, 1984. ISBN 978-1-315-13947-0. doi: 10.1201/9781315139470.
- L. F. Burgette and J. P. Reiter. Multiple Imputation for Missing Data via Sequential Regression Trees. *American Journal of Epidemiology*, 172(9):1070–1076, November 2010. ISSN 0002-9262, 1476-6256. doi: 10.1093/aje/kwq260.
- Stef Van Buuren and Karin Groothuis-Oudshoorn. **Mice** : Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software*, 45(3), 2011. ISSN 1548-7660. doi: 10.18637/jss.v045.i03.
- James R. Carpenter and Michael G. Kenward. *Multiple Imputation and Its Application*. Wiley, 1 edition, January 2013. ISBN 978-0-470-74052-1 978-1-119-94228-3. doi: 10.1002/9781119942283.
- Siqi Chen. *A New Multilevel Bayesian Nonparametric Algorithm and Its Application in Causal Inference*. PhD thesis, Texas A&M University., October 2020.
- Hugh A. Chipman, Edward I. George, and Robert E. McCulloch. BART: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4(1), March 2010. ISSN 1932-6157. doi: 10.1214/09-AOAS285.
- Jacob Cohen. Statistical power analysis for the behavioral sciences. *Computers, Environment and Urban Systems*, 14(1):71, January 1990. ISSN 01989715. doi: 10.1016/0198-9715(90)90050-4.
- Mei Dong and Aya Mitani. Multiple imputation methods for missing multilevel ordinal outcomes. *BMC Medical Research Methodology*, 23(1):112, May 2023. ISSN 1471-2288. doi: 10.1186/s12874-023-01909-5.
- Vincent Dorie. Dbarts: Discrete bayesian additive regression trees sampler, 2023a.
- Vincent Dorie. *Stan4bart: Bayesian Additive Regression Trees with Stan-Sampled Parametric Extensions*, 2023b.
- Vincent Dorie, George Perrett, Jennifer L. Hill, and Benjamin Goodrich. Stan and BART for Causal Inference: Estimating Heterogeneous Treatment Effects Using the Power of Stan and the Flexibility of Machine Learning. *Entropy*, 24(12):1782, December 2022. ISSN 1099-4300. doi: 10.3390/e24121782.
- Craig K. Enders. Multiple imputation as a flexible tool for missing data handling in clinical research. *Behaviour Research and Therapy*, 98:4–18, November 2017. ISSN 00057967. doi: 10.1016/j.brat.2016.11.008.
- Craig K. Enders, Stephen A. Mistler, and Brian T. Keller. Multilevel multiple imputation: A review and evaluation of joint modeling and chained equations imputation. *Psychological Methods*, 21(2):222–240, June 2016. ISSN 1939-1463, 1082-989X. doi: 10.1037/met0000063.
- Craig K. Enders, Timothy Hayes, and Han Du. A Comparison of Multilevel Imputation Schemes for Random Coefficient Models: Fully Conditional Specification and Joint Model Imputation with Random Covariance Matrices. *Multivariate Behavioral Research*, 53(5):695–713, September 2018a. ISSN 0027-3171, 1532-7906. doi: 10.1080/00273171.2018.1477040.

- Craig K. Enders, Brian T. Keller, and Roy Levy. A fully conditional specification approach to multilevel imputation of categorical and continuous variables. *Psychological Methods*, 23(2):298–317, June 2018b. ISSN 1939-1463, 1082-989X. doi: 10.1037/met0000148.
- Craig K. Enders, Han Du, and Brian T. Keller. A model-based imputation procedure for multilevel regression models with random coefficients, interaction effects, and nonlinear terms. *Psychological Methods*, 25(1):88–112, February 2020. ISSN 1939-1463, 1082-989X. doi: 10.1037/met0000228.
- Simon Grund, Oliver Lüdtke, and Alexander Robitzsch. Multiple imputation of missing covariate values in multilevel models with random slopes: A cautionary note. *Behavior Research Methods*, 48(2):640–649, June 2016. ISSN 1554-3528. doi: 10.3758/s13428-015-0590-3.
- Simon Grund, Oliver Lüdtke, and Alexander Robitzsch. Multiple Imputation of Missing Data for Multilevel Models: Simulations and Recommendations. *Organizational Research Methods*, 21(1):111–149, January 2018a. ISSN 1094-4281, 1552-7425. doi: 10.1177/1094428117703686.
- Simon Grund, Oliver Lüdtke, and Alexander Robitzsch. Multiple Imputation of Missing Data at Level 2: A Comparison of Fully Conditional and Joint Modeling in Multilevel Designs. *Journal of Educational and Behavioral Statistics*, 43(3):316–353, June 2018b. ISSN 1076-9986, 1935-1054. doi: 10.3102/1076998617738087.
- Simon Grund, Oliver Lüdtke, and Alexander Robitzsch. Multiple imputation of missing data in multilevel models with the R package mdmb: A flexible sequential modeling approach. *Behavior Research Methods*, 53(6):2631–2649, May 2021. ISSN 1554-3528. doi: 10.3758/s13428-020-01530-0.
- M. C. Gulliford, O. C. Ukoumunne, and S. Chinn. Components of Variance and Intraclass Correlations for the Design of Community-based Surveys and Intervention Studies: Data from the Health Survey for England 1994. *American Journal of Epidemiology*, 149(9):876–883, May 1999. ISSN 0002-9262, 1476-6256. doi: 10.1093/oxfordjournals.aje.a009904.
- M.C. Gulliford, G. Adams, O.C. Ukoumunne, R. Latinovic, S. Chinn, and M.J. Campbell. Intraclass correlation coefficient and outcome prevalence are associated in clustered binary data. *Journal of Clinical Epidemiology*, 58(3):246–251, March 2005. ISSN 08954356. doi: 10.1016/j.jclinepi.2004.08.012.
- T. J. Hastie, editor. *Statistical Models in S*. Routledge, 1st edition, 2017. ISBN 978-1-351-41422-7.
- Jennifer Hill, Antonio Linero, and Jared Murray. Bayesian Additive Regression Trees: A Review and Look Forward. *Annual Review of Statistics and Its Application*, 7(1):251–278, March 2020. ISSN 2326-8298, 2326-831X. doi: 10.1146/annurev-statistics-031219-041110.
- Joop Hox and J. Kyle Roberts, editors. *Handbook of Advanced Multilevel Analysis*. Routledge, 0 edition, January 2011. ISBN 978-1-136-95127-5. doi: 10.4324/9780203848852.
- Joop J. Hox, Mirjam Moerbeek, and Rens Van De Schoot. *Multilevel Analysis: Techniques and Applications*. Routledge, Third edition. | New York, NY : Routledge, 2017. |, 3 edition, September 2017. ISBN 978-1-315-65098-2. doi: 10.4324/9781315650982.
- Rachael A Hughes, Ian R White, Shaun R Seaman, James R Carpenter, Kate Tilling, and Jonathan Ac Sterne. Joint modelling rationale for chained equations. *BMC Medical Research Methodology*, 14(1):28, December 2014. ISSN 1471-2288. doi: 10.1186/1471-2288-14-28.
- Gareth James, Daniela Witten, Trevor Hastie, and Robert Tibshirani. *An Introduction to Statistical Learning: With Applications in R*. Springer Texts in Statistics. Springer US, New York, NY, 2021. ISBN 978-1-07-161417-4 978-1-07-161418-1. doi: 10.1007/978-1-0716-1418-1.
- Hyun Kang. The prevention and handling of the missing data. *Korean Journal of Anesthesiology*, 64(5): 402, 2013. ISSN 2005-6419, 2005-7563. doi: 10.4097/kjae.2013.64.5.402.
- Daniel Lee, Bob Carpenter, Peter Li, Mitzi Morris, Michael Betancourt, Maverickg, Marcus Brubaker, Rob Trangucci, Marco Inacio, Alp Kucukelbir, Stan Buildbot, Bgoodri, Seantalts, Jeffrey Arnold, Dustin Tran, Matt Hoffman, Charles Margossian, Martin Modrák, Avraham Adler, Krzysztof Sakrejda, Alexey Stukalov, Mike Lawrence, Rob J Goedman, Kevin S. Van Horn, Aki Vehtari, Jonah Gabry, Juan Sebastián Casallas, and Ben Bales. Stan-dev/stan: V2.17.1. Zenodo, December 2017.

- Shuqiong Lin and Wen Luo. A New Multilevel CART Algorithm for Multilevel Data with Binary Outcomes. *Multivariate Behavioral Research*, 54(4):578–592, July 2019. ISSN 0027-3171, 1532-7906. doi: 10.1080/00273171.2018.1552555.
- Oliver Lüdtke, Alexander Robitzsch, and Simon Grund. Multiple imputation of missing data in multilevel designs: A comparison of different strategies. *Psychological Methods*, 22(1):141–165, March 2017. ISSN 1939-1463, 1082-989X. doi: 10.1037/met0000096.
- Xiao-Li Meng. Multiple-imputation inferences with uncongenial sources of input. *Statistical science*, pages 538–558, 1994.
- Stephen A. Mistler and Craig K. Enders. A Comparison of Joint Model and Fully Conditional Specification Imputation for Multilevel Missing Data. *Journal of Educational and Behavioral Statistics*, 42(4):432–466, August 2017. ISSN 1076-9986, 1935-1054. doi: 10.3102/1076998617690869.
- Tim P. Morris, Ian R. White, and Michael J. Crowther. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11):2074–2102, May 2019. ISSN 0277-6715, 1097-0258. doi: 10.1002/sim.8086.
- David M. Murray and Jonathan L. Blitstein. Methods To Reduce The Impact Of Intraclass Correlation In Group-Randomized Trials. *Evaluation Review*, 27(1):79–103, February 2003. ISSN 0193-841X, 1552-3926. doi: 10.1177/0193841X02239019.
- Matteo Quartagno and James R. Carpenter. Substantive model compatible multilevel multiple imputation: A joint modeling approach. *Statistics in Medicine*, 41(25):5000–5015, November 2022. ISSN 0277-6715, 1097-0258. doi: 10.1002/sim.9549.
- R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria, 2023.
- Matthieu Resche-Rigon and Ian R White. Multiple imputation by chained equations for systematically and sporadically missing multilevel data. *Statistical Methods in Medical Research*, 27(6):1634–1649, June 2018. ISSN 0962-2802, 1477-0334. doi: 10.1177/0962280216666564.
- Jason D. Rights and Sonya K. Sterba. Quantifying explained variance in multilevel models: An integrative framework for defining R-squared measures. *Psychological Methods*, 24(3):309–338, June 2019. ISSN 1939-1463, 1082-989X. doi: 10.1037/met0000184.
- Donald B. Rubin. Inference and missing data. *Biometrika*, 63(3):581–592, 1976. ISSN 0006-3444, 1464-3510. doi: 10.1093/biomet/63.3.581.
- Donald B. Rubin. *Multiple Imputation for Nonresponse in Surveys*. Wiley, New York, 1987. ISBN 978-0-470-31669-6.
- Marie Salditt, Sarah Humberg, and Steffen Nestler. Gradient Tree Boosting for Hierarchical Data. *Multivariate Behavioral Research*, pages 1–27, January 2023. ISSN 0027-3171, 1532-7906. doi: 10.1080/00273171.2022.2146638.
- Gwonen Shieh. A comparison of two indices for the intraclass correlation coefficient. *Behavior Research Methods*, 44(4):1212–1223, December 2012. ISSN 1554-3528. doi: 10.3758/s13428-012-0188-y.
- Gabriella C. Silva and Roege Gutman. Multiple imputation procedures for estimating causal effects with multiple treatments with application to the comparison of healthcare providers. *Statistics in Medicine*, 41(1):208–226, January 2022. ISSN 0277-6715, 1097-0258. doi: 10.1002/sim.9231.
- Monica Taljaard, Allan Donner, and Neil Klar. Imputation Strategies for Missing Continuous Outcomes in Cluster Randomized Trials. *Biometrical Journal*, 50(3):329–345, June 2008. ISSN 0323-3847, 1521-4036. doi: 10.1002/bimj.200710423.
- Yaoyuan Vincent Tan, Carol A. C. Flannagan, and Michael R. Elliott. Predicting human-driving behavior to help driverless vehicles drive: Random intercept Bayesian Additive Regression Trees. 2016. doi: 10.48550/ARXIV.1609.07464.
- Stef Van Buuren. Multiple imputation of discrete and continuous data by fully conditional specification. *Statistical Methods in Medical Research*, 16(3):219–242, June 2007. ISSN 0962-2802, 1477-0334. doi: 10.1177/0962280206074463.

- Stef van Buuren. *Flexible Imputation of Missing Data*. Chapman & Hall/CRC Interdisciplinary Statistics Series. CRC Press, Taylor & Francis Group, Boca Raton London New York, second edition edition, 2018. ISBN 978-1-138-58831-8.
- James Wagner, Brady T. West, Michael R. Elliott, and Stephanie Coffey. Comparing the Ability of Regression Modeling and Bayesian Additive Regression Trees to Predict Costs in a Responsive Survey Design Context. *Journal of Official Statistics*, 36(4):907–931, December 2020. ISSN 2001-7367. doi: 10.2478/jos-2020-0043.
- Akbar K Waljee, Ashin Mukherjee, Amit G Singal, Yiwei Zhang, Jeffrey Warren, Ulysses Balis, Jorge Marrero, Ji Zhu, and Peter Dr Higgins. Comparison of imputation methods for missing laboratory data in medicine. *BMJ Open*, 3(8):e002847, August 2013. ISSN 2044-6055, 2044-6055. doi: 10.1136/bmjopen-2013-002847.
- Bruna Wundervald, Andrew Parnell, and Katarina Domijan. Hierarchical Embedded Bayesian Additive Regression Trees. 2022. doi: 10.48550/ARXIV.2204.07207.
- Dandan Xu, Michael J. Daniels, and Almut G. Winterstein. Sequential BART for imputation of missing covariates. *Biostatistics*, 17(3):589–602, July 2016. ISSN 1465-4644, 1468-4357. doi: 10.1093/biostatistics/kxw009.