

Master Research Report: Multilevel Multivariate Imputation by Chained Equations through Bayesian Additive Regression Trees

*Methodology and Statistics for the Behavioural, Biomedical and Social
Sciences*

Heleen Brügger

Word count:

Candidate Journal:

FETC Case Number:

Supervisors:

MSc. T. Volker

Dr. G. Vink

MSc. H. Oberman

3270

Computational Statistics & Data Analysis

23-1778

Utrecht University

Utrecht University

Utrecht University

1 Introduction

Incomplete data is a common challenge in many fields of research. Frequently, ad hoc strategies are used to deal with missing data, such as listwise deletion, pairwise deletion, mean imputation and regression imputation. However, these strategies typically lead to erroneous inferences in a realistic situation. The estimated may be biased and variances in the data are no longer correct [Austin et al., 2021, Enders, 2017, Kang, 2013, van Buuren, 2018]. Rubin defined three types of missing data mechanisms: Missing Completely At Random (MCAR) where the cause of the missing data is unrelated to the missing data, Missing At Random (MAR) where the missing data is related to the observed data, and Missing Not At Random (MNAR) where the missing data is related to unobserved data [Rubin, 1976].

Multiple imputation (MI) is considered a valid method for dealing with incomplete data and allows us to separate the missing data problem from the analysis problem [Audigier et al., 2018, Austin et al., 2021, Burgette and Reiter, 2010, Enders, 2017, Grund et al., 2021, Hughes et al., 2014, Mistler and Enders, 2017, Van Buuren, 2007, van Buuren, 2018]. MI is used to impute each missing value in the data set more than once given the observed data, thereby considering necessary variation associated with the missingness problem. The multiply imputed data sets are analyzed, and the corresponding inferences are pooled according to Rubin’s rules [Austin et al., 2021, Carpenter and Kenward, 2013, Rubin, 1987, van Buuren, 2018]. The models used to impute the missing data, the imputation models, should be at least as general as the analysis model and preferably all-encompassing [Bartlett et al., 2015, Enders et al., 2018a, Grund et al., 2016, 2018b, Meng, 1994]. Otherwise, the imputation model will not capture every aspect of the data and the completed data sets cannot be properly analysed. So, when the complexity of data increases, specifying the imputation models becomes more difficult [Grund et al., 2018b, van Buuren, 2018].

MI has been implemented in a multilevel data context [Audigier et al., 2018, Dong and Mitani, 2023, Enders et al., 2020, 2018a,b, 2016, Grund et al., 2016, 2018a,b, 2021, Lüdtke et al., 2017, Mistler and Enders, 2017, Quartagno and Carpenter, 2022, Resche-Rigon and White, 2018, Taljaard et al., 2008, van Buuren, 2018], where this problem of specifying imputation models as general as the analysis model becomes magnified. Multilevel data is hierarchically structured, where, for example, students are nested within schools, or patients are nested within hospitals [Hox and Roberts, 2011, Hox et al., 2017]. Thus, in these types of data sets there are level-1 and level-2 variables. Level-1 variables relate to the individual within a class and level-2 variables relate to the class as a whole. The recommended statistical technique to analyzing these models are multilevel models as it accounts for the specific dependencies in the multilevel data sets [Hox and Roberts, 2011, Hox et al., 2017, Lüdtke et al., 2017]. It can contain both level-1 and level-2 variables, random intercepts, random slopes, and cross-level interactions [Hox and Roberts, 2011, Hox et al., 2017]. Furthermore, the complexity of the multilevel analysis model is built step-wise with non-linearities [Hox and Roberts, 2011, Hox et al., 2017]. Thus, when defining imputation models for a multilevel data set one needs to account for the hierarchical structure, as well as the complicated non-linearities from cross-level interactions. Since ignoring the multilevel structure and imputing the data as if it were a single-level data set will underestimate the Intra-Class Correlation (ICC) [Hox and Roberts, 2011, Lüdtke et al., 2017, Taljaard et al., 2008, van Buuren, 2018]. The ICC can be interpreted as the expected correlation between two randomly sampled individuals from the same group or the proportion of the total variance at level-2 [Gulliford et al., 2005, Hox and Roberts, 2011, Shieh, 2012]. So, defining these models is quite challenging [Burgette and Reiter, 2010, Hox and Roberts, 2011, van Buuren, 2018].

Generally, multiple imputation operates under two frameworks: joint modeling (JM) and fully conditional specification (FCS) [Enders et al., 2018a,b, Hughes et al., 2014, Mistler and Enders, 2017, van Buuren, 2018]. JM employs a multivariate data distribution and regresses incomplete variables on all complete variables to impute missing values. FCS, or chained equations, iteratively imputes one variable with missing values at a time through conditional univariate distributions regressing an incomplete variable on complete variables and previously imputed variables [Enders et al., 2018a,b, 2016, Grund et al., 2018a, Hughes et al., 2014, Mistler and Enders, 2017, van Buuren, 2018]. JM and FCS use multivariate and univariate linear mixed models respectively in a multilevel context [Enders et al., 2018a, Mistler and Enders, 2017, Resche-Rigon and White, 2018], accounting for the hierarchical structure. Thus far, much focus has been on multilevel JM and FCS including random intercepts and slopes, but no cross-level interactions Enders et al. [2020, 2018a,b, 2016], Grund et al. [2016, 2018a]. Overall, FCS is believed to be more flexible than JM [Audigier et al., 2018, Burgette and Reiter, 2010, Grund et al., 2018a, Van Buuren, 2007] and, thus, may be better suited for multilevel data. Grund et al. [2018b] recommend using FCS to impute a multilevel data set with random intercepts, slopes and cross-level interactions or non-linear terms. FCS uses ‘passive imputation’ or treats interactions as ‘just another variable’ by defining an

imputation model for interactions. On the other hand, JM does not account for cross-level interactions at all [Grund et al., 2018b, van Buuren, 2018]. Still, the imputation model of multilevel models including cross-level interaction or non-linear terms in FCS is still a very complicated process [Grund et al., 2018b, 2021].

Using non-parametric tree-based models as imputation models might solve this problem. In a single-level imputation context, the use of tree-based, non-parametric models like regression trees, random forests or Bayesian Additive Regression Trees (BART) simplified imputation models and performed better than parametric methods: the imputations showed better confidence interval coverage of the population parameters, lower variance and lower bias, especially in non-linear and interactive contexts [Burgette and Reiter, 2010, Silva and Gutman, 2022, Xu et al., 2016]. Waljee et al. [2013] also found lower imputation error when imputing with a random forest algorithm compared to MICE, KNN and mean imputation and Stekhoven and Bühlmann [2012] found that the algorithm reduced computational time and could handle multivariate data consisting of both continuous and categorical data simultaneously.

Tree-based models use recursive partitioning to split the data into smaller subgroups based on the predictor variables maximizing the homogeneity of the subgroups. Tree-based models are non-parametric, which means that they do not assume a specific distribution of the data. Thus, they can handle non-linear relationships and interactions between the predictor variables well. Furthermore they handle continuous and categorical variables simultaneously [Breiman et al., 1984, Burgette and Reiter, 2010, Chipman et al., 2010, Hill et al., 2020, James et al., 2021, Lin and Luo, 2019, Salditt et al., 2023].

BART models have also been implemented in a multilevel prediction context. However, multilevel-BART models (M-BART) have predominantly been implemented with random intercepts and no random slopes and cross-level interactions [Chen, 2020, Tan et al., 2016, Wagner et al., 2020, Wundervald et al., 2022]. Wagner et al. [2020] have found that this random intercept M-BART model provided better predictions with a lower Mean Squared Error (MSE) compared to a parametric multilevel model, Tan et al. [2016] found higher Area Under the Curve values, and Chen [2020] found better predictions and better coverage compared to parametric models and a single-level BART model. Other researchers modeled the random intercept as an extra split on each terminal node within the BART algorithm and found a lower MSE compared to a standard BART model and parametric multilevel models [Wundervald et al., 2022]. Dorie et al. [2022] developed a multilevel BART model that included random intercepts and random slopes by combining BART with the Stan algorithm. However, the random intercept and slope are modeled by Stan, which is a parametric method. Their results showed that their algorithm ‘stan4bart’ showed better coverage of the population value and lower Root Mean Squared Error (RMSE) compared to BART models with varying intercept, BART models ignoring the multilevel structure, Bayesian Causal Forests (BCF), and parametric multilevel models.

In spite of these promising findings: tree-based model performing well in single-level imputation context [Burgette and Reiter, 2010, Silva and Gutman, 2022, Stekhoven and Bühlmann, 2012, Waljee et al., 2013, Xu et al., 2016] and M-BART models performing well in a multilevel prediction context [Chen, 2020, Dorie et al., 2022, Tan et al., 2016, Wagner et al., 2020, Wundervald et al., 2022], M-BART models have yet to be implemented in a multilevel multiple imputation context. Thus, my research question will be: *Can multivariate imputation by chained equations through a multilevel bayesian additive regression trees model improve the bias, variance and coverage of the estimates in a multilevel context compared to current practices?* Given the success of non-parametric models in single-level multiple imputation, I anticipate that employing multilevel BART models in a multilevel missing data context will reduce bias, accurately model variance, and improve estimate coverage compared to classical multilevel imputation through *2l.pmm*, *2l.lmer*, *2l.pan*, *2l.jomo*, *rf* and single-level *pmm* and complete case analysis in the R-package MICE [Buuren and Groothuis-Oudshoorn, 2011].

This research report is organised as follows: in section 2, will contain some theoretical background and describe the methods in which I will implement the M-BART model in a multilevel imputation context and Section 3 will provide some preliminary results.

2 Method

2.1 Theoretical background

Bayesian Additive Regression Trees (BART) is a sum-of-trees model proposed by Chipman et al. [Chipman et al., 2010]. Regression trees are its building blocks [Chipman et al., 2010, Hill et al., 2020, James et al., 2021]. Regression trees model non-linearities well and automatically through recursive binary partitioning of the predictor space [Burgette and Reiter, 2010, Hill et al., 2020]. Recursive binary partitioning

doesn't assume a specific data form; it divides the predictor space to maximize variance explanation by automatically identifying best fitting splits [Hastie, 2017, James et al., 2021, Salditt et al., 2023]. BART models can be described as:

$$y_i = f(\mathbf{x}_i) + \epsilon_i, \quad (1.1)$$

$$y_i = g(\mathbf{x}_i, T_1, M_1) + g(\mathbf{x}_i, T_2, M_2) + \dots + g(\mathbf{x}_i, T_k, M_k) + \epsilon_i, \quad (1.2)$$

where y_i is the outcome variable for person i , $f(\mathbf{x}_i)$ is the sum-of-trees many regression trees, and ϵ_i is the error term; $\epsilon \sim \mathcal{N}(0, \sigma^2)$. \mathbf{x} are the predictors included in the model, T_k is the k^{th} tree and M_k is the collection of leaf parameters within the k^{th} tree [Chipman et al., 2010, Hill et al., 2020, James et al., 2021]. Next to the sum-of-trees model, BART also includes a regularization prior that constrains the size and fit of each tree so that each contributes only a small part of the overall fit to prevent overfitting [Chipman et al., 2010, Hill et al., 2020, James et al., 2021]. The Bayesian back-fitting Markov Chain Monte Carlo (MCMC) algorithm is used to obtain estimates from BART. It updates each tree, conditional on the remaining trees, their associated parameters and σ , by fitting a new tree to the partial residuals, r_i , perturbing the tree from the previous iteration. The partial residuals, r_i , are defined as:

$$r_i = y_i - \sum_{k' < k} \hat{f}_{k'}^b(x_i) - \sum_{k' > k} \hat{f}_{k'}^{b-1}(x_i), \quad (2)$$

where $\hat{f}_{k'}^b(x_i)$ is the prediction of the k'^{th} tree in the b^{th} iteration for person i . The M-BART model including a random intercept can be identified as:

$$y_{ij} = \sum_{k=1}^m f(\mathbf{X}_{ij}; T_k, M_k) + \alpha_j + \epsilon_{ij}, \quad (3)$$

where, now, y_{ij} is the outcome variable for person i in cluster j and α_j is the random intercept for cluster j .

2.2 Simulation study

I will conduct a simulation study to examine the performance of the M-BART model in a multilevel multiple imputation context through FCS. I will compare the M-BART model to the following multilevel and single level imputation methods: *2l.pmm*, *2l.lmer*, *2l.pan*, *2l.jomo*, *rf* and *pmm* in MICE [Buuren and Groothuis-Oudshoorn, 2011]. The following five factors will be varied:

1. *Intraclass Correlation* (ICC = 0, .05, .3 and .5)
2. *Number of clusters* ($J = 30$ and 50)
3. *Within-cluster sample size* ($n_j = 5, 15, 25$ and 50)
4. *The Missing At Random (MAR) and Missing Completely At Random (MCAR) data rate* (0%, 25% and 50%)
5. *Within-group effect size*: (size of the regression coefficients $\beta = .2, .5$ and $.8$)

Table 1: Simulation design

Parameter	Values
Number of clusters	30, 50
Within-cluster sample size	5, 15, 35, 50
ICC	0, .05, .3, .5
Missing data mechanism	MAR, MCAR
Amount of missingness	0%, 25%, 50%
Within-group effect size	.2, .5, .8

All these values are realistic in practice and/or previously proposed [Enders et al., 2020, 2018b, Grund et al., 2018b, Gulliford et al., 1999, Hox et al., 2017, Murray and Blitstein, 2003]. The ICC can be interpreted as the expected correlation between two randomly sampled individuals from the same group or the proportion of the total variance at level-2 [Gulliford et al., 2005, Hox and Roberts, 2011, Shieh, 2012]. For each combination of varying parameters, a 1000 replicated data sets will be generated.

The simulation study will be performed in R [R Core Team, 2023] with the package MICE [Buuren and Groothuis-Oudshoorn, 2011] to perform the FCS imputations, which I will enhance by integrating a multilevel BART model. FCS multilevel imputation methods *2l.pmm*, *2l.lmer*, *2l.pan*, *2l.jomo*, *rf* and *pmm* in MICE and complete case analysis will serve as a benchmark. The population data-generating mechanism will be based on the following multilevel model:

$$y_{ij} = \beta_{0j} + \beta_{1j}X_{1ij} + \beta_{2j}X_{2ij} + \beta_{3j}X_{3ij} + \beta_{4j}X_{4ij} + \beta_{5j}X_{5ij} + \beta_{6j}X_{6ij} + \beta_{7j}X_{7ij} + \epsilon_{ij}, \quad (4)$$

$$\beta_{0j} = \gamma_{00} + \gamma_{01}Z_{1j} + v_{0j}, \quad (4.1)$$

$$\beta_{1j} = \gamma_{10} + \gamma_{11}Z_{1j} + v_{1j}, \quad (4.2)$$

$$\beta_{2j} = \gamma_{20} + \gamma_{21}Z_{1j} + v_{2j}, \quad (4.3)$$

$$\beta_{3j} = \gamma_{30} + \gamma_{32}Z_{2j} + v_{3j}, \quad (4.4)$$

$$\beta_{4j} = \gamma_{40} + v_{4j}, \quad (4.5)$$

$$\beta_{5j} = \gamma_{50} + v_{5j}, \quad (4.6)$$

$$\beta_{6j} = \gamma_{60} + v_{6j}, \quad (4.7)$$

$$\beta_{7j} = \gamma_{70}, \quad (4.8)$$

where y_{ij} is a continuous level 1 outcome variable for person i in group j and Z_{1j} and Z_{2j} are continuous level 2 variables. The random intercept β_{0j} is determined by the grand mean γ_{00} , the group effect $\gamma_{01}Z_{1j}$ and the group-level random residuals v_{0j} . The regression coefficients β_{1j} , β_{2j} , and β_{3j} for the continuous variables X_{1ij} , X_{2ij} , and X_{3ij} depend on the intercepts γ_{10} , γ_{20} , and γ_{30} , the cross-level interactions $\gamma_{11}Z_{1j}$, $\gamma_{21}Z_{1j}$, and $\gamma_{32}Z_{2j}$, and the random slopes v_{1j} , v_{2j} , and v_{3j} . The regression coefficients β_{4j} , β_{5j} and β_{6j} are determined by the intercepts γ_{40} , γ_{50} and γ_{60} and the random slopes v_{4j} , v_{5j} and v_{6j} . The regression coefficient β_{7j} is determined by the intercept γ_{70} . The residuals and random slopes v_{0j} , v_{1j} , v_{2j} , v_{3j} , v_{4j} , v_{5j} , v_{6j} and ϵ_{ij} follow a zero-mean normal distribution. The variance of v_{0j} , the group-level random residuals, were scaled such that the ICC value was as specified. v_{1j} , v_{2j} , v_{3j} , v_{4j} , v_{5j} , and v_{6j} all have a variance of 1. X_1 , X_2 , X_3 , X_4 , X_5 , X_6 and X_7 are multivariate normally distributed: $\mathbf{X}_{ij} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, with $\boldsymbol{\mu} = (0, 0, 0, 0, 0, 0, 0)$ and $\text{diag}(\boldsymbol{\Sigma}) = (6.25, 9, 4, 11.56, 4, 2.5, 19.36)$ with no co-variances. The level-2 variables Z_1 and Z_2 are also multivariate normally distributed: $\mathbf{Z}_j \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, with $\boldsymbol{\mu} = (0, 0)$ and $\text{diag}(\boldsymbol{\Sigma}) = (1, 2.56)$ with no co-variances.

The estimates will be evaluated on their relative bias (the difference between the average estimate and the true value), modeled variance and the 95% confidence interval coverage [Oberman and Vink, 2023], which will be calculated as follows [Morris et al., 2019]:

$$\text{Bias} = \frac{1}{n_{\text{sim}}} \sum_{i=1}^{n_{\text{sim}}} (\hat{\theta}_i - \theta), \quad (5)$$

$$\text{Coverage} = \frac{1}{n_{\text{sim}}} \sum_{i=1}^{n_{\text{sim}}} I(\theta \in [\hat{\theta}_{\min,i}, \hat{\theta}_{\max,i}]), \quad (6)$$

$$SE = \frac{1}{n_{\text{sim}}} \sum_{i=1}^{n_{\text{sim}}} \sqrt{\frac{\sigma_{\hat{\theta}_i}^2}{N}}, \quad (7)$$

where $\hat{\theta}_i$ is the estimated parameter, θ is the true value, n_{sim} is the number of simulated data sets, $\hat{\theta}_{\min,i}$ and $\hat{\theta}_{\max,i}$ are the lower and upper bounds of the 95% confidence interval, $\sigma_{\hat{\theta}_i}^2$ is the variance of the estimated parameter and N is the size of each simulated data set.

In this research report, the preliminary results will pertain to the evaluation of different BART models in terms of the bias and MSE of the estimates. The MSE will be calculated as follows [Morris et al.,

2019]:

$$MSE = \frac{1}{n_{\text{sim}}} \sum_{i=1}^{n_{\text{sim}}} (\hat{\theta}_i - \theta)^2, \quad (8)$$

where $\hat{\theta}_i$ is the estimated parameter, θ is the true value, and n_{sim} is the number of simulated data sets. In order to reduce computational time, 6 data sets will be simulated for each scenario. Furthermore, since missing data mechanisms are not relevant for this research report, they will not be incorporated in the simulated data sets. 576 data sets will be simulated in total. 4 different models will be compared: a single level BART, single level BART with group-dummy's, a multilevel BART model incorporating a random intercept [Chen, 2020, Tan et al., 2016, Wagner et al., 2020, Wundervald et al., 2022], and a multilevel BART model combined with Stan to model the random parts of the models [Dorie et al., 2022]. The first three models will be performed in R [R Core Team, 2023] with the package *dbarts* [Dorie, 2023a] and the last will performed with the R-package *stan4bart* [Dorie, 2023b]. The default arguments from the function *rbart_vi* will be used for all models, as well as the default priors.

3 Results

The results of this research report are shown in figure 1 and 2. Figure 1 shows the average relative bias for all models, simulated data sets, and every *ICC* value. On the x-axis we can see the different simulated data sets with their names specifying the parameters used in generation. Figure 2 shows the average Mean Squared Error (MSE) for all models, data sets and *ICC* values.

In figure 1 we can see that when there is no multilevel structure in the data set, $ICC = 0$, the models perform similarly in terms of relative bias: overall, the bias is around zero. We can see a slight increase in relative bias when the group size decreases, or more specifically, when the total sample size is small relative bias seems to slightly increase. When we increase the amount of multilevel structure in the data set, $ICC > 0$, we can see a divide in the performance of the models. Even with a little multilevel structure, $ICC = .05$, the single level BART model (*bart*) increases in relative bias. The single level BART model shows higher levels of relative bias when the total sample size is small. However, when the total sample size is large, 2500, the single level BART model performs better than the single level BART model incorporating a group-dummy. The same pattern is observed for the other models: when $ICC = .05$, the relative bias decreases when the total sample size increases. This pattern is further continued when increasing the *ICC*: the relative bias decreases when the total sample size increases for both $ICC = .3$ and $ICC = .5$. However, when considering the *stan4bart* model, which models the random parts of the model in Stan and the fixed parts in BART, the relative bias stay considerably constant when increasing the *ICC*. While the relative bias is higher when the total sample size is small, it does not seem to significantly increase when introducing more multilevel structure. This cannot be said for the *bart* and *gbart* models, which show a slight increase in relative bias when increasing the *ICC*, especially in the lower total sample sizes. The *rbart* model, which incorporates a random intercept, shows a similar pattern as the *stan4bart* model, but with overall more relative bias.

Figure 1: Bias of the estimates for all simulated data sets, all ICC values, and differing within group effect sizes using four models. bart = single level BART model, gbart = single level BART model with group-dummys, rbart = multilevel BART model incorporating a random intercept, stan4bart = multilevel BART model combined with Stan to model the random parts of the models. Data set names show the total sample size, number of groups, and group size. Bias is based on 6 replicates for every data set.





Figure 2 shows a similar pattern as figure 1. When there is no multilevel structure in data sets, $ICC = 0$ the models perform well and almost exactly the same. When introducing multilevel structure, we can start to see a divide in the performance of the models. When $ICC = .05$ the models bart, gbart and rbart perform similarly, with gbart outperforming the other two models when the group sizes are 35 or larger. However, stan4bart consistently outperforms the other three models in terms of MSE. Introducing more multilevel structure to the data, $ICC = .3$, divides the performance of the models further: in small data sets bart now has the highest MSE with gbart performing only a little better. rbart performs better than bart and gbart, but when the data sets increase in size, it performs similar to bart with gbart outperforming them both. stan4bart is still the best overall lowest MSE. Increasing the ICC further to .5 exaggerates these patterns. Over all ICC values, increasing the within group effect size increases the MSE of the models.

Figure 2: Mean Squared Error (MSE) of the estimates for all simulated data sets, all ICC values, and differing within group effect sizes using four models. bart = single level BART model, gbart = single level BART model with group-dummies, rbart = multilevel BART model incorporating a random intercept, stan4bart = multilevel BART model combined with Stan to model the random parts of the models. Data set names show the total sample size, number of groups, and group size. MSE is based on 6 replicates for every data set.



4 Discussion

References

- Audigier, V., White, I. R., Jolani, S., Debray, T. P. A., Quartagno, M., Carpenter, J., Van Buuren, S., and Resche-Rigon, M. (2018). Multiple Imputation for Multilevel Data with Continuous and Binary Variables. *Statistical Science*, 33(2).
- Austin, P. C., White, I. R., Lee, D. S., and Van Buuren, S. (2021). Missing Data in Clinical Research: A Tutorial on Multiple Imputation. *Canadian Journal of Cardiology*, 37(9):1322–1331.
- Bartlett, J. W., Seaman, S. R., White, I. R., Carpenter, J. R., and for the Alzheimer’s Disease Neuroimaging Initiative* (2015). Multiple imputation of covariates by fully conditional specification: Accommodating the substantive model. *Statistical Methods in Medical Research*, 24(4):462–487.
- Breiman, L., Friedman, J. H., Olshen, R. A., and Stone, C. J. (1984). *Classification And Regression Trees*. Routledge, 1 edition.
- Burgette, L. F. and Reiter, J. P. (2010). Multiple Imputation for Missing Data via Sequential Regression Trees. *American Journal of Epidemiology*, 172(9):1070–1076.
- Buuren, S. V. and Groothuis-Oudshoorn, K. (2011). **Mice** : Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software*, 45(3).
- Carpenter, J. R. and Kenward, M. G. (2013). *Multiple Imputation and Its Application*. Wiley, 1 edition.
- Chen, S. (2020). *A New Multilevel Bayesian Nonparametric Algorithm and Its Application in Causal Inference*. PhD thesis, Texas A&M University.
- Chipman, H. A., George, E. I., and McCulloch, R. E. (2010). BART: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4(1).
- Dong, M. and Mitani, A. (2023). Multiple imputation methods for missing multilevel ordinal outcomes. *BMC Medical Research Methodology*, 23(1):112.
- Dorie, V. (2023a). Dbarts: Discrete bayesian additive regression trees sampler.
- Dorie, V. (2023b). *Stan4bart: Bayesian Additive Regression Trees with Stan-Sampled Parametric Extensions*.
- Dorie, V., Perrett, G., Hill, J. L., and Goodrich, B. (2022). Stan and BART for Causal Inference: Estimating Heterogeneous Treatment Effects Using the Power of Stan and the Flexibility of Machine Learning. *Entropy*, 24(12):1782.
- Enders, C. K. (2017). Multiple imputation as a flexible tool for missing data handling in clinical research. *Behaviour Research and Therapy*, 98:4–18.
- Enders, C. K., Du, H., and Keller, B. T. (2020). A model-based imputation procedure for multilevel regression models with random coefficients, interaction effects, and nonlinear terms. *Psychological Methods*, 25(1):88–112.
- Enders, C. K., Hayes, T., and Du, H. (2018a). A Comparison of Multilevel Imputation Schemes for Random Coefficient Models: Fully Conditional Specification and Joint Model Imputation with Random Covariance Matrices. *Multivariate Behavioral Research*, 53(5):695–713.
- Enders, C. K., Keller, B. T., and Levy, R. (2018b). A fully conditional specification approach to multilevel imputation of categorical and continuous variables. *Psychological Methods*, 23(2):298–317.
- Enders, C. K., Mistler, S. A., and Keller, B. T. (2016). Multilevel multiple imputation: A review and evaluation of joint modeling and chained equations imputation. *Psychological Methods*, 21(2):222–240.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2016). Multiple imputation of missing covariate values in multilevel models with random slopes: A cautionary note. *Behavior Research Methods*, 48(2):640–649.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2018a). Multiple Imputation of Missing Data at Level 2: A Comparison of Fully Conditional and Joint Modeling in Multilevel Designs. *Journal of Educational and Behavioral Statistics*, 43(3):316–353.

- Grund, S., Lüdtke, O., and Robitzsch, A. (2018b). Multiple Imputation of Missing Data for Multilevel Models: Simulations and Recommendations. *Organizational Research Methods*, 21(1):111–149.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2021). Multiple imputation of missing data in multilevel models with the R package mdmb: A flexible sequential modeling approach. *Behavior Research Methods*, 53(6):2631–2649.
- Gulliford, M., Adams, G., Ukoumunne, O., Latinovic, R., Chinn, S., and Campbell, M. (2005). Intraclass correlation coefficient and outcome prevalence are associated in clustered binary data. *Journal of Clinical Epidemiology*, 58(3):246–251.
- Gulliford, M. C., Ukoumunne, O. C., and Chinn, S. (1999). Components of Variance and Intraclass Correlations for the Design of Community-based Surveys and Intervention Studies: Data from the Health Survey for England 1994. *American Journal of Epidemiology*, 149(9):876–883.
- Hastie, T. J., editor (2017). *Statistical Models in S*. Routledge, 1st edition.
- Hill, J., Linero, A., and Murray, J. (2020). Bayesian Additive Regression Trees: A Review and Look Forward. *Annual Review of Statistics and Its Application*, 7(1):251–278.
- Hox, J. and Roberts, J. K., editors (2011). *Handbook of Advanced Multilevel Analysis*. Routledge, 0 edition.
- Hox, J. J., Moerbeek, M., and Van De Schoot, R. (2017). *Multilevel Analysis: Techniques and Applications*. Routledge, Third edition. — New York, NY : Routledge, 2017. —, 3 edition.
- Hughes, R. A., White, I. R., Seaman, S. R., Carpenter, J. R., Tilling, K., and Sterne, J. A. (2014). Joint modelling rationale for chained equations. *BMC Medical Research Methodology*, 14(1):28.
- James, G., Witten, D., Hastie, T., and Tibshirani, R. (2021). *An Introduction to Statistical Learning: With Applications in R*. Springer Texts in Statistics. Springer US, New York, NY.
- Kang, H. (2013). The prevention and handling of the missing data. *Korean Journal of Anesthesiology*, 64(5):402.
- Lin, S. and Luo, W. (2019). A New Multilevel CART Algorithm for Multilevel Data with Binary Outcomes. *Multivariate Behavioral Research*, 54(4):578–592.
- Lüdtke, O., Robitzsch, A., and Grund, S. (2017). Multiple imputation of missing data in multilevel designs: A comparison of different strategies. *Psychological Methods*, 22(1):141–165.
- Meng, X.-L. (1994). Multiple-imputation inferences with uncongenial sources of input. *Statistical science*, pages 538–558.
- Mistler, S. A. and Enders, C. K. (2017). A Comparison of Joint Model and Fully Conditional Specification Imputation for Multilevel Missing Data. *Journal of Educational and Behavioral Statistics*, 42(4):432–466.
- Morris, T. P., White, I. R., and Crowther, M. J. (2019). Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11):2074–2102.
- Murray, D. M. and Blitstein, J. L. (2003). Methods To Reduce The Impact Of Intraclass Correlation In Group-Randomized Trials. *Evaluation Review*, 27(1):79–103.
- Oberman, H. I. and Vink, G. (2023). Toward a standardized evaluation of imputation methodology. *Biometrical Journal*, page 2200107.
- Quartagno, M. and Carpenter, J. R. (2022). Substantive model compatible multilevel multiple imputation: A joint modeling approach. *Statistics in Medicine*, 41(25):5000–5015.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.
- Resche-Rigon, M. and White, I. R. (2018). Multiple imputation by chained equations for systematically and sporadically missing multilevel data. *Statistical Methods in Medical Research*, 27(6):1634–1649.
- Rubin, D. B. (1976). Inference and missing data. *Biometrika*, 63(3):581–592.

- Rubin, D. B. (1987). *Multiple Imputation for Nonresponse in Surveys*. Wiley, New York.
- Salditt, M., Humberg, S., and Nestler, S. (2023). Gradient Tree Boosting for Hierarchical Data. *Multivariate Behavioral Research*, pages 1–27.
- Shieh, G. (2012). A comparison of two indices for the intraclass correlation coefficient. *Behavior Research Methods*, 44(4):1212–1223.
- Silva, G. C. and Gutman, R. (2022). Multiple imputation procedures for estimating causal effects with multiple treatments with application to the comparison of healthcare providers. *Statistics in Medicine*, 41(1):208–226.
- Stekhoven, D. J. and Bühlmann, P. (2012). MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics*, 28(1):112–118.
- Taljaard, M., Donner, A., and Klar, N. (2008). Imputation Strategies for Missing Continuous Outcomes in Cluster Randomized Trials. *Biometrical Journal*, 50(3):329–345.
- Tan, Y. V., Flannagan, C. A. C., and Elliott, M. R. (2016). Predicting human-driving behavior to help driverless vehicles drive: Random intercept Bayesian Additive Regression Trees.
- Van Buuren, S. (2007). Multiple imputation of discrete and continuous data by fully conditional specification. *Statistical Methods in Medical Research*, 16(3):219–242.
- van Buuren, S. (2018). *Flexible Imputation of Missing Data*. Chapman & Hall/CRC Interdisciplinary Statistics Series. CRC Press, Taylor & Francis Group, Boca Raton London New York, second edition edition.
- Wagner, J., West, B. T., Elliott, M. R., and Coffey, S. (2020). Comparing the Ability of Regression Modeling and Bayesian Additive Regression Trees to Predict Costs in a Responsive Survey Design Context. *Journal of Official Statistics*, 36(4):907–931.
- Waljee, A. K., Mukherjee, A., Singal, A. G., Zhang, Y., Warren, J., Balis, U., Marrero, J., Zhu, J., and Higgins, P. D. (2013). Comparison of imputation methods for missing laboratory data in medicine. *BMJ Open*, 3(8):e002847.
- Wundervald, B., Parnell, A., and Domijan, K. (2022). Hierarchical Embedded Bayesian Additive Regression Trees.
- Xu, D., Daniels, M. J., and Winterstein, A. G. (2016). Sequential BART for imputation of missing covariates. *Biostatistics*, 17(3):589–602.