

Master Thesis:

Multilevel Multivariate Imputation by Chained Equations through Bayesian Additive Regression Trees

Methodology and Statistics for the Behavioural, Biomedical and Social Sciences

Heleen Brüggen



Word count:

4036

Candidate Journal:

Computational Statistics & Data Analysis

FETC Case Number:

23-1778

Supervisors:

T. Volken MSc.

Utrecht University

Dr. G. Vink

Utrecht University

H. Oberman MSc.

Utrecht University

Contents

1	Introduction	2
2	Method	3
2.1	Theoretical background	3
2.1.1	Bayesian Additive Regression Trees (BART)	3
2.1.2	Multilevel-BART (M-BART)	5
2.1.3	stan4bart	5
2.2	Simulation study	6
2.2.1	Data generating mechanism	6
2.2.2	Simulation design	7
2.2.3	Missing data generation	8
2.2.4	Evaluation	9
3	Results	9
3.1	Bias	9
3.2	Coverage	15
3.3	Confidence interval width	17
4	Discussion	21
5	Conclusion	22
6	Appendix	23
	References	28

1 Introduction

1. ask someone to read for flow (mama, merlijn, tuanke???)

Incomplete data is a common challenge in many fields of research. Frequently used ad hoc strategies to deal with missing data, such as complete case analysis or mean imputation, often lead to erroneous inferences in realistic situations. Missingness can follow a multivariate mechanism that may depend on observed data or even unobserved data, leading to biased estimates and inaccurate variance estimates when using one of these ad hoc strategies (Austin et al., 2021; Enders, 2017; Kang, 2013; Little and Rubin, 2002; van Buuren, 2018). Multiple imputation (MI; Rubin, 1987) is proven to be an effective method for dealing with multivariate incomplete data supported by a considerable amount of methodological research (Audigier et al., 2018; Austin et al., 2021; Burgette and Reiter, 2010; Enders, 2017; Grund et al., 2021; Hughes et al., 2014; Little and Rubin, 2002; Mistler and Enders, 2017a; Van Buuren, 2007; van Buuren, 2018).

MI separates the missing data problem from the analysis problem (Audigier et al., 2018; Austin et al., 2021; Bartlett et al., 2015; Burgette and Reiter, 2010; Carpenter and Kenward, 2013; Enders, 2017; Grund et al., 2021; Hughes et al., 2014; Little and Rubin, 2002; Mistler and Enders, 2017a; Van Buuren, 2007; van Buuren, 2018). A statistical model specifying the variables used for imputation, i.e. the imputation model, is defined for every variable with missing values. Each missing value in the dataset is imputed m times by drawing values from their posterior predictive distribution conditional on the observed data and parameters from the imputation model. By repeatedly drawing values from the posterior predictive distributions – in other words, the distribution of plausible replacement values – the necessary variation associated with the missingness problem is considered. After imputation, each of the imputed datasets are analyzed according to the model of interest, i.e. the substantive analysis model. Then, their m corresponding model parameters are pooled together according to Rubin's rules (Rubin, 1987). One central requirement for MI is the concept of congeniality; the imputation model should be at least as general as the analysis model and preferably all-encompassing (Bartlett et al., 2015; Enders et al., 2018a; Grund et al., 2016, 2018b; Little and Rubin, 2002; Meng, 1994). If not, the imputation model will not be compatible with the analysis model and the pooled estimates of the latter may be biased.

When MI is applied in a multilevel data context, concerns regarding the concept of congeniality become more pronounced (Audigier et al., 2018; Dong and Mitani, 2023; Enders et al., 2020, 2018a,b, 2016; Grund et al., 2016, 2018a,b, 2021; Lüdtke et al., 2017; Mistler and Enders, 2017a; Quartagno and Carpenter, 2022; Resche-Rigon and White, 2018; Taljaard et al., 2008; van Buuren, 2018). Multilevel data is hierarchically structured, where, for example, students are nested within classes within schools or patients within hospitals (Hox and Roberts, 2011; Hox et al., 2017). When analyzing multilevel data, this hierarchical structure should be taken into consideration. Ignoring it will underestimate the intra-class correlation (ICC) and standard errors, as conventional statistical analyses assume independence of observations (Hox and Roberts, 2011; Lüdtke et al., 2017; Taljaard et al., 2008; van Buuren, 2018). The ICC can be interpreted as the proportion of the total variance at level-2 (Gulliford et al., 2005; Hox and Roberts, 2011; Shieh, 2012). Accounting for this structure, can be done using multilevel models (MLMs; Hox and Roberts, 2011; Hox et al., 2017; Lüdtke et al., 2017). MLMs can contain variables relating to the individual level – level-1 variables – or to the grouping structure – level-2 variables or potentially higher order structures. For example, imagine a case where students are nested within classes. Here, the academic performance of a student is a level-1 variable, whereas the teacher's experience is a level-2 variable. Additionally, MLMs allow you to specify random intercepts, indicating that some classes have students that significantly perform better or worse academically on average; random slopes, indicating that the relationship between the performance of students and the outcome variable differs between classes; and cross-level interactions, indicating that the effect of performance of students can differ with the teacher's experience (Hox and Roberts, 2011; Hox et al., 2017). Typically, the complexity of the multilevel analysis model is built step-wise with non-linearities, meaning the analysis model is not determined beforehand: predictors, random intercepts, random slopes, and cross-level interactions are added in a stepwise manner to the model (Hox and Roberts, 2011; Hox et al., 2017). Thus, ensuring congeniality for the imputation model can be complex, since the final analysis model is not pre-determined. Furthermore, including the hierarchical structure along with cross-level interactions or other complicated non-linearities in imputation models is quite challenging (Burgette and Reiter, 2010; Hox and Roberts, 2011; van Buuren, 2018), also because very complex models might not converge (van Buuren, 2018).

A popular and flexible implementation of MI in a multilevel context, is fully conditional specification (FCS), otherwise known as chained equations (Audigier et al., 2018; Burgette and Reiter, 2010; Grund et al., 2018a; Van Buuren, 2007). FCS employs univariate linear mixed models to account for the hierarchical

structure of multilevel models (Enders et al., 2018a; Mistler and Enders, 2017a; Resche-Rigon and White, 2018) and iteratively imputes each incomplete variable conditional on observed and previously imputed variables (Enders et al., 2018a,b, 2016; Grund et al., 2018a; Hughes et al., 2014; Mistler and Enders, 2017a; van Buuren, 2018). Furthermore, it can impute non-linearities, such as cross-level interactions, by using ‘passive imputation’ or defining a separate imputation model for the non-linearities (Grund et al., 2018b; van Buuren, 2018). However, including these non-linearities in FCS is still very complicated (Grund et al., 2018b, 2021; van Buuren, 2018). FCS can also handle random intercepts and slopes, yet, once again, correctly specifying an imputation model accounting for these random effects can be challenging (Grund et al., 2018b, 2021; van Buuren, 2018).

Non-parametric, tree-based models might alleviate these complexities when defining imputation models. They do not assume a specific data distribution. So, they implicitly model non-linear relationships and can simultaneously handle continuous and categorical variables (Breiman et al., 1984; Burgette and Reiter, 2010; Chipman et al., 2010; Hill et al., 2020; James et al., 2021; Lin and Luo, 2019; Salditt et al., 2023). Studies showed that the use of tree-based, non-parametric models like regression trees, random forests, or Bayesian Additive Regression Trees (BART) in imputation of single-level data simplified the imputation process (Burgette and Reiter, 2010; Silva and Gutman, 2022; Waljee et al., 2013; Xu et al., 2016). They showed better model parameter estimates than parametric methods. Specifically, the imputations showed better confidence interval coverage of the parameters, lower variance and lower bias, especially in non-linear and interactive contexts (Burgette and Reiter, 2010; Silva and Gutman, 2022; Xu et al., 2016). Waljee et al. (2013) also found lower missclassification error rate for the predicted class as well as lower imputation error when imputing with a random forest algorithm compared to multivariate imputation by chained equations (`mice`) using linear, logistic, and polytomous logistic regression imputation models, K-nearest neighbors (KNN) and mean imputation.

In prediction, multilevel-BART models (M-BART) have predominantly been implemented with random intercepts only (Chen, 2020; Tan et al., 2016; Wagner et al., 2020; Wundervald et al., 2022). Wagner et al. (2020) have found that this random intercept M-BART model provided better predictions with a lower mean squared error (MSE) compared to a parametric MLM, Tan et al. (2016) found higher area under the curve (AUC) values compared to a singel-level BART model and linear logistic random intercept model, and Chen (2020) found better predictions and better coverage of the parameter estimates compared to parametric models and a single-level BART model. Other researchers modeled the random intercept as an extra split on each terminal node and found a lower MSE compared to a standard BART model and parametric MLMs (Wundervald et al., 2022). Dorie et al. (2022) developed a multilevel BART model that included random intercepts, random slopes and cross-level interactions by modeling these random parts with a Stan (Lee et al., 2017) model and the fixed parts with a BART model. Their results showed that their algorithm `stan4bart` showed better coverage of the population values and lower root mean squared error (RMSE) compared to BART models with varying intercept, BART models ignoring the multilevel structure, bayesian causal forests, and parametric MLMs.

Despite these promising findings, M-BART models have yet to be implemented in a multilevel multiple imputation context. Thus, my research question will be: *Can multivariate imputation by chained equations through a multilevel bayesian additive regression trees model improve the bias, variance, and coverage of the multilevel model parameter estimates compared to current practices?* Given the success of non-parametric models in single-level MI, I anticipate that employing M-BART models in a multilevel missing data context will reduce bias, accurately model variance, and improve estimate coverage compared to conventional implementations of multilevel MI, single-level MI, and complete case analysis in the R-package `mice` (Buuren and Groothuis-Oudshoorn, 2011).

2 Method

2.1 Theoretical background

2.1.1 Bayesian Additive Regression Trees (BART)

BART is a sum-of-trees model proposed by Chipman et al. (2010) with regression trees as its building blocks (Chipman et al., 2010; Hill et al., 2020; James et al., 2021). Regression trees recursively split the data into binary subgroups based on the predictors included in the model. At each step down the tree, these splits are based on the predictor that minimizes the variability within the subgroups from all predictors. Observations are then assigned to a certain subgroup according to these splits. This is continued until a certain stopping criterion is reached; for example, we desire a minimal number of

observations with in a subgroup (Breiman et al., 1984; Hastie, 2017; James et al., 2021; Salditt et al., 2023). Recursive binary partitioning of the predictor space doesn't assume a specific data form. This making regression trees, and as a consequence, BART, non-parametric models (Breiman et al., 1984; Hastie, 2017; James et al., 2021; Salditt et al., 2023) and allows regression trees to model non-linearities and other complicated relationships well and automatically (Burgette and Reiter, 2010; Hill et al., 2020). Chipman et al. (2010) define the BART model as:

$$f(\mathbf{x}) = \sum_{k=1}^K g(\mathbf{x}; T_k, M_k), \quad (1)$$

where $f(\mathbf{x})$ is the overall fit of the model: the sum of K regression trees, \mathbf{x} are the predictor variables, T_k is the k^{th} tree and M_k is the collection of leaf parameters within the k^{th} tree, i.e. the collection of predictions for its terminal nodes (Chipman et al., 2006, 1998, 2010; Hill et al., 2020; James et al., 2021). The data are assumed to arise from a model with additive normally distributed errors: $Y = \sum_{k=1}^K g(\mathbf{x}; T_k, M_k) + \epsilon, \epsilon \sim \mathcal{N}(0, \sigma^2)$. Next to the sum-of-trees model, BART also includes a regularization prior that constrains the size and fit of each tree so that each contributes only a small part of the variation in the outcome variables to prevent overfitting. The prior is imposed over all parameters of the sum-of-trees model, specifically, $(T_1, M_1), \dots, (T_K, M_K)$ and σ . However, the specification of the regularization prior is simplified by a series of independence assumptions:

$$\begin{aligned} p((T_1, M_1), \dots, (T_K, M_K), \sigma) &= \left[\prod_k p(T_k, M_k) \right] p(\sigma), \\ &= \left[\prod_k p(M_k|T_k) p(T_k) \right] p(\sigma), \\ p(M_k|T_k) &= \prod_j p(\mu_{jk}|T_k), \end{aligned} \quad (2)$$

where $\mu_{jk} \in M_k$. These assumptions state that the trees (T_k), leaf parameters ($\mu_j|T_k$), and the standard deviation (σ) are independent of each other. Thus, priors only need to be specified for those parameters (Chipman et al., 2006, 1998, 2010; Hill et al., 2020). Chipman et al. (1998) define an independent prior for each tree. The probability that a node at depth d splits is defined as:

$$\alpha(1+d)^{-\beta}, \alpha \in (0, 1), \beta \in [0, \infty), \quad (3)$$

where the default specification put forth by Chipman et al. (2006, 2010) is $\alpha = .95$ and $\beta = 2$. This specification sets the probability of a tree with 1, 2, 3, 4, and 5 nodes at .05, .55, .28, .09, and .03 respectively. Thus, smaller trees are favoured. Chipman et al. (2006, 2010) also provide a default specification for the prior for the leaf parameters. They propose to rescale the response value to the interval $[-.5, .5]$. Then, the leaf parameter prior is defined as:

$$\mu_{jk} \sim \mathcal{N}(0, \sigma_\mu^2), \text{ with } \sigma_\mu^2 = \frac{.5}{t\sqrt{K}}, \quad (4)$$

where t is a preselected number and K is the number of trees. This prior shrinks the tree parameters μ_{jk} towards 0, decreasing the effect of the individual tree components. If t or K increase, more shrinkage is applied. Chipman et al. (2006, 2010) found good results with and recommend using $t = 2$ – or values between 1 and 3 – as a default choice. Furthermore, Chipman et al. (2006, 2010) propose the conjugate inverse chi-square distribution as the prior for the residual standard deviation $\sigma^2 \sim \nu\lambda/\chi_\nu^2$. They represent the degrees of freedom, λ , as the probability that the BART residual standard deviation, σ , is less than the estimated residual standard deviation from a linear regression model, $\hat{\sigma}_{\text{OLS}}$. Their default specification of the hyperparameters is $\nu = 3$ and $\Pr(\sigma < \hat{\sigma}_{\text{OLS}}) = .9$ (Chipman et al., 2006, 1998, 2010; Hill et al., 2020).

BARTs are estimated using the Bayesian back-fitting Markov Chain Monte Carlo (MCMC) algorithm (Chipman et al., 2006, 1998, 2010; Hill et al., 2020; James et al., 2021). Each tree is initialized with a single root node with the mean response value divided by the number of trees ($\hat{f}_k^1(x) = \frac{1}{nK} \sum_{i=1}^n y_i$, with sample size n). Then, each pair (T_k, M_k) is updated considering the remaining trees, their associated parameters,

and the residual standard deviation (σ) by sampling from the following conditional distribution:

$$(T_k, M_k) | T_{k'}, M_{k'}, \sigma, y. \quad (5)$$

However, this conditional distribution only depends on $(T_{k'}, M_{k'}, y)$ through the partial residuals:

$$r_i = y_i - \sum_{k' < k} \hat{f}_{k'}^b(x_i) - \sum_{k' > k} \hat{f}_{k'}^{b-1}(x_i), \text{ with } i = 1, \dots, n, \quad (6)$$

where $\hat{f}_k^b(x_i)$ is the prediction of the k^{th} tree in the b^{th} iteration for person i and sample size n . Thus, updating each pair (T_k, M_k) simplifies to proposing a new tree fit to the partial residuals, r_i , treating them as the data, by perturbing the tree from the previous iteration. Perturbations entail either *growing*, *pruning*, or *changing* a tree. *Growing* means adding additional splits, *pruning* removes splits, and *changing* changes decision rules. The algorithm stops after the specified number of iterations (Chipman et al., 2006, 1998, 2010; Hill et al., 2020; James et al., 2021).

2.1.2 Multilevel-BART (M-BART)

1. check flow and understanding
2. incorporate feedback
3. change title??? $-_i$ should also be changed in many other places!!!

Tan et al. (2016); Wagner et al. (2020) and Dorie et al. (2024) define an M-BART model including a random intercept. The BART model (1) is extended to include a random intercept by:

$$m(\mathbf{x}) = \sum_{k=1}^K g(\mathbf{x}; T_k, M_k) + \alpha_j, \quad (7)$$

where, now, $f(\mathbf{x})$ is the overall fit of the model incorporating random intercept α_j for cluster j and. So, the data are now assumed to arise from the following model:

$$Y_{ij} = \sum_{k=1}^K g(\mathbf{x}; T_k, M_k) + \alpha_j + \epsilon_{ij}, \quad \epsilon_{ij} \sim \mathcal{N}(0, \sigma^2), \quad \alpha_j \sim \mathcal{N}(0, \tau^2), \quad (8)$$

where $\alpha_j \perp \epsilon_{ij}$. Now the joint prior distribution (2) becomes:

$$\begin{aligned} p((T_1, M_1), \dots, (T_K, M_K), \sigma) &= \left[\prod_k p(T_k, M_k) \right] p(\sigma) p(\tau), \\ &= \left[\prod_k p(M_k | T_k) p(T_k) \right] p(\sigma) p(\tau), \\ p(M_k | T_k) &= \prod_j p(\mu_{jk} | T_k). \end{aligned} \quad (9)$$

A Metropolis within Gibbs procedure is used to draw values from the posterior. First, the Gibbs sample for σ , τ , and α_j are obtained from their respective posterior distributions. Then, we obtain $\tilde{Y}_{ij} = Y_{ij} - \alpha_j$ and view $\tilde{Y}_{ij} | \mathbf{X}_j$ as a BART model. So, \tilde{Y} is now used as the outcome variable in the BART algorithm described in the previous section, 2.1.1. (Tan et al., 2016; Wagner et al., 2020). Dorie et al. (2024) implemented this algorithm within the R-package `dbarts` with the function `rbart_vi()`. Where, the default prior for the random intercept is $\tau \sim \text{Cauchy}(0, 2.5)$: a Cauchy distribution with a scale parameter 2.5 times the original scale.

2.1.3 stan4bart

1. check flow and understanding
2. incorporate feedback

Dorie et al. (2022) developed a multilevel BART model that included random intercepts, random slopes, and cross-level interactions. They extend a Bayesian linear, mixed model with a BART model (1). The resulting model is:

$$h(\mathbf{x}) = \mathbf{x}^\beta \boldsymbol{\beta} + f(\mathbf{x}; T_K, M_K) + \boldsymbol{\lambda} \mathbf{w}, \quad (10)$$

where \mathbf{x}^β is a vector of 1 – for the intercept – and the linear predictors; $\boldsymbol{\beta}$ is a vector of linear, parametric coefficients; $\boldsymbol{\lambda}$ is a vector of all parametric random slopes and intercepts; \mathbf{w} is a vector of the coefficients for the random slopes and intercepts; and $f(\mathbf{x}; T_K, M_K)$ is a non-parametric, sum-of-trees BART model (Dorie et al., 2022). So, the data are assumed to arise from the following model:

$$Y_{ij} = \mathbf{x}^\beta \boldsymbol{\beta} + f(\mathbf{x}; T_K, M_K) + \boldsymbol{\lambda} \mathbf{w} + \epsilon_{ij}, \quad \epsilon_{ij} \sim \mathcal{N}(0, \sigma^2), \quad \boldsymbol{\lambda} \sim \mathcal{N}(0, \boldsymbol{\Sigma}_\lambda), \quad (11)$$

where $\boldsymbol{\Sigma}_\lambda$ is the variance-covariance matrix for the random intercept and slopes. The model is implemented as a Gibbs sampler: a Hamiltonian Monte Carlo, no-U-turn sampler with a diagonal Euclidean adaptation matrix is used to jointly sample the linear, parametric components given the non-parametric components. The non-parametric components are sampled using the BART algorithm described in section 2.1.1 (Dorie et al., 2022). To accomplish this, a parametric Stan model (Lee et al., 2017) fits equation 10 with $f(\mathbf{x}; T_K, M_K)$ as a generic linear offset. Dorie et al. (2022) combine a custom mutable Stan sampler object with a BART sampler with a fixed variance and offset term: First, the Stan sampler collects the current draws of the BART model into $\text{vec}_i f(\mathbf{x}_i; T_K, M_K)$ and uses this to draw $\beta, \lambda, \sigma, \boldsymbol{\Sigma}_\lambda | \mathbf{Y}, \text{vec}_i f(\mathbf{x}_i; T_K, M_K)$. Then, σ and $\text{vec}_i [\mathbf{x}_i^\beta \boldsymbol{\beta} + \boldsymbol{\lambda} \mathbf{w}_i]$ are passed to BART, which produces $M_k, T_k | \mathbf{Y}, \text{vec}_i [\mathbf{x}_i^\beta \boldsymbol{\beta} + \boldsymbol{\lambda} \mathbf{w}_i], \sigma, M_{k'}, T_{k'}$. Then, the cycle is completed by passing $\text{vec}_i f(\mathbf{x}_i; T_K, M_K)$ back to Stan. The process is continued for the set amount of posterior samples intended for inference. This algorithm is implemented in the R-package `stan4bart` (Dorie, 2023).

2.2 Simulation study

2.2.1 Data generating mechanism

1. check R reference

We assembled a simulation study to evaluate the performance of multilevel BART models in a multilevel imputation context. The population data-generating mechanism is based on the following MLM:

$$y_{ij} = \beta_{0j} + \sum_{k=1}^7 \beta_{kj} X_{kij} + \epsilon_{ij}, \quad X_{kij} \sim \mathcal{MVN}(0, \boldsymbol{\Sigma}_x), \quad (12a)$$

$$\beta_{0j} = \gamma_{00} + \sum_{p=1}^2 \gamma_{0q} Z_{pj} + v_{0j}, \quad (12b)$$

$$\beta_{kj} = \gamma_{k0} + \sum_{p=1}^2 \gamma_{kp} Z_{pj} + v_{kj}, \quad Z_{pj} \sim \mathcal{MVN}(0, \boldsymbol{\Sigma}_z), \quad (12c)$$

where y_{ij} is a continuous level-1 outcome variable for person i in group j and X_{kij} are 7 continuous level-1 variables and Z_{pj} are 2 continuous level-2 variables. The predictors are multivariate normally distributed with means of 0 and variance-covariance matrix $\boldsymbol{\Sigma}_x$ and $\boldsymbol{\Sigma}_z$, respectively:

$$\boldsymbol{\Sigma}_x = \begin{pmatrix} 6.25 & & & & & & \\ 2.25 & 9 & & & & & \\ 1.5 & 1.8 & 4 & & & & \\ 2.25 & 3.06 & 2.04 & 11.56 & & & \\ 1.5 & 1.8 & 1.2 & 2.04 & 4 & & \\ 1.125 & 1.35 & 0.9 & 1.53 & .9 & 2.25 & \\ 3.3 & 3.96 & 2.64 & 4.488 & 2.64 & 1.98 & 19.36 \end{pmatrix}, \quad (13a)$$

$$\boldsymbol{\Sigma}_z = \begin{pmatrix} 1 & & \\ .48 & 2.56 \end{pmatrix}. \quad (13b)$$

The covariances between the variables are calculated such that the correlation between the variables is .3,

aligned with Cohen's (1990) medium effect size benchmark. The residuals are normally distributed as,

$$\epsilon_{ij} \sim \mathcal{N}(0, 25). \quad (14)$$

The random intercept β_{0j} is determined by the overall intercept γ_{00} , the 2 group-level effects $\gamma_{0q}Z_{pj}$ and the group-level random residuals v_{0j} . The overall intercept γ_{00} is set to 10 and the group-level effects γ_{01} and γ_{02} to .5. The 7 regression coefficients β_{kj} for the continuous variables X_{kij} depend on the intercepts γ_{k0} , the cross-level interactions $\gamma_{kp}Z_{pj}$, and the random slopes v_{kj} . The 7 intercepts, or within-group effect sizes, γ_{k0} are set to .5, the cross-level interactions γ_{11} , γ_{21} , and γ_{32} are set to .35.

$$\gamma_{00} = 10, \quad \gamma_{0p} = \begin{pmatrix} .5 \\ .5 \\ .5 \\ .5 \\ .5 \\ .5 \\ .5 \end{pmatrix}, \quad \gamma_{k0} = \begin{pmatrix} .5 \\ .5 \\ .5 \\ .5 \\ .5 \\ .5 \\ .5 \end{pmatrix}, \quad \gamma_{kp} = \begin{pmatrix} .35 & 0 \\ .35 & 0 \\ 0 & .35 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}. \quad (15)$$

The random slopes are multivariate normally distributed with a mean of 0 and a variance-covariance matrix \mathbf{T} shown in equation 16. Again, the covariances are calculated to yield a correlation of .3.

$$\mathbf{v}_j \sim \mathcal{MVN}(0, \mathbf{T}), \quad \mathbf{T} = \begin{pmatrix} t_{00} & & & & & & & \\ .3 & 1 & & & & & & \\ .3 & .3 & 1 & & & & & \\ .3 & .3 & .3 & 1 & & & & \\ 0 & 0 & 0 & 0 & 0 & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (16)$$

The variance of v_{0j} , the group-level random residuals t_{00} , are scaled such that the specified ICC values as in table 1 was obtained. The following formula is used to calculate v_{0j} following the variance decomposition from Rights and Sterba (2019):

$$\text{ICC} = \frac{\gamma^b' \phi^b \gamma^b + \tau_{00}}{\gamma^w' \phi^w \gamma^w + \gamma^b' \phi^b \gamma^b + \text{tr}(\mathbf{T}\Sigma) + \tau_{00} + \sigma^2}, \quad (17)$$

where γ^b and γ^w are the level-1 and level-2 fixed effects; ϕ^b is the variance-covariance matrix of a vector with 1, for the intercept, and all level-2 predictors; ϕ^w is the variance-covariance matrices of all cluster-mean-centered level-1 predictors; τ_{00} is the variance of the random intercept; \mathbf{T} is the variance-covariance matrix of the random intercept and slopes; Σ is the variance-covariance matrix of a vector containing 1, for the intercept, and the level-1 variables; and σ^2 is the residual variance. The value for τ_{00} is calculated using the function `uniroot()` in R (R Core Team, 2023).

2.2.2 Simulation design

1. check flow
2. add difference in simulation for stan4bart (when how many data sets and why)
3. change order of how this are discussed
4. check explanation of how I implemented the imputation methods

Table 1 shows the design factors considered in the simulation study. These factors are either grounded in prior research or deemed realistic in real-world applications (Enders et al., 2020, 2018b; Grund et al., 2018b; Gulliford et al., 1999; Hox et al., 2017; Murray and Blitstein, 2003). According to Kreft and de Leeuw (2007), 30 groups is the smallest acceptable number in multilevel research and 50 groups is frequent in organizational research (Maas and Hox, 2005). Group sizes of 15 are typical in educational research (Lüdtke et al., 2017) and group sizes of 50 are often used in simulation studies (Akkaya Hocagil

and Yucel, 2023; Enders et al., 2020, 2018a,b; Grund et al., 2018b; Maas and Hox, 2005). The ICC was chosen to be .5, which is often used as an upper limit in methodological research (Enders et al., 2020, 2018a,b; Grund et al., 2018b; Mistler and Enders, 2017b; Salditt et al., 2023). Oberman and Vink (2023) recommend including both Missing Completely At Random (MCAR) and Missing At Random (MAR) missingness mechanisms in simulation studies. They pose that the statistical properties of the imputation method are not deemed sound if it cannot yield valid inferences under MCAR. Furthermore, they pose that including observed-data-dependend missingness – for example, MAR – is of utmost importance in evaluating the imputation method’s performance. The amount of missingness in data sets is varied between 0% and 50%. 0% missingness is included as an additional benchmark and 50% missingess is often used in simulation studies as a high amount of missingness (Grund et al., 2016; Lüdtke et al., 2017; Schouten and Vink, 2021). For each combination of design factors, 100 datasets are simulated. 5 different imputation methods are compared:

1. conventional single-level imputation with PMM (predictive mean matching),
2. conventional multilevel imputation with PMM,
3. single-level BART imputation,
4. multilevel BART imputation accounting for random intercepts (Chen, 2020; Tan et al., 2016; Wagner et al., 2020),
5. multilevel BART imputation accounting for random effects and cross-level interactions (Dorie et al., 2022).

The first and second methods are implemented with the R-packages `mice` and `miceadds` (Robitzsch et al., 2024). The conventional single-level imputation is implemented with the imputation method `pmm` and the conventional multilevel imputation is implemented with the `2l.pmm` method for level-1 variables and `2lonly.mean` for level-2 variables.

The third, single-level BART, fourth, random intercept BART and fifth method, multilevel BART methods are implemented by writing new method-functions in R (R Core Team, 2023) for the package `mice`. The functions `bart` and `rbart_vi` from the `dbarts` package were used for the single-level and random intercept BART imputation methods (Dorie et al., 2024). The function `stan4bart` from the package `stan4bart` was used for the multilevel BART imputation method accounting for random effects and cross-level interactions (Dorie, 2023). The functions were written such that they can be used as imputation methods in the `mice` package. All three functions are implementend as follows: for every variable to be imputed, a respective BART model is fitted based on the predictor matrix. Then, the fitted values – the posterior means – are extracted for the observed and missing values. Imputations for the missing values are then obtain using predictive mean matching by matching the predicted values for the observed cases to the predicted values for the missing cases. The code for these functions can be found in the appendix – listing 1, 2, and 3.

For all imputation methods, the incomplete data sets are imputed 5 times with 10 iterations each. Then, each of the 5 imputed datasets are then analyzed using the R-package `lme4` (Bates et al., 2015) with an MLM reflecting the population generating mechanism: $y = 1 + x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + z_1 + z_2 + x_1 * z_1 + x_2 * z_1 + x_3 * z_2 + (1 + x_1 + x_2 + x_3 | group)$. The estimates from the 5 imputed datasets are pooled together using the R-package `mice` (Buuren and Groothuis-Oudshoorn, 2011). These pooled estimates are compared on the bias, coverage, and the width of the 95% confidence intervals.

As an additional benchmark, the imputation methods will also be compared to analyses using listwise deletion, i.e. complete case analysis, and using the true data without missing values.

2.2.3 Missing data generation

Missing values in the variables are introduced by multivariate amputation using the function `ampute()` (Schouten et al., 2018) from package `mice`. As can be seen in table 1, the missing data mechanism is either Missing Completely At Random (MCAR) or Missing At Random (MAR). The missing data mechanism

is said to be MCAR when the cause of the missing data is unrelated to the data and MAR when the missing data is related to the observed data (Rubin, 1976). The amount of missingness is either 0% or 50%, which is defined as the percentage of cases that have at least one missing value.

For both MCAR and MAR, all possible patterns with 1 to 5 missing values out of the 10 variables ($x_1, x_2, x_3, x_4, x_5, x_6, x_7, z_1, z_2$, and y) per case are generated. They have the same relative frequency of occurrence in the data sets. So, 50% of the cases had 1 to 5 missing values.

For the MAR mechanism, the weighted sum of scores on the observed variables is used to predict the probability of missingness for a case. The weights of the variables x_4 and z_1 are set to 2 and 1.5 respectively when they remain observed in a specific pattern, while the weights of the other variables that remain observed in a specific pattern are set to 1. The type of missingness is set to ‘RIGHT’ meaning that cases with a higher weighted sum of scores have a higher probability of becoming incomplete. So, this means that cases with higher values on x_4 and z_1 are more likely to become incomplete.

In summary, either no missing values are introduced (0%), or up to 5 missing values are introduced in 50% of the cases. When data is MAR, the probability of a value being missing depends on the observed values of all other variables, with variables x_4 and z_1 having a greater influence on this probability.

2.2.4 Evaluation

1. add relative bias explanation/definition
2. add relative bias evaluation

The estimates from the analysis models are evaluated in terms of absolute bias, coverage of 95% confidence intervals, with their respective Monte Carlo SE (MCSE), and the width of the 95% confidence intervals (Morris et al., 2019; Oberman and Vink, 2023):

$$\text{Bias} = \frac{1}{n_{\text{sim}}} \sum_{t=1}^{n_{\text{sim}}} (\hat{\theta}_t - \theta), \quad \text{MCSE}_{\text{Bias}} = \sqrt{\frac{\sum_{t=1}^{n_{\text{sim}}} (\hat{\theta}_t - \bar{\theta})^2}{n_{\text{sim}}(n_{\text{sim}} - 1)}}, \quad (18)$$

$$\text{Coverage} = \frac{1}{n_{\text{sim}}} \sum_{t=1}^{n_{\text{sim}}} \mathbb{1}(\hat{\theta}_{\text{low},i} \leq \theta \leq \hat{\theta}_{\text{upp},i}), \quad \text{MCSE}_{\text{Coverage}} = \sqrt{\frac{\text{Cov}\hat{\text{erage}}(1 - \text{Cov}\hat{\text{erage}})}{n_{\text{sim}}}}, \quad (19)$$

$$\text{CIW} = \frac{1}{n_{\text{sim}}} \sum_{t=1}^{n_{\text{sim}}} (\hat{\theta}_{\text{upp},i} - \hat{\theta}_{\text{low},i}), \quad (20)$$

where $\hat{\theta}_t$ is the estimated parameter in simulation t , θ is the true value, $\bar{\theta}$ is the mean of $\hat{\theta}_t$, and n_{sim} is the number of simulated datasets. The lower and upper bounds of the 95% confidence intervals are denoted as $\hat{\theta}_{\text{low},i}$ and $\hat{\theta}_{\text{upp},i}$ respectively. The coverage is the proportion of the 95% confidence intervals that contain the true value.

Enders et al. (2018a); Morris et al. (2019); Oberman and Vink (2023); van Buuren (2018) suggest that a coverage of 95% is acceptable. Poor coverage, i.e. below 95%, indicates biased estimates or too narrow intervals. While, coverage above 95% indicates that efficiency could still be gained. The width of the confidence intervals is a measure of the statistical precision of the estimates: a smaller width indicates a more precise estimate (Oberman and Vink, 2023; van Buuren, 2018).

3 Results

1. check flow
2. check importance of results described
3. check if some figure can be moved to appendix (- i , some thing may not be relevant enough?)

3.1 Bias

Figures 1 through 6 show the absolute bias of the estimates of the linear mixed model for all imputation methods in consideration with Monte Carlo SE. The absolute bias of the overall intercept; the level-1 effects; the level-2 effects; cross-level interactions; the random slopes; and the residual and intercept variance are shown in figures 1, 2, 3, 4, 5, and 6 respectively.

First, the estimates of the fixed effects – the overall intercept, γ_{00} ; level-1 effects $\gamma_{10} : \gamma_{70}$; level-2 effects γ_{01} and γ_{02} ; and the cross-level interactions γ_{11}, γ_{21} , and γ_{32} – will be considered in terms of absolute bias. Then, the random structure of the model – the random intercept v_0 ; the random slopes v_1, v_2 , and v_3 ; and the residual variance, ϵ_{ij} – will be considered.

From figure 1 it can be seen that when the data is MAR, the overall intercept is acceptably biased – the absolute biases fall between the 10% relative bias lines – for all imputation methods when sample size is smallest – i.e. 30 groups of size 15. When the group size is increased to 50, for imputation methods PMM, 2l.PMM, BART, and R-BART it is close to zero. Yet, stan4bart increased in bias, now overestimating the intercept. Eventhough, the simulation uncertainty for stan4bart does still encompass the zero-bias line. With 50 groups of size 15, all imputation methods underestimate the intercept beyond the 10% relative bias line. Yet, increasing the group size to 50, slightly decreases the bias for PMM, 2l.PMM, BART, and R-BART. Stan4bart now overestimates the intercept, yet the simulation uncertainty still encompasses the zero-bias line. Overall, when the data is MAR, BART performs best out of all imputation methods. While stan4bart is the worst in terms of absolute bias, its MCSE always encompasses the zero-bias line. Listwise deletion underestimates the intercept compared to the 0%, “true” data. When the data is MCAR, there seem to be less differences in performance between the imputation methods. All methods overestimate the intercept with 30 groups of size 15 and are acceptably biased with the other groups conditions. However, stan4bart overestimates the intercept with 50 groups of size 50.




Figure 1: Absolute bias of the overall intercept of the linear mixed model with respective Monte Carlo SE for all simulated data sets over 100 simulations with ICC = .5. The dashed lines represent $\pm 10\%$ relative bias.

Figure 2 shows the absolute bias of the level-1 effects. When the sample size is smallest – i.e. 30 groups of size 15 – fluctuations in bias is more common for all methods when the data is MAR or MCAR. 2l.PMM, PMM, and BART seem to perform best out of all imputation methods for both missingness mechanisms, especially with a larger total sample size. Stan4bart seems to overestimate some fixed effects, which increases with more and larger groups and when the data is MAR. Overall, R-BART has the worst performance in terms of absolute bias, especially when the data is MAR with more groups.




Figure 2: Absolute bias of the level-1 effects of the linear mixed model with respective Monte Carlo SE for all simulated data sets over 100 simulations with $\text{ICC} = .5$.

Considering the level-2 effects from figure 3 – γ_{01} and γ_{02} –, stan4bart performs the worst out of all imputation methods for all conditions: it greatly underestimates the level-2 effects. 2l.PMM performs best out of all imputation methods, even though still over- or underestimating the level-2 effects at times and seemingly performing slightly worse under MCAR. PMM and R-BART consistently underestimate the level-2 effects for all conditions. However, BART performs slightly better, at times even mimicking the performance of 2l.PMM. Overall, only when the data is MAR do all methods seem to benefit from smaller groups when there are 50 groups, reducing in bias when they are smaller, and larger groups when there are 30 groups. Thus, preferring a larger sample size.




Figure 3: Absolute bias of the level-2 effects of the linear mixed model with respective Monte Carlo SE for all simulated data sets over 100 simulations with ICC = .5.

For the cross-level interactions – γ_{11}, γ_{21} , and γ_{32} –, stan4bart has, again, the worst performance out of all methods. Figure 4 shows that stan4bart consistently underestimates the cross-level interaction for all conditions. For both MAR and MCAR, listwise deletion performs largely acceptable in terms of bias, often staying within the 10% relative bias lines. 2l.PMM performs best out of all imputation methods, however still underestimating the cross-level interactions regularly. Furthermore, 2l.PMM performs slightly better under MCAR than MAR. BART consistently outperforms PMM and R-BART, however, still underestimating the cross-level interactions. BART, R-BART and PMM, perform slightly better under MCAR compared to MAR when group size are 15.




Figure 4: Absolute bias of the cross-level interactions of the linear mixed model with respective Monte Carlo SE for all simulated data sets over 100 simulations with ICC = .5.

The absolute bias for the random slopes in figure 5 show that overall, stan4bart has the best performance out of all imputation methods. When the data is MAR, stan4bart provides acceptable biases when group sizes are 15. However, when group sizes are 50, stan4bart underestimates the random slopes. This underestimation reduces when there are more groups. PMM, BART and R-BART have the worst performance: they consistently underestimate the random slopes for all factor conditions. While 2l.PMM does perform better than PMM, BART, and R-BART, it still underestimates the random slopes for almost all conditions. For most conditions, all methods seem to perform worse under MCAR and when groups are larger. Listwise deletion performs largely acceptable in terms of bias, most of the time staying within the 10% relative bias lines for both MAR and MCAR. However, under MCAR, listwise deletion does tend to slightly overestimate the random slopes, but to an acceptable extend – not more than the acceptable 10% relative bias.




Figure 5: Absolute bias of the random slopes of the linear mixed model with respective Monte Carlo SE for all simulated data sets over 100 simulations with ICC = .5.

Lastly, the absolute bias for the intercept and residual variance will be discussed. From figure 6 it can be seen that stan4bart and 2l.PMM have an acceptable bias for the intercept variance for almost all conditions. Stan4bart seems to slightly overestimate the intercept variance when the data is MAR compared to MCAR, however, still within the 10% relative bias lines. Also, stan4bart improves in absolute bias when there are more groups in the data set. 2l.PMM pretty consistently underestimates – not more than 10% in terms of relative bias – the intercept variance for all conditions. PMM, R-BART and BART routinely underestimate the intercept variance for all conditions. From these three imputation methods, BART performs best, however still far below the acceptable threshold. Furthermore, BART seems to improve in bias when there are larger groups. Listwise deletion shows a very slight bias, underestimating the intercept variance when the data is MAR.

Looking at the residual variance, listwise deletion is the only method that is acceptable biased for all – or any – condition. All other imputation methods routinely overestimate the residual variance. Stan4bart has the best performance followed by 2l.PMM, BART, PMM, and R-BART, in that order. Overall, the bias seems to be markedly consistent across all conditions. However, stan4bart seems to increase in bias when the total sample size increases and 2l.PMM seems to decrease in bias with more groups in the data set.




Figure 6: Bias of the e_{ij} and u_0

3.2 Coverage

Figures 7 and 8 show the 95% confidence interval coverage of the estimates of the linear mixed model – the overall intercept and the level-1 effects; and the level-2 effects – for all imputation methods in consideration with Monte Carlo SE.

First, the estimates of the fixed effects – the overall intercept, γ_{00} and level-1 effects $\gamma_{10} : \gamma_{70}$ – will be considered in terms of coverage. Then, the level-2 effects – γ_{01} and γ_{02} – will be considered.

Figure 7 shows that the coverage of the overall intercept, γ_{00} , is best for imputation method 2l.PMM when the data is MAR. However, when the data is MCAR, stan4bart performs best. 2l.PMM shows slight undercoverage in the smallest sample size when the data is MCAR. While, stan4bart shows more fluctuations in coverage when the data is MAR. For all conditions, PMM shows undercoverage, especially when there are 50 groups and the data is MAR. R-BART also shows consistent undercoverage of the intercept for all conditions except with 30 groups of size 15 when the data is MAR. When the data is MAR, BART undercovers the intercept for all conditions. However, when the data is MCAR and there are 60 groups, BART shows acceptable coverage.

Next, the coverage of the level-1 effects in figure 7 shows that 2l.PMM performs best out of all imputation methods for all conditions. At times even showing overcoverage. Stan4bart also shows a good performance in terms of coverage, however, it regularly overcovers and sometimes undercovers the level-1 effects. PMM at times shows under- or overcoverage, while BART more consistently shows undercoverage. R-BART has most fluctuations in coverage: when group sizes are 50, it shows undercoverage as low as around 47,5% and overcoverage as high as around 98%. Furthermore, stan4bart shows great coverage of the level-1 effects with a 100 simulations – i.e. when the data is MCAR with 30 groups of size 15.




Figure 7: Coverage of the estimates

Lastly, the coverage of the level-2 effects and cross-level interactions in figure 8 shows that BART has the overall worst coverage for all conditions, with coverage ranging from 65% to 82.5%. PMM also routinely undercovers the level-2 effects, which worsens with group sizes of 50. Then, R-BART tends to overcover the level-2 effects, but undercovers the level-2 effect of $z_2 - \gamma_{02}$ – when there are 50 groups of size 50 for both MAR and MCAR. Overall, 2l.PMM performs best in terms of coverage. Eventhough, 2l.PMM shows slight undercoverage when the data is MAR and there are 50 groups.

Then, the coverage of the cross-level interactions shows that 2l.PMM has the best coverage of the imputation methods. However, 2l.PMM still shows a slight under- or overcoverage at times. Listwise deletion also performs considerably good, showing better coverage when the data is MAR compared MCAR. When the group size is 15, PMM; BART; and R-BART, show an acceptable coverage – sometimes under- or overcovering the cross-level interactions –, but show considerable undercoverage when the group size is increased to 50. Stan4bart has the worst coverage of the cross-level interactions. It shows considerable undercoverage for all conditions, especially when the group size is increased to 50.




Figure 8: Coverage of the estimates

3.3 Confidence interval width

Figures 9, 10, and 11 show the 95% confidence interval width of the estimates of the linear mixed model – the overall intercept and the level-1 effects; and the level-2 effects – for all imputation methods in consideration.

First, the estimates of the fixed effects – the overall intercept, γ_{00} and level-1 effects $\gamma_{10} : \gamma_{70}$ – will be considered in terms of coverage. Then, the level-2 effects – γ_{01} and γ_{02} – will be considered.

First the confidence interval width of the intercept, γ_{00} for all imputation methods will be considered. Figure 9 shows that the confidence interval width of the intercept is smallest for PMM suggesting the highest statistical efficiency. However, paired with the coverage estimates from figure 7, PMM seems to be efficient but routinely undercovers the intercept. They same pattern can be seen for BART and R-BART, while showing smaller confidence interval widths, they routinely undercover the intercept. 2l.PMM, stan4bart and listwise deletion show confidence interval widths that somewhat mimic the “true” data with 0% missing values. Though their confidence interval widths are larger, they oft show acceptable coverage.




Figure 9: Confidence interval width of the estimates

For the level-1 effects, $\gamma_{10} : \gamma_{70}$, the confidence interval width of the “true” data with 0% missing values is oft smallest. Listwise deletion and stan4bart are closest in mimicking the width of the “true” data and figure 7 shows generally show acceptable coverage of the level-1 effects. 2l.PMM and BART have overall slightly larger confidence intervals, with BARTs intervals being smaller than 2l.PMM, however, as mentioned before, 2l.PMM has the best coverage of the level-1 effects while BART shows more undercoverage. Then, PMM and R-BART show the largest confidence intervals. PMM shows both under- and overcoverage in figure 7, while R-BART shows considerable undercoverage when its confidence intervals are smallest and overcoverage when its confidence intervals are largest – for example when the data is MAR with 50 groups of 50 for effect γ_{70} and γ_{60} . All methods seem to show a similar pattern in the confidence interval width when the total sample size increases: the confidence interval width decreases towards the “true” data’s confidence interval width.




Figure 10: Confidence interval width of the estimates

From figure 11 we can see that 2l.PMM has largest confidence interval width of all imputation methods, however, considerably closely mimicking the confidence interval from the 0% missing data. The other imputation methods, PMM; BART; R-BART; and stan4bart, show smaller confidence intervals. Listwise deletion performs similar to 2l.PMM and the 0% missing data. Furthermore, PMM; BART; R-BART; and stan4bart show a similar pattern in the confidence interval width when the total sample size increases: the confidence interval width decreases leading to more undercoverage – as can be seen in figure 8. At the same time, 2l.PMM and listwise deletion show a decrease in width when there are more groups in the data, which is a pattern mirrored by the “true” data as well.




Figure 11: Confidence interval width of the estimates

The width of the 95% confidence intervals for the cross-level interaction – shown in figure 12 show a similar pattern for all methods: when group sizes are 15, all methods have confidence intervals larger than the 0% missing data. However, when group sizes are 50, the confidence intervals decrease in size for all method, either being smaller than or similar to the 0% missing data. This pattern is also reflected in figure 8, where these smaller intervals result in undercoverage.




Figure 12: Confidence interval width of the estimates

4 Discussion

1. Interpretation of results

- (a) overall best performance of 2l.PMM
- (b) overall worst performance of single level method PMM and BART
- (c) R-BART really bad in intercept variance, fixed effects, level-2 effects, cross-level, residual variance, random slopes $-\zeta$ at times performing even worse than BART. Would be interesting to see how it would perform if there are no random slopes in the data, thus only random intercepts (which it is supposedly modelling).
- (d) Listwise deletion performs really good, even under MAR which is surprising $-\zeta$ might be modelling a special case.
- (e) stan4bart has an okay performance. Certainly one that hopefully spraks interest. It performs badly in level-2 effects (fixed and cross-level), however it might be interesting how this would perform with including passive imputation (?? dont know if that is even possible). However, this would make the imputation method more complex and less user-friendly $-\zeta$ it would look very similar to the imputation method 2l.PMM in terms of specification. stan4bart does outperform 2l.PMM in terms of random effects and residual variance. one big downside of this method was the computational time (loop 4 took around 3 days to run).

2. comparison with previous research

3. limitations of this study

- (a) 100 replications, sometimes 20
- (b) computational time
- (c) not many different factors
- (d) matching method
- (e)

4. future research recommendations

- (a) interesting explorations for stan4bart: reduce computational time, matching procedure for predictive mean matching, passive imputation, MAR and MCAR, different amount of missingness, different effect sizes, smaller sample sizes.

5 Conclusion

1. summarise the main findings
2. reiterate the importance of this study

6 Appendix

Listing 1: Imputation function for single-level BART

```

1  mice.impute.bart <- function(y, ry, x, wy = NULL, use.matcher = FALSE, donors = 5L,
2  ...){
3      install.on.demand("dbarts", ...)
4      if (is.null(wy)) {
5          wy <- !ry
6      }
7
8      # Parameter estimates
9      fit <- dbarts::bart(x, y, keeptrees = TRUE, verbose = FALSE)
10
11     yhatobs <- fitted(fit, type = "ev", sample = "train")[ry]
12     yhatmis <- fitted(fit, type = "ev", sample = "train")[wy]
13
14     # Find donors
15     if (use.matcher) {
16         idx <- matcher(yhatobs, yhatmis, k = donors)
17     } else {
18         idx <- matchindex(yhatobs, yhatmis, donors)
19     }
20
21     return(y[ry][idx])
22 }
```

Listing 2: Imputation function for random intercept BART

```

1  mice.impute.2l.rbart <- function(y, ry, x, wy = NULL, type, use.matcher = FALSE,
2  donors = 5L, ...){
3      install.on.demand("dbarts", ...)
4      if (is.null(wy)) {
5          wy <- !ry
6      }
7
8      clust <- names(type[type == -2])
9      effects <- names(type[type != -2])
10     X <- x[, effects, drop = FALSE]
11
12     model <- paste0(
13         "y ~ ", paste0(colnames(X), collapse = " + "))
14
15     fit <- dbarts::rbart_vif(formula = formula(model), group.by = clust, data =
16     data.frame(y, x), verbose = FALSE, n.threads = 1, n.samples = 500L, n.burn = 500L, ...)
17
18     yhatobs <- fitted(fit, type = "ev", sample = "train")[ry]
19     yhatmis <- fitted(fit, type = "ev", sample = "train")[wy]
20
21     # Find donors
22     if (use.matcher) {
23         idx <- matcher(yhatobs, yhatmis, k = donors)
24     } else {
25         idx <- matchindex(yhatobs, yhatmis, donors)
26     }
27
28     return(y[ry][idx])
29 }
```

Listing 3: Imputation function for multilevel BART with random effects and cross-level interactions

```

1  mice.impute.2l.bart <- function(y, ry, x, wy = NULL, type, intercept = TRUE, use.
2  matcher = FALSE, donors = 5L, ...){
3      install.on.demand("stan4bart", ...)
4      if (is.null(wy)) {
5          wy <- !ry
6      }
7
8      if (intercept) {
9          x <- cbind(1, as.matrix(x))
10         type <- c(2, type)
11     }
12
13     fit <- stan4bart(y = y, x = x, type = type, intercept = intercept, use.matcher =
14     use.matcher, donors = donors, ...)
```

```

10     names(type)[1] <- colnames(x)[1] <- "(Intercept)"
11 }
12
13 clust <- names(type[type == -2])
14 rande <- names(type[type == 2])
15 fixe <- names(type[type > 0])
16
17 lev <- unique(x[, clust])
18
19 X <- x[, fixe, drop = FALSE]
20 Z <- x[, rande, drop = FALSE]
21 xobs <- x[ry, , drop = FALSE]
22 yobs <- y[ry]
23 Xobs <- X[ry, , drop = FALSE]
24 Zobs <- Z[ry, , drop = FALSE]
25
26 # create formula
27 fr <- ifelse(length(rande) > 1,
28   paste0("+ (1 +", paste(rande[-1L], collapse = "+")),
29   " + (1 "
30 )
31 randmodel <- paste0(
32   "y ~ bart(", paste0(fixe[-1L], collapse = " + "), ")",
33   fr, "| ", clust, ")"
34 )
35 fit <- eval(parse(text = paste("stan4bart::stan4bart(", randmodel,
36   ", data = data.frame(y, x),
37   verbose = -1,
38   bart_args = list(k = 2.0, n.samples = 500L, n.burn = 500L, n.thin = 1L, n.
39 threads = 1))",
40   collapse = ""))
41
42 yhatobs <- fitted(fit, type = "ev", sample = "train")[ry]
43 yhatmis <- fitted(fit, type = "ev", sample = "train")[wy]
44
45 # Find donors
46 if (use.matcher) {
47   idx <- matcher(yhatobs, yhatmis, k = donors)
48 } else {
49   idx <- matchindex(yhatobs, yhatmis, donors)
50 }
51
52 return(y[ry][idx])
53 }
```

References

- Akkaya Hocagil, T. and Yucel, R. M. (2023). A computationally efficient sequential regression imputation algorithm for multilevel data. *Journal of Applied Statistics*, pages 1–21.
- Audigier, V., White, I. R., Jolani, S., Debray, T. P. A., Quartagno, M., Carpenter, J., Van Buuren, S., and Resche-Rigon, M. (2018). Multiple Imputation for Multilevel Data with Continuous and Binary Variables. *Statistical Science*, 33(2).
- Austin, P. C., White, I. R., Lee, D. S., and Van Buuren, S. (2021). Missing Data in Clinical Research: A Tutorial on Multiple Imputation. *Canadian Journal of Cardiology*, 37(9):1322–1331.
- Bartlett, J. W., Seaman, S. R., White, I. R., Carpenter, J. R., and for the Alzheimer’s Disease Neuroimaging Initiative* (2015). Multiple imputation of covariates by fully conditional specification: Accommodating the substantive model. *Statistical Methods in Medical Research*, 24(4):462–487.
- Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of Statistical Software*, 67(1).
- Breiman, L., Friedman, J. H., Olshen, R. A., and Stone, C. J. (1984). *Classification And Regression Trees*. Routledge, 1 edition.
- Burgette, L. F. and Reiter, J. P. (2010). Multiple Imputation for Missing Data via Sequential Regression Trees. *American Journal of Epidemiology*, 172(9):1070–1076.
- Buuren, S. V. and Groothuis-Oudshoorn, K. (2011). **Mice** : Multivariate Imputation by Chained Equations in *R*. *Journal of Statistical Software*, 45(3).
- Carpenter, J. R. and Kenward, M. G. (2013). *Multiple Imputation and Its Application*. Wiley, 1 edition.
- Chen, S. (2020). *A New Multilevel Bayesian Nonparametric Algorithm and Its Application in Causal Inference*. PhD thesis, Texas A&M University.
- Chipman, H., George, E., and McCulloch, R. (2006). Bayesian Ensemble Learning. In *Advances in Neural Information Processing Systems*, volume 19. MIT Press.
- Chipman, H. A., George, E. I., and McCulloch, R. E. (1998). Bayesian CART Model Search. *Journal of the American Statistical Association*, 93(443):935–948.
- Chipman, H. A., George, E. I., and McCulloch, R. E. (2010). BART: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4(1).
- Cohen, J. (1990). Statistical power analysis for the behavioral sciences. *Computers, Environment and Urban Systems*, 14(1):71.
- Dong, M. and Mitani, A. (2023). Multiple imputation methods for missing multilevel ordinal outcomes. *BMC Medical Research Methodology*, 23(1):112.
- Dorie, V. (2023). *Stan4bart: Bayesian Additive Regression Trees with Stan-Sampled Parametric Extensions*.
- Dorie, V., Chipman, H., McCulloch, R., Dadgar, A., Team, R. C., Draheim U., G., Bosmans, M., Tournayre, C., Petch, M., Valle, R. d. L., Johnson G., S., Frigo, M., Zaitseff, J., Veldhuizen, T., Maisonneuve, L., Pakin, S., and Daniel G., R. (2024). Dbarts: Discrete Bayesian Additive Regression Trees Sampler.
- Dorie, V., Perrett, G., Hill, J. L., and Goodrich, B. (2022). Stan and BART for Causal Inference: Estimating Heterogeneous Treatment Effects Using the Power of Stan and the Flexibility of Machine Learning. *Entropy*, 24(12):1782.
- Enders, C. K. (2017). Multiple imputation as a flexible tool for missing data handling in clinical research. *Behaviour Research and Therapy*, 98:4–18.
- Enders, C. K., Du, H., and Keller, B. T. (2020). A model-based imputation procedure for multilevel regression models with random coefficients, interaction effects, and nonlinear terms. *Psychological Methods*, 25(1):88–112.

- Enders, C. K., Hayes, T., and Du, H. (2018a). A Comparison of Multilevel Imputation Schemes for Random Coefficient Models: Fully Conditional Specification and Joint Model Imputation with Random Covariance Matrices. *Multivariate Behavioral Research*, 53(5):695–713.
- Enders, C. K., Keller, B. T., and Levy, R. (2018b). A fully conditional specification approach to multilevel imputation of categorical and continuous variables. *Psychological Methods*, 23(2):298–317.
- Enders, C. K., Mistler, S. A., and Keller, B. T. (2016). Multilevel multiple imputation: A review and evaluation of joint modeling and chained equations imputation. *Psychological Methods*, 21(2):222–240.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2016). Multiple imputation of missing covariate values in multilevel models with random slopes: A cautionary note. *Behavior Research Methods*, 48(2):640–649.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2018a). Multiple Imputation of Missing Data at Level 2: A Comparison of Fully Conditional and Joint Modeling in Multilevel Designs. *Journal of Educational and Behavioral Statistics*, 43(3):316–353.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2018b). Multiple Imputation of Missing Data for Multilevel Models: Simulations and Recommendations. *Organizational Research Methods*, 21(1):111–149.
- Grund, S., Lüdtke, O., and Robitzsch, A. (2021). Multiple imputation of missing data in multilevel models with the R package mdmb: A flexible sequential modeling approach. *Behavior Research Methods*, 53(6):2631–2649.
- Gulliford, M., Adams, G., Ukoumunne, O., Latinovic, R., Chinn, S., and Campbell, M. (2005). Intra-class correlation coefficient and outcome prevalence are associated in clustered binary data. *Journal of Clinical Epidemiology*, 58(3):246–251.
- Gulliford, M. C., Ukoumunne, O. C., and Chinn, S. (1999). Components of Variance and Intra-class Correlations for the Design of Community-based Surveys and Intervention Studies: Data from the Health Survey for England 1994. *American Journal of Epidemiology*, 149(9):876–883.
- Hastie, T. J., editor (2017). *Statistical Models in S*. Routledge, 1st edition.
- Hill, J., Linero, A., and Murray, J. (2020). Bayesian Additive Regression Trees: A Review and Look Forward. *Annual Review of Statistics and Its Application*, 7(1):251–278.
- Hox, J. and Roberts, J. K., editors (2011). *Handbook of Advanced Multilevel Analysis*. Routledge, 0 edition.
- Hox, J. J., Moerbeek, M., and Van De Schoot, R. (2017). *Multilevel Analysis: Techniques and Applications*. Routledge, Third edition. | New York, NY : Routledge, 2017. |, 3 edition.
- Hughes, R. A., White, I. R., Seaman, S. R., Carpenter, J. R., Tilling, K., and Sterne, J. A. (2014). Joint modelling rationale for chained equations. *BMC Medical Research Methodology*, 14(1):28.
- James, G., Witten, D., Hastie, T., and Tibshirani, R. (2021). *An Introduction to Statistical Learning: With Applications in R*. Springer Texts in Statistics. Springer US, New York, NY.
- Kang, H. (2013). The prevention and handling of the missing data. *Korean Journal of Anesthesiology*, 64(5):402.
- Kreft, I. and de Leeuw, J. (2007). *Introducing Multilevel Modeling*. Introducing Statistical Methods. SAGE, Los Angeles, Calif., reprinted edition.
- Lee, D., Carpenter, B., Li, P., Morris, M., Betancourt, M., Maverickg, Brubaker, M., Trangucci, R., Inacio, M., Kucukelbir, A., Buildbot, S., Bgoodri, Seantalts, Arnold, J., Tran, D., Hoffman, M., Margossian, C., Modrák, M., Adler, A., Sakrejda, K., Stukalov, A., Lawrence, M., Goedman, R. J., Van Horn, K. S., Vehtari, A., Gabry, J., Casallas, J. S., and Bales, B. (2017). Stan-dev/stan: V2.17.1. Zenodo.
- Lin, S. and Luo, W. (2019). A New Multilevel CART Algorithm for Multilevel Data with Binary Outcomes. *Multivariate Behavioral Research*, 54(4):578–592.
- Little, R. J. A. and Rubin, D. B. (2002). *Statistical Analysis with Missing Data*. Wiley Series in Probability and Statistics. Wiley, 1 edition.

- Lüdtke, O., Robitzsch, A., and Grund, S. (2017). Multiple imputation of missing data in multilevel designs: A comparison of different strategies. *Psychological Methods*, 22(1):141–165.
- Maas, C. J. M. and Hox, J. J. (2005). Sufficient Sample Sizes for Multilevel Modeling. *Methodology*, 1(3):86–92.
- Meng, X.-L. (1994). Multiple-imputation inferences with uncongenial sources of input. *Statistical science*, pages 538–558.
- Mistler, S. A. and Enders, C. K. (2017a). A Comparison of Joint Model and Fully Conditional Specification Imputation for Multilevel Missing Data. *Journal of Educational and Behavioral Statistics*, 42(4):432–466.
- Mistler, S. A. and Enders, C. K. (2017b). A Comparison of Joint Model and Fully Conditional Specification Imputation for Multilevel Missing Data. *Journal of Educational and Behavioral Statistics*, 42(4):432–466.
- Morris, T. P., White, I. R., and Crowther, M. J. (2019). Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11):2074–2102.
- Murray, D. M. and Blitstein, J. L. (2003). Methods To Reduce The Impact Of Intraclass Correlation In Group-Randomized Trials. *Evaluation Review*, 27(1):79–103.
- Oberman, H. I. and Vink, G. (2023). Toward a standardized evaluation of imputation methodology. *Biometrical Journal*, page 2200107.
- Quartagno, M. and Carpenter, J. R. (2022). Substantive model compatible multilevel multiple imputation: A joint modeling approach. *Statistics in Medicine*, 41(25):5000–5015.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.
- Resche-Rigon, M. and White, I. R. (2018). Multiple imputation by chained equations for systematically and sporadically missing multilevel data. *Statistical Methods in Medical Research*, 27(6):1634–1649.
- Rights, J. D. and Sterba, S. K. (2019). Quantifying explained variance in multilevel models: An integrative framework for defining R-squared measures. *Psychological Methods*, 24(3):309–338.
- Robitzsch, A., Simon Grund, and Henke, T. (2024). Miceadds: Some Additional Multiple Imputation Functions, Especially for ‘mice’.
- Rubin, D. B. (1976). Inference and missing data. *Biometrika*, 63(3):581–592.
- Rubin, D. B. (1987). *Multiple Imputation for Nonresponse in Surveys*. Wiley, New York.
- Salditt, M., Humberg, S., and Nestler, S. (2023). Gradient Tree Boosting for Hierarchical Data. *Multivariate Behavioral Research*, pages 1–27.
- Schouten, R. M., Lugtig, P., and Vink, G. (2018). Generating missing values for simulation purposes: A multivariate amputation procedure. *Journal of Statistical Computation and Simulation*, 88(15):2909–2930.
- Schouten, R. M. and Vink, G. (2021). The Dance of the Mechanisms: How Observed Information Influences the Validity of Missingness Assumptions. *Sociological Methods & Research*, 50(3):1243–1258.
- Shieh, G. (2012). A comparison of two indices for the intraclass correlation coefficient. *Behavior Research Methods*, 44(4):1212–1223.
- Silva, G. C. and Gutman, R. (2022). Multiple imputation procedures for estimating causal effects with multiple treatments with application to the comparison of healthcare providers. *Statistics in Medicine*, 41(1):208–226.
- Taljaard, M., Donner, A., and Klar, N. (2008). Imputation Strategies for Missing Continuous Outcomes in Cluster Randomized Trials. *Biometrical Journal*, 50(3):329–345.
- Tan, Y. V., Flannagan, C. A. C., and Elliott, M. R. (2016). Predicting human-driving behavior to help driverless vehicles drive: Random intercept Bayesian Additive Regression Trees.

- Van Buuren, S. (2007). Multiple imputation of discrete and continuous data by fully conditional specification. *Statistical Methods in Medical Research*, 16(3):219–242.
- van Buuren, S. (2018). *Flexible Imputation of Missing Data*. Chapman & Hall/CRC Interdisciplinary Statistics Series. CRC Press, Taylor & Francis Group, Boca Raton London New York, second edition edition.
- Wagner, J., West, B. T., Elliott, M. R., and Coffey, S. (2020). Comparing the Ability of Regression Modeling and Bayesian Additive Regression Trees to Predict Costs in a Responsive Survey Design Context. *Journal of Official Statistics*, 36(4):907–931.
- Waljee, A. K., Mukherjee, A., Singal, A. G., Zhang, Y., Warren, J., Balis, U., Marrero, J., Zhu, J., and Higgins, P. D. (2013). Comparison of imputation methods for missing laboratory data in medicine. *BMJ Open*, 3(8):e002847.
- Wundervald, B., Parnell, A., and Domijan, K. (2022). Hierarchical Embedded Bayesian Additive Regression Trees.
- Xu, D., Daniels, M. J., and Winterstein, A. G. (2016). Sequential BART for imputation of missing covariates. *Biostatistics*, 17(3):589–602.