

# Mathematical modelling of between-pathogen interactions in the human host

Application to influenza viruses and pneumococcus

Hélène Arduin

Biostatistiques, Biomathématique, Pharmacopidémiologie  
et Maladies Infectieuses

UMR 1181 Inserm, UVSQ, Institut Pasteur



**Inserm**



UNIVERSITÉ DE  
VERSAILLES  
ST-QUENTIN-EN-YVELINES  
université PARIS-SACLAY



 Institut Pasteur

June 26, 2019

# Introduction

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

Presence of  $P_1$  :

- ▶  $P_2$ 's growth
- ▶  $P_2$ 's severity
- ▶  $P_2$ 's infection duration

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

Presence of  $P_1$  :

increases →


- ▶  $P_2$ 's growth
- ▶  $P_2$ 's severity
- ▶  $P_2$ 's infection duration

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

## **Synergistic interaction**

increases →



Presence of  $P_1$  :


- ▶  $P_2$ 's growth
- ▶  $P_2$ 's severity
- ▶  $P_2$ 's infection duration

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

## Synergistic interaction

increases →



Presence of  $P_1$  :

- ▶  $P_2$ 's growth
- ▶  $P_2$ 's severity
- ▶  $P_2$ 's infection duration


← decreases

Co-circulation of many pathogens in the environnement and inside the human host.

→ **possibilities of interactions during human infection**

## Synergistic interaction

increases →




Presence of  $P_1$  :

- ▶  $P_2$ 's growth
- ▶  $P_2$ 's severity
- ▶  $P_2$ 's infection duration

## Antagonistic interaction

← decreases





Often suggested in the literature

*Bosch et al. 2013, Mina et al. 2014, Opatowski et al. 2018*

## First evidence: 1918 influenza pandemic

- ▶ Majority of deaths caused by secondary bacterial infections
  - ▶ *Streptococcus pneumoniae*, *Staphylococcus aureus*,  
*Haemophilus influenzae*

*Brundage et al. 2008, Morens et al. 2008, Joseph et al. 2013*

- Explore, through modelling, the impact of interactions at the individual scale on infection dynamics at the population scale



- Explore, through modelling, the impact of interactions at the individual scale on infection dynamics at the population scale



- Assess, through a simulation study, the ability of several methods to detect an interaction from ecological data



# **SimFI: Simulator of Flu in Interaction**

**Objectif** Stochastic and realistic simulator of the co-circulation of influenza and another pathogen in a virtual human population.

**Objectif** Stochastic and realistic simulator of the co-circulation of influenza and another pathogen in a virtual human population.


→ **Agent-based model**

- ▶ Precise description at the individual scale
- ▶ Emergence of a global dynamic at the population scale
- ▶ Study the links between these two scales

Available on the NetLogo platform

*Wilensky et al. 1999*






<i>Risk</i>	<i>acquisition</i>	<i>transmission</i>	<i>infection</i>
<div> <div>Second pathogen</div> <div>Influenza</div> </div>	<b>Susceptible</b>	<b>Infected</b>	<b>Colonised</b>
<b>Susceptible</b>	$\beta$ <i>no interaction</i>	$\beta$ <i>no interaction</i>	$p$ <i>no interaction</i>
<b>Infected</b>	$\beta \times A$ <i>acquisition mechanism</i>	$\beta \times \Theta$ <i>transmission mechanism</i>	$p \times \Pi$ <i>pathogénicité mechanism</i>



<i>Risk</i>	<i>acquisition</i>	<i>transmission</i>	<i>infection</i>
<div> <div>Second pathogen</div> <div>Influenza</div> </div>	<b>Susceptible</b>	<b>Infected</b>	<b>Colonised</b>
<b>Susceptible</b>	$\beta$ <i>no interaction</i>	$\beta$ <i>no interaction</i>	$p$ <i>no interaction</i>
<b>Infected</b>	$\beta \times A$ <i>acquisition mechanism</i>	$\beta \times \Theta$ <i>transmission mechanism</i>	$p \times \Pi$ <i>pathogénicité mechanism</i>

## Interaction scenario

- ▶  $A = \Theta = \Pi = 1 \rightarrow$  baseline scenario, no interaction
- ▶  $A \in [2; 50]$  ;  $\Theta = \Pi = 1$
- ▶  $\Theta \in [2; 17]$  ;  $A = \Pi = 1$
- ▶  $\Pi \in [5; 50]$  ;  $A = \Theta = 1$



## Validation criteria

- ▶ Influenza *Sentinelles*
  - ▶ Annual number of cases
  - ▶ Epidemic onset
  - ▶ Peak timing
  - ▶ Epidemic duration
- ▶ Pneumococcal infections
  - ▶ Annual number of cases *Epibac*
  - ▶ Seasonality *Météo France*

## Detection of the interaction

Numerous studies but no gold standard


1. Regression models
2. Dynamical models


Numerous studies but no gold standard

1. Regression models
2. Dynamical models

**Are these methods adequate to detect such an interaction?**







$$F(I_t) = a + bG_{t+i} + c \cos \frac{2\pi t}{52} + d \sin \frac{2\pi t}{52} + \epsilon_t$$

$F$  transformation  
(linear, Poisson, or  
negative binomiale)

$I_t$  pneumococcal infections

$G_{t+i}$  influenza infections

$t$  week

$i$  time lag



$$F(I_t) = a + bG_{t+i} + c \cos \frac{2\pi t}{52} + d \sin \frac{2\pi t}{52} + \epsilon_t$$


→ 1 134 000 regressions

## Results comparison

Student t-test:  $t = \frac{\hat{b}}{SE_b}$

## Softwares and computing resources





$$\begin{cases} \frac{dS}{dt} = \lambda C(t) - \beta(t) \frac{S(t)C(t)}{N} \\ \frac{dC}{dt} = -\lambda C(t) + \beta(t) \frac{S(t)C(t)}{N} \\ \frac{dI}{dt} = p(t)C(t) \end{cases} \quad \begin{cases} \beta(t) = \beta_0(1 + \xi \times G(t)) \\ p(t) = p_0(1 + \pi \times G(t)) \end{cases}$$

4 parameters to estimate:  $\beta_0$ ,  $\xi$ ,  $p_0$  et  $\pi$ .


→ 36 000 estimations and their confidence intervals

## Likelihood maximisation

- ▶ Poisson likelihood
- ▶ Estimation: NSGA-2 algorithm *Deb et al. 2002*
- ▶ Confidence intervals : profiled likelihood on the OpenMOLE platform *Reuillon et al. 2015*

## Softwares and computing resources





# Conclusion

**SimFI model** Reconstruction of multi-scales dynamics, better understanding of the impact of interactions

**Interaction detection** Importance of the method, limits of the analysis of aggregated surveillance data

**SimFI model** Reconstruction of multi-scales dynamics, better understanding of the impact of interactions

**Interaction detection** Importance of the method, limits of the analysis of aggregated surveillance data

## Public Health applications

- ▶ Integrating ecological aspects of interactions
- ▶ Evaluate the global impact of public health policies
  - ▶ optimise drug use
  - ▶ fight globally against infectious diseases



**Thank you for your attention!**