

Quantifier rules: universal elimination

WEEK 7 . TOPIC INTRODUCTION

UNIVERSAL ELIMINATION

1. $\forall x \Phi(x)$
2. $\Phi[x/c]$ $E\forall 1$

The $\Phi[x/c]$ part means 'Take Φ and replace every instance of x with c '

For this rule, you can pick *any name you want* for c .


UNIVERSAL ELIMINATION

1. $\forall x \Phi(x)$
2. $\Phi[x/c]$ $E\forall 1$

The $\Phi[x/c]$ part means 'Take Φ and replace every instance of x with c '

E.g.:

1. $\forall x (Fx \rightarrow Gx)$:assumption
2. $Fa \rightarrow Ga$: $E\forall 1$



You must replace every instance of x with your new constant.

UNIVERSAL ELIMINATION

1. $\forall x \Phi(x)$
2. $\Phi[x/c]$ $E\forall 1$

The $\Phi[x/c]$ part means 'Take Φ and replace every instance of x with c '

E.g.:

1. $\forall x (Fx \rightarrow Gax)$:assumption
2. $Fa \rightarrow Gaa$: $E\forall 1$

UNIVERSAL ELIMINATION

1. $\forall x \Phi(x)$
2. $\Phi[x/c]$ $E\forall 1$

The $\Phi[x/c]$ part means 'Take Φ and replace every instance of x with c '

Check-in: Does this make sense?
Why can I replace a variable
with *any* constant I want?

A: Because the universal
quantifier is saying that for all
objects, such and such is true!