

# MASD

Lecture 2  
08.09.2022

François Lauze

Lecture will start at 10.15

# Objectives

After today's lecture, you should

- ▶ Know the definition of the derivative of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  and have an intuitive understanding of what it means (Sections 2.7-2.8).

# Objectives

After today's lecture, you should

- ▶ Know the definition of the derivative of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  and have an intuitive understanding of what it means (Sections 2.7-2.8).
- ▶ Know some differentiation rules (Sections 3.1 and 3.2). More rules and their use will follow next week.

# Limits of Functions - From Lecture 1

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

- ▶ What does the statement  $\lim_{x \rightarrow a} f(x) = L$  mean?

## Limits of Functions - From Lecture 1

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

- ▶ What does the statement  $\lim_{x \rightarrow a} f(x) = L$  mean?
- ▶ Let  $f$  be a function defined in some open interval containing  $a$  (except perhaps at  $a$ ).

# Limits of Functions - From Lecture 1

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .


- ▶ What does the statement  $\lim_{x \rightarrow a} f(x) = L$  mean?
- ▶ Let  $f$  be a function defined in some open interval containing  $a$  (except perhaps at  $a$ ).
- ▶ Pick any  $\varepsilon > 0$ .

# Limits of Functions - From Lecture 1

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

- ▶ What does the statement  $\lim_{x \rightarrow a} f(x) = L$  mean?
- ▶ Let  $f$  be a function defined in some open interval containing  $a$  (except perhaps at  $a$ ).
- ▶ Pick any  $\varepsilon > 0$ .
- ▶  $\lim_{x \rightarrow a} f(x) = L$  if there is  $\delta > 0$  such that for all  $x$

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$




# Limits of Functions - From Lecture 1

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

- ▶ What does the statement  $\lim_{x \rightarrow a} f(x) = L$  mean?
- ▶ Let  $f$  be a function defined in some open interval containing  $a$  (except perhaps at  $a$ ).
- ▶ Pick any  $\varepsilon > 0$ .
- ▶  $\lim_{x \rightarrow a} f(x) = L$  if there is  $\delta > 0$  such that for all  $x$

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$



- ▶  $f(x)$  can be made arbitrarily close to  $L$  if  $x$  is close enough to  $a$  but not equal to  $a$ .





## Continuous Functions - From Lecture 1

- ▶ A function  $f$  defined on an interval  $I$  (could be entire  $\mathbb{R}$ ) is *continuous* at  $a \in I \iff \lim_{x \rightarrow a} f(x) = f(a)$ . This implicitly requires that
  - ▶  $f(a)$  is defined ( $a$  is in the domain of  $f$ ),
  - ▶  $\lim_{x \rightarrow a} f(x)$  exists.



## Continuous Functions - From Lecture 1

- ▶ A function  $f$  defined on an interval  $I$  (could be entire  $\mathbb{R}$ ) is *continuous* at  $a \in I \iff \lim_{x \rightarrow a} f(x) = f(a)$ . This implicitly requires that
  - ▶  $f(a)$  is defined ( $a$  is in the domain of  $f$ ),
  - ▶  $\lim_{x \rightarrow a} f(x)$  exists.
- ▶ Remember that  $x$  can approach  $a$  from both the left and the right side and limits from left and from right both **must be equal** to  $f(a)$  (unless  $I$  is a closed interval with  $a$  being one of its endpoints - then only right limit at  $a$  or only left limit at  $b$  is needed).




# Continuous Functions - From Lecture 1

- ▶ A function  $f$  defined on an interval  $I$  (could be entire  $\mathbb{R}$ ) is *continuous* at  $a \in I \iff \lim_{x \rightarrow a} f(x) = f(a)$ . This implicitly requires that
  - ▶  $f(a)$  is defined ( $a$  is in the domain of  $f$ ),
  - ▶  $\lim_{x \rightarrow a} f(x)$  exists.
- ▶ Remember that  $x$  can approach  $a$  from both the left and the right side and limits from left and from right both **must be equal** to  $f(a)$  (unless  $I$  is a closed interval with  $a$  being one of its endpoints - then only right limit at  $a$  or only left limit at  $b$  is needed).




- ▶ The function  $f$  is *continuous* if it is continuous at every point  $a$  of its domain.

# Tangents Revisited



- ▶ When  $x$  approaches  $a$  then  $Q = (x, f(x))$  approaches  $P = (a, f(a))$  on  $C$  defined by  $y = f(x)$ . The slope  $m_{PQ}$  of the secant line approaches the slope of the tangent line.


# Tangents Revisited



- ▶ When  $x$  approaches  $a$  then  $Q = (x, f(x))$  approaches  $P = (a, f(a))$  on  $C$  defined by  $y = f(x)$ . The slope  $m_{PQ}$  of the secant line approaches the slope of the tangent line.
- ▶ The slope of the secant line is given by

$$m_{PQ} = \frac{f(x) - f(a)}{x - a}$$

# Tangents Revisited



- ▶ When  $x$  approaches  $a$  then  $Q = (x, f(x))$  approaches  $P = (a, f(a))$  on  $C$  defined by  $y = f(x)$ . The slope  $m_{PQ}$  of the secant line approaches the slope of the tangent line.
- ▶ The slope of the secant line is given by

$$m_{PQ} = \frac{f(x) - f(a)}{x - a}$$

- ▶ The slope of the tangent line of  $C$  at  $P$  is given by

$$m_P = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

# Tangents Revisited

- Consider the function  $f(x) = x^2$ . What is the tangent slope at point  $P = (2, 4)$ ?

$$m_P = \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x - 2)(x + 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 4$$

# Tangents Revisited

- Let  $h = x - a$ . Then  $m_{PQ}$  and  $m_P$  can be written as

$$m_{PQ} = \frac{f(x) - f(a)}{x - a} = \frac{f(a + h) - f(a)}{h}$$

$$m_P = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$



# Tangents Revisited

- Let  $h = x - a$ . Then  $m_{PQ}$  and  $m_P$  can be written as

$$m_{PQ} = \frac{f(x) - f(a)}{x - a} = \frac{f(a + h) - f(a)}{h}$$

$$m_P = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

- Consider the function  $f(x) = \frac{3}{x}$ . What is the tangent slope at point  $P = (3, 1)$ ?

$$m_P = \lim_{h \rightarrow 0} \frac{f(3 + h) - f(3)}{h} = \lim_{h \rightarrow 0} \frac{\frac{3}{3+h} - 1}{h} = \lim_{h \rightarrow 0} \frac{\frac{3 - (3+h)}{3+h}}{h} =$$

$$\lim_{h \rightarrow 0} \frac{-h}{h(3 + h)} = \lim_{h \rightarrow 0} -\frac{1}{3 + h} = -\frac{1}{3}$$

## Velocity Revisited

- ▶ What is the velocity of an object at a specific time?

## Velocity Revisited

- ▶ What is the velocity of an object at a specific time?
- ▶ Assume that an object starts in the origo and moves along the  $x$ -axis. Its position (in meters)  $s$  at time  $t$  (in seconds) is given by

$$s(t) = 3t^2, t \geq 0$$

## Velocity Revisited

- ▶ What is the velocity of an object at a specific time?
- ▶ Assume that an object starts in the origo and moves along the  $x$ -axis. Its position (in meters)  $s$  at time  $t$  (in seconds) is given by

$$s(t) = 3t^2, t \geq 0$$

- ▶ What is the average velocity of the object between times  $t_0$  and  $t$ ?

$$\text{av. velocity} = \frac{\text{distance travelled}}{\text{time elapsed}} = \frac{s(t) - s(t_0)}{t - t_0} = 3 \frac{t^2 - t_0^2}{t - t_0}$$

## Velocity Revisited

- ▶ What is the velocity of an object at a specific time?
- ▶ Assume that an object starts in the origo and moves along the  $x$ -axis. Its position (in meters)  $s$  at time  $t$  (in seconds) is given by

$$s(t) = 3t^2, t \geq 0$$


- ▶ What is the average velocity of the object between times  $t_0$  and  $t$ ?

$$\text{av. velocity} = \frac{\text{distance travelled}}{\text{time elapsed}} = \frac{s(t) - s(t_0)}{t - t_0} = 3 \frac{t^2 - t_0^2}{t - t_0}$$



- ▶ What is the instantaneous velocity at time  $t_0$ ?

$$\text{inst. velocity} = \lim_{t \rightarrow t_0} \frac{s(t) - s(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{s(t_0 + h) - s(t_0)}{h}$$




# Velocity Revisited



# Velocity Revisited







# Velocity Revisited





# Velocity Revisited



- Limits play central role in differentiation - main concept in calculus.

# Derivatives

- ▶ Computing tangents of a curve at a given position or velocity of a moving object at a given time are closely related. They both amount to the determination of the *rate of change* of a function.

# Derivatives

- ▶ Computing tangents of a curve at a given position or velocity of a moving object at a given time are closely related. They both amount to the determination of the *rate of change* of a function.
- ▶ Many problems in sciences and engineering require calculations of rates of change.

# Derivatives

- ▶ Computing tangents of a curve at a given position or velocity of a moving object at a given time are closely related. They both amount to the determination of the *rate of change* of a function.
- ▶ Many problems in sciences and engineering require calculations of rates of change.
- ▶ Given some function  $f$ , its *derivative at some  $a$*  (da. afledte i  $a$ ) in the domain of  $f$  is

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

$f'(a)$  is undefined if the limit does not exist.

# Derivatives

- ▶ Computing tangents of a curve at a given position or velocity of a moving object at a given time are closely related. They both amount to the determination of the *rate of change* of a function.
- ▶ Many problems in sciences and engineering require calculations of rates of change.
- ▶ Given some function  $f$ , its *derivative at some  $a$*  (da. afledte i  $a$ ) in the domain of  $f$  is

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

$f'(a)$  is undefined if the limit does not exist.

- ▶ Derivative  $f'(a)$  at  $a$  is a number.

# Derivatives as Functions

- ▶ Given a function  $f(x)$ , its *derivative function* (da. afledt funktion)  $f'$  is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

where  $x$  is a variable.

# Derivatives as Functions

- ▶ Given a function  $f(x)$ , its *derivative function* (da. afledt funktion)  $f'$  is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

where  $x$  is a variable.

- ▶ The domain of  $f'$  is the set  $\{x | f'(x) \text{ exists}\}$ . **Take poll 9 here**

# Derivatives as Functions

- ▶ Given a function  $f(x)$ , its *derivative function* (derivative function)  $f'$  is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

where  $x$  is a variable.

- ▶ The domain of  $f'$  is the set  $\{x | f'(x) \text{ exists}\}$ . **Take poll 9 here**
- ▶ Common alternative notations (assuming that  $y = f(x)$ ):

$$f'(x) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx}f(x) = Df(x) = D_x f(x)$$



# Derivatives as Functions

- ▶ Given a function  $f(x)$ , its *derivative function* (derivative function)  $f'$  is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

where  $x$  is a variable.

- ▶ The domain of  $f'$  is the set  $\{x | f'(x) \text{ exists}\}$ . **Take poll 9 here**
- ▶ Common alternative notations (assuming that  $y = f(x)$ ):


$$f'(x) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx}f(x) = Df(x) = D_x f(x)$$

- ▶ Leibnitz notation:  $\frac{dy}{dx}$

# Differentiability

- The *derivative* of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  at some  $x_0$  in the domain of  $f$  is the limit


$$\frac{df}{dx}(x_0) = f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$



# Differentiability

- The *derivative* of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  at some  $x_0$  in the domain of  $f$  is the limit

$$\frac{df}{dx}(x_0) = f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$




- A function  $f$  is *differentiable at some*  $x_0$  in the domain of  $f$  if  $f'(a)$  exists.

# Differentiability



- ▶ The *derivative* of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  at some  $x_0$  in the domain of  $f$  is the limit

$$\frac{df}{dx}(x_0) = f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$





- ▶ A function  $f$  is *differentiable at some*  $x_0$  in the domain of  $f$  if  $f'(a)$  exists.
- ▶ When the derivative  $f'$  of  $f$  exists in every point  $x_0$  in an open interval  $]a, b[ \in \mathbb{R}$ , then  $f$  is *differentiable* in  $]a, b[$  ( $a$  could be  $-\infty$  and/or  $b$  could be  $\infty$ ).

# Does a derivative always exist?



- If  $f$  is differentiable at  $x_0$  then it is continuous in  $x_0$ .

# Does a derivative always exist?



- ▶ If  $f$  is differentiable at  $x_0$  then it is continuous in  $x_0$ .
- ▶ There are continuous functions that are not differentiable (e.g.  $f(x) = |x|$  is not differentiable at 0).

Proof:  $f$  is differentiable in  $a \implies f$  is continuous in  $a$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (f(x) + f(a) - f(a)) =$$

Proof:  $f$  is differentiable in  $a \implies f$  is continuous in  $a$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (f(x) + f(a) - f(a)) =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} [f(x) - f(a)] =$$



Proof:  $f$  is differentiable in  $a \implies f$  is continuous in  $a$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (f(x) + f(a) - f(a)) =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} [f(x) - f(a)] =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} \left[ \frac{f(x) - f(a)}{x - a} (x - a) \right] =$$

Proof:  $f$  is differentiable in  $a \implies f$  is continuous in  $a$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (f(x) + f(a) - f(a)) =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} [f(x) - f(a)] =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} \left[ \frac{f(x) - f(a)}{x - a} (x - a) \right] =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \lim_{x \rightarrow a} (x - a) =$$

Proof:  $f$  is differentiable in  $a \implies f$  is continuous in  $a$

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (f(x) + f(a) - f(a)) =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} [f(x) - f(a)] =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} \left[ \frac{f(x) - f(a)}{x - a} (x - a) \right] =$$

$$\lim_{x \rightarrow a} f(a) + \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \lim_{x \rightarrow a} (x - a) =$$


$$\lim_{x \rightarrow a} f(a) + f'(a) * 0 = f(a)$$

## Higher Order Derivatives

- ▶  $f''$  is the *second order* derivative of  $f$  and the derivative of  $f'$ . It can be interpreted as the rate of change of the slope of the curve defined by  $f$ .

## Higher Order Derivatives

- ▶  $f''$  is the *second order* derivative of  $f$  and the derivative of  $f'$ . It can be interpreted as the rate of change of the slope of the curve defined by  $f$ .
- ▶ If  $f$  is the distance covered by a moving object as the function of time,  $f'$  is its instantaneous velocity as the function of time (or the rate of position change), and  $f''$  is its instantaneous acceleration as the function of time (or the rate of velocity change).



# Derivatives of Constant Functions

- **Constant Rule:** Let  $f(x) = c$  where  $c$  is an arbitrary constant. Then  $f'(x) = 0$ . Let  $a \in \mathbb{R}$ .

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \frac{c - c}{h} = 0$$

# Derivatives of Power Functions

- **Power Rule:** Let  $f(x) = x^n$  where  $n$  is a positive integer. Then  $f'(x) = nx^{n-1}$ . Let  $a \in \mathbb{R}$

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} =$$

# Derivatives of Power Functions

- **Power Rule:** Let  $f(x) = x^n$  where  $n$  is a positive integer. Then  $f'(x) = nx^{n-1}$ . Let  $a \in \mathbb{R}$

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} =$$

$$\lim_{x \rightarrow a} \frac{(x - a)(x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1})}{x - a} =$$



# Derivatives of Power Functions

- **Power Rule:** Let  $f(x) = x^n$  where  $n$  is a positive integer. Then  $f'(x) = nx^{n-1}$ . Let  $a \in \mathbb{R}$

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} =$$

$$\lim_{x \rightarrow a} \frac{(x - a)(x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1})}{x - a} =$$

$$\lim_{x \rightarrow a} (x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1}) =$$

# Derivatives of Power Functions

- **Power Rule:** Let  $f(x) = x^n$  where  $n$  is a positive integer. Then  $f'(x) = nx^{n-1}$ . Let  $a \in \mathbb{R}$

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} =$$

$$\lim_{x \rightarrow a} \frac{(x - a)(x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1})}{x - a} =$$

$$\lim_{x \rightarrow a} (x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1}) =$$

$$a^{n-1} + a^{n-2}a + \dots + aa^{n-2} + a^{n-1} = na^{n-1}$$

# Derivatives of Power Functions

- **Power Rule:** Let  $f(x) = x^n$  where  $n$  is a positive integer. Then  $f'(x) = nx^{n-1}$ . Let  $a \in \mathbb{R}$

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} =$$

$$\lim_{x \rightarrow a} \frac{(x - a)(x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1})}{x - a} =$$

$$\lim_{x \rightarrow a} (x^{n-1} + x^{n-2}a + \dots + xa^{n-2} + a^{n-1}) =$$

$$a^{n-1} + a^{n-2}a + \dots + aa^{n-2} + a^{n-1} = na^{n-1}$$

- Generalizes to  $n$  being any real number (proof in Section 3.6).

## New Derivatives from Old Derivatives

- **Constant Multiple Rule:** Let  $g(x) = cf(x)$  where  $c$  is a constant and  $f$  is a differentiable function.

$$\begin{aligned} g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} = \\ &= \lim_{h \rightarrow 0} c \frac{f(x+h) - f(x)}{h} = c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = cf'(x) \end{aligned}$$

where the second last equality follows from one of the limit laws.

## Sum Rule

- Let  $F(x) = f(x) + g(x)$  where both  $f$  and  $g$  are differentiable.

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} = \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right] = \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \\ &= f'(x) + g'(x) \end{aligned}$$

where the second last equality follows from one of the limit laws.

## Sum Rule

- Let  $F(x) = f(x) + g(x)$  where both  $f$  and  $g$  are differentiable.

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} = \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right] = \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \\ &= f'(x) + g'(x) \end{aligned}$$

where the second last equality follows from one of the limit laws.

- **Difference Rule:** similar.

## Sum Rule

- ▶ Let  $F(x) = f(x) + g(x)$  where both  $f$  and  $g$  are differentiable.

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} = \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right] = \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \\ &= f'(x) + g'(x) \end{aligned}$$

where the second last equality follows from one of the limit laws.

- ▶ **Difference Rule:** similar.
- ▶ These rules can be used to differentiate any polynomial.

# Differentiating Polynomials

► Let  $f(x) = x^7 - 4x^5 + 13x^4 - x + 19$

$$f'(x) = 7x^6 - 4*5x^4 + 13*4x^3 - 1x^0 + 0 = 7x^6 - 20x^4 + 52x^3 - 1$$



# Exponential Functions

► Let  $f(x) = b^x$ ..

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} =$$

$$\lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} = \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h} = b^x f'(0)$$

# Exponential Functions

- ▶ Let  $f(x) = b^x$ ..

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} = \\ \lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} &= \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h} = b^x f'(0) \end{aligned}$$

- ▶ This shows that if  $f(x) = b^x$  is differentiable at 0 then it is differentiable everywhere.

# Exponential Functions

- ▶ Let  $f(x) = b^x$ ..

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} =$$

$$\lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} = \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h} = b^x f'(0)$$

- ▶ This shows that if  $f(x) = b^x$  is differentiable at 0 then it is differentiable everywhere.
- ▶ If  $f(x) = 2^x$  then  $f'(x) = f'(0)2^x$  and  $f'(0)=0.693...$

# Exponential Functions

- ▶ Let  $f(x) = b^x$ ..

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} =$$

$$\lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} = \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h} = b^x f'(0)$$

- ▶ This shows that if  $f(x) = b^x$  is differentiable at 0 then it is differentiable everywhere.
- ▶ If  $f(x) = 2^x$  then  $f'(x) = f'(0)2^x$  and  $f'(0)=0.693...$
- ▶ If  $f(x) = 3^x$  then  $f'(x) = f'(0)3^x$  and  $f'(0)=1.099...$

# Exponential Functions

- ▶ Let  $f(x) = b^x$ ..

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} = \\ \lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} &= \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h} = b^x f'(0) \end{aligned}$$

- ▶ This shows that if  $f(x) = b^x$  is differentiable at 0 then it is differentiable everywhere.
- ▶ If  $f(x) = 2^x$  then  $f'(x) = f'(0)2^x$  and  $f'(0)=0.693...$
- ▶ If  $f(x) = 3^x$  then  $f'(x) = f'(0)3^x$  and  $f'(0)=1.099...$
- ▶ Determine constant  $e$  such that the exponential function  $f(x) = e^x$  satisfies  $f'(0) = 1$ . This happens for  $e=2.71828...$   
Then  $f'(x) = f(x)$ .

## Product Rule (da. Produktregeln)

- ▶ Let  $F(x) = f(x)g(x)$  where both  $f$  and  $g$  are differentiable.  
Then

$$F'(x) = f(x)g'(x) + f'(x)g(x)$$

## Product Rule (da. Produktregeln)

- ▶ Let  $F(x) = f(x)g(x)$  where both  $f$  and  $g$  are differentiable. Then

$$F'(x) = f(x)g'(x) + f'(x)g(x)$$

- ▶ Let  $F(x) = x^3 \cos x$ . Hence,  $f(x) = x^3$  and  $g(x) = \cos x$

$$F'(x) = x^3(-\sin x) + 3x^2 \cos x = -x^3 \sin x + 3x^2 \cos x$$

# Product Rule - Proof

$$\begin{aligned}F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} = \\&\lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} = \\&\lim_{h \rightarrow 0} \frac{f(x+h)[g(x+h) - g(x)]}{h} + \lim_{h \rightarrow 0} \frac{g(x)[f(x+h) - f(x)]}{h} = \\&\lim_{h \rightarrow 0} \left[ f(x+h) \frac{g(x+h) - g(x)}{h} \right] + \lim_{h \rightarrow 0} \left[ g(x) \frac{f(x+h) - f(x)}{h} \right] = \\&\lim_{h \rightarrow 0} f(x+h) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} + \lim_{h \rightarrow 0} g(x) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \\&f(x)g'(x) + g(x)f'(x)\end{aligned}$$



## Quotient Rule (da. Brøkreglen)

- Let  $F(x) = \frac{f(x)}{g(x)}$  where both  $f$  and  $g$  are differentiable.

$$F'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$$

## Quotient Rule (da. Brøkreglen)

- Let  $F(x) = \frac{f(x)}{g(x)}$  where both  $f$  and  $g$  are differentiable.

$$F'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$$

- Let  $F(x) = \frac{2-x^2}{2x^3+x+3}$ . Hence  $f(x) = 2 - x^2$  and  $g(x) = 2x^3 + x + 3$ .

$$F'(x) = \frac{-2x(2x^3 + x + 3) - (2 - x^2)(6x^2 + 1)}{(2x^3 + x + 3)^2} =$$

$$\frac{2x^4 - 13x^2 - 6x - 2}{(2x^3 + x + 3)^2}$$

# Quotient Rule - Proof

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} =$$

# Quotient Rule - Proof

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} =$$

# Quotient Rule - Proof

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} = \\ &\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\ &\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \end{aligned}$$

## Quotient Rule - Proof

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} = \\ &\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\ &\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\ &\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{h} \right\} = \end{aligned}$$

## Quotient Rule - Proof

$$\begin{aligned}F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{h} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ \frac{f(x+h)g(x) - f(x)g(x)}{h} + \frac{f(x)g(x) - f(x)g(x+h)}{h} \right] \right\} =\end{aligned}$$

## Quotient Rule - Proof

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{h} \right\} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ \frac{f(x+h)g(x) - f(x)g(x)}{h} + \frac{f(x)g(x) - f(x)g(x+h)}{h} \right] \right\} =$$

$$\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ g(x) \frac{f(x+h) - f(x)}{h} - f(x) \frac{g(x+h) - g(x)}{h} \right] \right\} =$$



## Quotient Rule - Proof

$$\begin{aligned}F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{h} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ \frac{f(x+h)g(x) - f(x)g(x)}{h} + \frac{f(x)g(x) - f(x)g(x+h)}{h} \right] \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ g(x) \frac{f(x+h) - f(x)}{h} - f(x) \frac{g(x+h) - g(x)}{h} \right] \right\} = \\&\lim_{h \rightarrow 0} \frac{1}{g(x+h)g(x)} \left[ \lim_{h \rightarrow 0} g(x) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} - \lim_{h \rightarrow 0} f(x) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \right] =\end{aligned}$$

## Quotient Rule - Proof

$$\begin{aligned}F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{f(x+h)}{g(x+h)} - \frac{f(x)}{g(x)}}{h} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{h} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{g(x+h)g(x)} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \frac{f(x+h)g(x) - f(x)g(x) + f(x)g(x) - f(x)g(x+h)}{h} \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ \frac{f(x+h)g(x) - f(x)g(x)}{h} + \frac{f(x)g(x) - f(x)g(x+h)}{h} \right] \right\} = \\&\lim_{h \rightarrow 0} \left\{ \frac{1}{g(x+h)g(x)} \left[ g(x) \frac{f(x+h) - f(x)}{h} - f(x) \frac{g(x+h) - g(x)}{h} \right] \right\} = \\&\lim_{h \rightarrow 0} \frac{1}{g(x+h)g(x)} \left[ \lim_{h \rightarrow 0} g(x) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} - \lim_{h \rightarrow 0} f(x) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \right] = \\&\frac{g(x)f'(x) - f(x)g'(x)}{\lim_{h \rightarrow 0} g(x+h) \lim_{h \rightarrow 0} g(x)} = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}\end{aligned}$$

# Summary

By now, you should be familiar with

- ▶ the definition of function limits,
- ▶ the most familiar rules for function limits, and be able to use them for deriving limits of new functions,
- ▶ the definition of continuity, and be able to use it to prove continuity of other functions,
- ▶ the definition of the derivative of a function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,
- ▶ basic differentiation rules.