

MASD

Lecture 3
13.09.2022

François Lauze (material from Pawel Winter and
others)

Lecture will start at 8.15

Objectives

We cover sections 3.4-3.6 and 3.10

- ▶ Differentiation of composite functions: Chain rule
- ▶ Implicit differentiation
- ▶ Differentiation of inverse functions
- ▶ Differentiation of logarithmic functions
- ▶ Logarithmic differentiation
- ▶ Linear approximations of functions
- ▶ Differentials

Chain Rule (da. Kædereglen)

- ▶ Assume that the function g is differentiable at x and the function f is differentiable at $g(x)$.

Chain Rule (da. Kædereglen)

- ▶ Assume that the function g is differentiable at x and the function f is differentiable at $g(x)$.
- ▶ Consider a composite function $F = f \circ g$ defined by $F(x) = f(g(x))$.

Chain Rule (da. Kædereglen)

- ▶ Assume that the function g is differentiable at x and the function f is differentiable at $g(x)$.
- ▶ Consider a composite function $F = f \circ g$ defined by $F(x) = f(g(x))$.
- ▶ F is differentiable at x and $F'(x) = f'(g(x))g'(x)$.

Chain Rule (da. Kædereglen)

- ▶ Assume that the function g is differentiable at x and the function f is differentiable at $g(x)$.
- ▶ Consider a composite function $F = f \circ g$ defined by $F(x) = f(g(x))$.
- ▶ F is differentiable at x and $F'(x) = f'(g(x))g'(x)$.
- ▶ Let $u = g(x)$ and $y = f(u)$. Then using Leibnitz notation

$$F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$
- ▶ Let $y = f(g(x)) = f(u) = u^3$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$
- ▶ Let $y = f(g(x)) = f(u) = u^3$
- ▶ $f'(g(x)) = f'(u) = \frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$
- ▶ Let $y = f(g(x)) = f(u) = u^3$
- ▶ $f'(g(x)) = f'(u) = \frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$
- ▶ $g'(x) = \frac{du}{dx} = \frac{d}{dx}(4x^5 - 7x^3 + 14x^2 - 5) = 20x^4 - 21x^2 + 28x$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$
- ▶ Let $y = f(g(x)) = f(u) = u^3$
- ▶ $f'(g(x)) = f'(u) = \frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$
- ▶ $g'(x) = \frac{du}{dx} = \frac{d}{dx}(4x^5 - 7x^3 + 14x^2 - 5) = 20x^4 - 21x^2 + 28x$
- ▶ $F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = 3u^2(20x^4 - 21x^2 + 28x) =$

Chain Rule - Example I

- ▶ Let $F(x) = f(g(x)) = (4x^5 - 7x^3 + 14x^2 - 5)^3$. This is a composite function.
- ▶ Let $u = g(x) = 4x^5 - 7x^3 + 14x^2 - 5$
- ▶ Let $y = f(g(x)) = f(u) = u^3$
- ▶ $f'(g(x)) = f'(u) = \frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$
- ▶ $g'(x) = \frac{du}{dx} = \frac{d}{dx}(4x^5 - 7x^3 + 14x^2 - 5) = 20x^4 - 21x^2 + 28x$
- ▶ $F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = 3u^2(20x^4 - 21x^2 + 28x) =$
 $3(4x^5 - 7x^3 + 14x^2 - 5)^2(20x^4 - 21x^2 + 28x)$

Chain Rule - Example II

- ▶ Let $F(x) = \sqrt{3x^2 - 7x + 12}$. This is a composite function.

Chain Rule - Example II

- ▶ Let $F(x) = \sqrt{3x^2 - 7x + 12}$. This is a composite function.
- ▶ Let $y = \sqrt{u}$ and $u = 3x^2 - 7x + 12$

Chain Rule - Example II

- ▶ Let $F(x) = \sqrt{3x^2 - 7x + 12}$. This is a composite function.
- ▶ Let $y = \sqrt{u}$ and $u = 3x^2 - 7x + 12$
- ▶ $\frac{dy}{du} = \frac{d}{du}(\sqrt{u}) = \frac{d}{du}(u^{\frac{1}{2}}) = \frac{1}{2}u^{-\frac{1}{2}} = \frac{1}{2\sqrt{u}}$

Chain Rule - Example II

- ▶ Let $F(x) = \sqrt{3x^2 - 7x + 12}$. This is a composite function.
- ▶ Let $y = \sqrt{u}$ and $u = 3x^2 - 7x + 12$
- ▶ $\frac{dy}{du} = \frac{d}{du}(\sqrt{u}) = \frac{d}{du}(u^{\frac{1}{2}}) = \frac{1}{2}u^{-\frac{1}{2}} = \frac{1}{2\sqrt{u}}$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(3x^2 - 7x + 12) = 6x - 7$

$$F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{2\sqrt{u}}(6x - 7) = \frac{6x - 7}{2\sqrt{3x^2 - 7x + 12}}$$

Chain Rule - Example III

- ▶ Let $F(x) = \sin(4x)$. This is a composite function.

Chain Rule - Example III

- ▶ Let $F(x) = \sin(4x)$. This is a composite function.
- ▶ Let $y = \sin(u)$ and $u = 4x$

Chain Rule - Example III

- ▶ Let $F(x) = \sin(4x)$. This is a composite function.
- ▶ Let $y = \sin(u)$ and $u = 4x$
- ▶ $\frac{dy}{du} = \frac{d}{du}(\sin(u)) = \cos(u)$

Chain Rule - Example III

- ▶ Let $F(x) = \sin(4x)$. This is a composite function.
- ▶ Let $y = \sin(u)$ and $u = 4x$
- ▶ $\frac{dy}{du} = \frac{d}{du}(\sin(u)) = \cos(u)$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(4x) = 4$

$$F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \cos(u) * 4 = 4 \cos(4x)$$

Chain Rule - Example IV

- ▶ Let $F(x) = \cos^3(x)$. This is a composite function.

Chain Rule - Example IV

- ▶ Let $F(x) = \cos^3(x)$. This is a composite function.
- ▶ Let $y = u^3$ and $u = \cos(x)$

Chain Rule - Example IV

- ▶ Let $F(x) = \cos^3(x)$. This is a composite function.
- ▶ Let $y = u^3$ and $u = \cos(x)$
- ▶ $\frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$

Chain Rule - Example IV

- ▶ Let $F(x) = \cos^3(x)$. This is a composite function.
- ▶ Let $y = u^3$ and $u = \cos(x)$
- ▶ $\frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(\cos(x)) = -\sin(x)$

Chain Rule - Example IV

- ▶ Let $F(x) = \cos^3(x)$. This is a composite function.
- ▶ Let $y = u^3$ and $u = \cos(x)$
- ▶ $\frac{dy}{du} = \frac{d}{du}(u^3) = 3u^2$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(\cos(x)) = -\sin(x)$
- ▶ $F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = 3u^2(-\sin(x)) = -3\cos^2(x)\sin(x)$

Chain Rule - Example V

- ▶ Let $F(x) = e^{-x}$. This is a composite function.

Chain Rule - Example V

- ▶ Let $F(x) = e^{-x}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = -x$

Chain Rule - Example V

- ▶ Let $F(x) = e^{-x}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = -x$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$

Chain Rule - Example V

- ▶ Let $F(x) = e^{-x}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = -x$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(-x) = -1$

Chain Rule - Example V

- ▶ Let $F(x) = e^{-x}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = -x$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(-x) = -1$
- ▶ $F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = e^u(-1) = -e^{-x}$

Chain Rule - Example VI

- ▶ Let $F(x) = e^{4x^3-6x+1}$. This is a composite function.

Chain Rule - Example VI

- ▶ Let $F(x) = e^{4x^3-6x+1}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = 4x^3 - 6x + 1$

Chain Rule - Example VI

- ▶ Let $F(x) = e^{4x^3-6x+1}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = 4x^3 - 6x + 1$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$

Chain Rule - Example VI

- ▶ Let $F(x) = e^{4x^3-6x+1}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = 4x^3 - 6x + 1$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(4x^3 - 6x + 1) = 12x^2 - 6$

Chain Rule - Example VI

- ▶ Let $F(x) = e^{4x^3-6x+1}$. This is a composite function.
- ▶ Let $y = e^u$ and $u = 4x^3 - 6x + 1$
- ▶ $\frac{dy}{du} = \frac{d}{du}(e^u) = e^u$
- ▶ $\frac{du}{dx} = \frac{d}{dx}(4x^3 - 6x + 1) = 12x^2 - 6$
- ▶ $F'(x) = \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = e^u(12x^2 - 6) = (12x^2 - 6)e^{4x^3-6x+1}$

Chain Rule - Example VII

- ▶ Determine $\frac{d}{dx}(b^x)$

Chain Rule - Example VII

- ▶ Determine $\frac{d}{dx}(b^x)$
- ▶ We know that $b = e^{\ln b}$ for any $b > 0$. Therefore

$$b^x = (e^{\ln b})^x = e^{(\ln b)x}$$

Chain Rule - Example VII

- ▶ Determine $\frac{d}{dx}(b^x)$
- ▶ We know that $b = e^{\ln b}$ for any $b > 0$. Therefore

$$b^x = (e^{\ln b})^x = e^{(\ln b)x}$$

- ▶ Applying chain rule yields

$$\frac{d}{dx}(b^x) = \frac{d}{dx}(e^{(\ln b)x}) = e^{(\ln b)x} \frac{d}{dx}[(\ln b)x] = e^{(\ln b)x} \ln b = b^x \ln b$$

where the 2. equality is due to the chain rule while the 3. equality holds because $\ln b$ is a constant.

Chain Rule - Proof Attempt

- Definition of the derivative of $F = f \circ g$ at some a where g is differentiable at a and f is differentiable at $g(a)$:

$$F'(a) = (f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a}$$

Chain Rule - Proof Attempt

- Definition of the derivative of $F = f \circ g$ at some a where g is differentiable at a and f is differentiable at $g(a)$:

$$F'(a) = (f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a}$$

- If $g(x) \neq g(a)$ for any x near a , then

$$\lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} \left[\frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \frac{g(x) - g(a)}{x - a} \right] =$$

$$\lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a)$$

Chain Rule - Proof Attempt

- Definition of the derivative of $F = f \circ g$ at some a where g is differentiable at a and f is differentiable at $g(a)$:

$$F'(a) = (f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a}$$

- If $g(x) \neq g(a)$ for any x near a , then

$$\lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} \left[\frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \frac{g(x) - g(a)}{x - a} \right] =$$

$$\lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a)$$

- Unfortunately, there are functions such that $g(x) - g(a) = 0$ when x is arbitrarily close to a . For example $g(x) = x^2 \sin\left(\frac{1}{x}\right)$ when x is close to 0.

Chain Rule - Proof

► Let

$$Q(g(x)) = \begin{cases} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} & \text{if } g(x) \neq g(a) \\ f'(g(a)) & \text{if } g(x) = g(a) \end{cases}$$

Chain Rule - Proof

► Let

$$Q(g(x)) = \begin{cases} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} & \text{if } g(x) \neq g(a) \\ f'(g(a)) & \text{if } g(x) = g(a) \end{cases}$$

► If $g(x) \neq g(a)$, then $Q(g(x)) \frac{g(x) - g(a)}{x - a} = \frac{f(g(x)) - f(g(a))}{x - a}$

Chain Rule - Proof

► Let

$$Q(g(x)) = \begin{cases} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} & \text{if } g(x) \neq g(a) \\ f'(g(a)) & \text{if } g(x) = g(a) \end{cases}$$

► If $g(x) \neq g(a)$, then $Q(g(x)) \frac{g(x) - g(a)}{x - a} = \frac{f(g(x)) - f(g(a))}{x - a}$

► If $g(x) = g(a)$, $x \neq a$, then

$$Q(g(x)) \frac{g(x) - g(a)}{x - a} = f'(g(a)) * 0 = 0 = \frac{f(g(x)) - f(g(a))}{x - a}$$

Chain Rule - Proof

► Let

$$Q(g(x)) = \begin{cases} \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} & \text{if } g(x) \neq g(a) \\ f'(g(a)) & \text{if } g(x) = g(a) \end{cases}$$

► If $g(x) \neq g(a)$, then $Q(g(x)) \frac{g(x) - g(a)}{x - a} = \frac{f(g(x)) - f(g(a))}{x - a}$

► If $g(x) = g(a)$, $x \neq a$, then

$$Q(g(x)) \frac{g(x) - g(a)}{x - a} = f'(g(a)) * 0 = 0 = \frac{f(g(x)) - f(g(a))}{x - a}$$

$$\lim_{x \rightarrow a} \left[Q(g(x)) \frac{g(x) - g(a)}{x - a} \right] = \lim_{x \rightarrow a} Q(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} =$$

$$\left[\lim_{x \rightarrow a} Q(g(x)) \right] g'(a) \text{ (since } g \text{ is differentiable at } a \text{)}$$

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.
- ▶ f is differentiable at $g(a)$ by assumption.

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.
- ▶ f is differentiable at $g(a)$ by assumption.
- ▶ Therefore Q is continuous at $g(a)$, by definition of the derivative.

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.
- ▶ f is differentiable at $g(a)$ by assumption.
- ▶ Therefore Q is continuous at $g(a)$, by definition of the derivative.
- ▶ g is continuous at a because it is differentiable at a .

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.
- ▶ f is differentiable at $g(a)$ by assumption.
- ▶ Therefore Q is continuous at $g(a)$, by definition of the derivative.
- ▶ g is continuous at a because it is differentiable at a .
- ▶ Therefore $Q \circ g$ is continuous at a .

Chain Rule - Proof Continued

- ▶ We need to show that $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$
- ▶ Q is defined wherever f is defined.
- ▶ f is differentiable at $g(a)$ by assumption.
- ▶ Therefore Q is continuous at $g(a)$, by definition of the derivative.
- ▶ g is continuous at a because it is differentiable at a .
- ▶ Therefore $Q \circ g$ is continuous at a .
- ▶ So $\lim_{x \rightarrow a} Q(g(x)) = Q(g(a)) = f'(g(a))$.

Implicit Differentiation- Application of Chain Rule

- ▶ Equations with two variables x and y cannot always be formulated by functions with x as the only independent variable.

Implicit Differentiation- Application of Chain Rule

- ▶ Equations with two variables x and y cannot always be formulated by functions with x as the only independent variable.
- ▶ For example, $x^2 + y^2 = 25$ is an equation defining a circle with the center in origo and with radius 5. The equation **implicitly** defines functions $g(x)$ satisfying $x^2 + g(x)^2 = 25$.

Implicit Differentiation- Application of Chain Rule


- ▶ Equations with two variables x and y cannot always be formulated by functions with x as the only independent variable.
- ▶ For example, $x^2 + y^2 = 25$ is an equation defining a circle with the center in origo and with radius 5. The equation **implicitly** defines functions $g(x)$ satisfying $x^2 + g(x)^2 = 25$.
- ▶ $x^2 + y^2 = 25$ implicitly defines 2 functions

$$g_1(x) = \sqrt{25 - x^2} \text{ and } g_2(x) = -\sqrt{25 - x^2}$$

on the closed interval $[-5, 5]$. g_1 defines the function whose graph is the upper half-circle while g_2 defines the function whose graph is the lower half-circle.


Implicit Differentiation

- For the equation $x^3 + y^3 = 2xy$ it is however not that easy to identify implicitly defined functions.



Implicit Differentiation

- ▶ For the equation $x^3 + y^3 = 2xy$ it is however not that easy to identify implicitly defined functions.



- ▶ But such equations define curves. One may still need to determine derivatives (slopes of tangents) of differentiable functions without bothering about the functions itself.

Implicit Differentiation - Example I: $x^2 + y^2 = 25$

- ▶ Consider y as an (unknown) function of x .

Implicit Differentiation - Example 1: $x^2 + y^2 = 25$

- ▶ Consider y as an (unknown) function of x .
- ▶ Differentiate both sides with respect to x (applying the chain rule to y^2) and solve w.r.t. $\frac{dy}{dx}$.

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}(25) \iff \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \iff$$

$$2x + \frac{d}{dy}(y^2) \frac{dy}{dx} = 0 \iff 2x + 2y \frac{dy}{dx} = 0 \iff \frac{dy}{dx} = -\frac{x}{y}$$


Implicit Differentiation - Example 1: $x^2 + y^2 = 25$

- ▶ Consider y as an (unknown) function of x .
- ▶ Differentiate both sides with respect to x (applying the chain rule to y^2) and solve w.r.t. $\frac{dy}{dx}$.

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}(25) \iff \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 0 \iff$$

$$2x + \frac{d}{dy}(y^2) \frac{dy}{dx} = 0 \iff 2x + 2y \frac{dy}{dx} = 0 \iff \frac{dy}{dx} = -\frac{x}{y}$$

- ▶ At point $(3, 4)$, $x = 3$ and $y = 4$ and $\frac{dy}{dx} = -\frac{3}{4}$



Implicit Differentiation - Example II: $(x - y)^2 = x + y - 1$

$$\frac{d}{dx}[(x - y)^2] = \frac{d}{dx}[x + y - 1] \iff$$

Implicit Differentiation - Example II: $(x - y)^2 = x + y - 1$

$$\frac{d}{dx}[(x - y)^2] = \frac{d}{dx}[x + y - 1] \iff$$

$$2(x - y)\left(1 - \frac{dy}{dx}\right) = 1 + \frac{dy}{dx} \iff$$

Implicit Differentiation - Example II: $(x - y)^2 = x + y - 1$

$$\frac{d}{dx}[(x - y)^2] = \frac{d}{dx}[x + y - 1] \iff$$

$$2(x - y)\left(1 - \frac{dy}{dx}\right) = 1 + \frac{dy}{dx} \iff$$

$$2x - 2y - 2(x - y)\frac{dy}{dx} = 1 + \frac{dy}{dx} \iff$$

Implicit Differentiation - Example II: $(x - y)^2 = x + y - 1$

$$\frac{d}{dx}[(x - y)^2] = \frac{d}{dx}[x + y - 1] \iff$$

$$2(x - y)\left(1 - \frac{dy}{dx}\right) = 1 + \frac{dy}{dx} \iff$$

$$2x - 2y - 2(x - y)\frac{dy}{dx} = 1 + \frac{dy}{dx} \iff$$

$$2x - 2y - 1 = (2x - 2y + 1)\frac{dy}{dx} \iff$$

Implicit Differentiation - Example II: $(x - y)^2 = x + y - 1$

$$\frac{d}{dx}[(x - y)^2] = \frac{d}{dx}[x + y - 1] \iff$$

$$2(x - y)\left(1 - \frac{dy}{dx}\right) = 1 + \frac{dy}{dx} \iff$$

$$2x - 2y - 2(x - y)\frac{dy}{dx} = 1 + \frac{dy}{dx} \iff$$

$$2x - 2y - 1 = (2x - 2y + 1)\frac{dy}{dx} \iff$$

$$\frac{dx}{dy} = \frac{2x - 2y - 1}{2x - 2y + 1}$$

Implicit Differentiation - Example III: $x^3 + y^3 = 2xy$

- ▶ Changed notation.

Implicit Differentiation - Example III: $x^3 + y^3 = 2xy$

- ▶ Changed notation.
- ▶ Regard y as a function of x . Differentiate both sides with respect to x (applying chain rule to y^3 on the left side and product rule to the right side).

$$3x^2 + 3y^2y' = 2xy' + 2y$$

where $y' = \frac{dy}{dx}$.

Implicit Differentiation - Example III: $x^3 + y^3 = 2xy$

- ▶ Changed notation.
- ▶ Regard y as a function of x . Differentiate both sides with respect to x (applying chain rule to y^3 on the left side and product rule to the right side).

$$3x^2 + 3y^2y' = 2xy' + 2y$$

where $y' = \frac{dy}{dx}$.

- ▶ Solve for y'

$$3y^2y' - 2xy' = 2y - 3x^2 \iff (3y^2 - 2x)y' = 2y - 3x^2 \iff$$

$$y' = \frac{2y - 3x^2}{3y^2 - 2x} \text{ if } 3y^2 - 2x \neq 0$$

Implicit Differentiation - Example III: $x^3 + y^3 = 2xy$

- ▶ Changed notation.
- ▶ Regard y as a function of x . Differentiate both sides with respect to x (applying chain rule to y^3 on the left side and product rule to the right side).

$$3x^2 + 3y^2y' = 2xy' + 2y$$

where $y' = \frac{dy}{dx}$.

- ▶ Solve for y'

$$3y^2y' - 2xy' = 2y - 3x^2 \iff (3y^2 - 2x)y' = 2y - 3x^2 \iff$$

$$y' = \frac{2y - 3x^2}{3y^2 - 2x} \text{ if } 3y^2 - 2x \neq 0$$

- ▶ Point $(1, 1)$ satisfies the equation and therefore the derivative y' (slope of the tangent) in that point is

$$y' = \frac{2 * 1 - 3 * 1^2}{3 * 1^2 - 2 * 1} = -1$$

Derivatives of Inverse Functions

- ▶ A one-to-one function f has an inverse function g iff $g(f(x)) = x$ or $f(g(x)) = x$

Derivatives of Inverse Functions

- ▶ A one-to-one function f has an inverse function g iff $g(f(x)) = x$ or $f(g(x)) = x$
- ▶ An inverse of f is usually denoted by f^{-1}

Derivatives of Inverse Functions

- ▶ A one-to-one function f has an inverse function g iff $g(f(x)) = x$ or $f(g(x)) = x$
- ▶ An inverse of f is usually denoted by f^{-1}
- ▶ It can be shown (App. F) that the inverse function f^{-1} of one-to-one differentiable function f is differentiable except where f^{-1} has a vertical tangent.

Derivatives of Inverse Functions

- ▶ A one-to-one function f has an inverse function g iff $g(f(x)) = x$ or $f(g(x)) = x$
- ▶ An inverse of f is usually denoted by f^{-1}
- ▶ It can be shown (App. F) that the inverse function f^{-1} of one-to-one differentiable function f is differentiable except where f^{-1} has a vertical tangent.
- ▶ Let us apply implicit differentiation and chain rule to $f(g(x)) = x$

$$\frac{d}{dx}[f(g(x))] = \frac{d}{dx}[x] \iff f'(g(x))g'(x) = 1 \iff g'(x) = \frac{1}{f'(g(x))}$$

Derivatives of Inverse Functions

- ▶ A one-to-one function f has an inverse function g iff $g(f(x)) = x$ or $f(g(x)) = x$
- ▶ An inverse of f is usually denoted by f^{-1}
- ▶ It can be shown (App. F) that the inverse function f^{-1} of one-to-one differentiable function f is differentiable except where f^{-1} has a vertical tangent.
- ▶ Let us apply implicit differentiation and chain rule to $f(g(x)) = x$

$$\frac{d}{dx}[f(g(x))] = \frac{d}{dx}[x] \iff f'(g(x))g'(x) = 1 \iff g'(x) = \frac{1}{f'(g(x))}$$

- ▶ $f(x) = e^x$. We know that $f'(x) = e^x$ and $g(y) = \ln y$ is the inverse of f . Hence, $g'(y) = \frac{1}{f'(\ln y)} = \frac{1}{e^{\ln y}} = \frac{1}{y}$

Derivatives of Logarithmic Functions

- ▶ The logarithmic function $y = \log_b x$ is the inverse of the one-to-one exponential function $b^y = b^{\log x} = x$ which is differentiable. So the logarithmic function is also differentiable.

Derivatives of Logarithmic Functions

- ▶ The logarithmic function $y = \log_b x$ is the inverse of the one-to-one exponential function $b^y = b^{\log_b x} = x$ which is differentiable. So the logarithmic function is also differentiable.
- ▶ Applying implicit differentiation to $b^y = x$

$$\frac{d}{dx}(b^y) = \frac{d}{dx}(x) \iff \frac{d}{dy}(b^y) \frac{dy}{dx} = 1 \iff b^y \ln b \frac{dy}{dx} = 1 \iff$$

$$\frac{dy}{dx} = \frac{1}{b^y \ln b} = \frac{1}{x \ln b} \iff \frac{d}{dx}(\log_b x) = \frac{1}{x \ln b}$$

Derivatives of Logarithmic Functions

- ▶ The logarithmic function $y = \log_b x$ is the inverse of the one-to-one exponential function $b^y = b^{\log_b x} = x$ which is differentiable. So the logarithmic function is also differentiable.
- ▶ Applying implicit differentiation to $b^y = x$

$$\frac{d}{dx}(b^y) = \frac{d}{dx}(x) \iff \frac{d}{dy}(b^y) \frac{dy}{dx} = 1 \iff b^y \ln b \frac{dy}{dx} = 1 \iff$$

$$\frac{dy}{dx} = \frac{1}{b^y \ln b} = \frac{1}{x \ln b} \iff \frac{d}{dx}(\log_b x) = \frac{1}{x \ln b}$$

- ▶ If $b = e$, we get

$$\frac{dy}{dx}(\ln x) = \frac{1}{x}$$

Derivatives of Logarithmic Functions - Example I

- Determine $f'(x)$ for $f(x) = \ln |x|$

$$f(x) = \begin{cases} \ln x & \text{if } x > 0 \\ \ln(-x) & \text{if } x < 0 \end{cases}$$

Derivatives of Logarithmic Functions - Example I

- Determine $f'(x)$ for $f(x) = \ln |x|$

$$f(x) = \begin{cases} \ln x & \text{if } x > 0 \\ \ln(-x) & \text{if } x < 0 \end{cases}$$

- Then

$$f'(x) = \begin{cases} \frac{1}{x} & \text{if } x > 0 \\ \frac{1}{-x}(-1) = \frac{1}{x} & \text{if } x < 0 \end{cases}$$

Derivatives of Logarithmic Functions - Example I

- Determine $f'(x)$ for $f(x) = \ln |x|$

$$f(x) = \begin{cases} \ln x & \text{if } x > 0 \\ \ln(-x) & \text{if } x < 0 \end{cases}$$

- Then

$$f'(x) = \begin{cases} \frac{1}{x} & \text{if } x > 0 \\ \frac{1}{-x}(-1) = \frac{1}{x} & \text{if } x < 0 \end{cases}$$

- It follows that $f'(x) = \frac{1}{x}$ for all $x \neq 0$.

Derivatives of Logarithmic Functions - Example II

- ▶ Determine $f'(x)$ for $f(x) = \ln(x^3 + 1)$.

Derivatives of Logarithmic Functions - Example II

- ▶ Determine $f'(x)$ for $f(x) = \ln(x^3 + 1)$.
- ▶ To use the chain rule, let $u = x^3 + 1$ and $y = \ln u$. Then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{u} \frac{du}{dx} = \frac{1}{x^3 + 1} (3x^2) = \frac{3x^2}{x^3 + 1}$$

Derivatives of Logarithmic Functions - Example II

- ▶ Determine $f'(x)$ for $f(x) = \ln(x^3 + 1)$.
- ▶ To use the chain rule, let $u = x^3 + 1$ and $y = \ln u$. Then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{u} \frac{du}{dx} = \frac{1}{x^3 + 1} (3x^2) = \frac{3x^2}{x^3 + 1}$$

- ▶ More generally,

$$\frac{d}{dx}(\ln u) = \frac{1}{u} \frac{du}{dx}$$

Derivatives of Logarithmic Functions - Example II

- ▶ Determine $f'(x)$ for $f(x) = \ln(x^3 + 1)$.
- ▶ To use the chain rule, let $u = x^3 + 1$ and $y = \ln u$. Then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{u} \frac{du}{dx} = \frac{1}{x^3 + 1} (3x^2) = \frac{3x^2}{x^3 + 1}$$

- ▶ More generally,

$$\frac{d}{dx}(\ln u) = \frac{1}{u} \frac{du}{dx}$$

- ▶ Determine $f'(x) = \ln(\sin x)$. Let $u = \sin x$ and $y = \ln u$. Then

$$\frac{d}{dx} \ln(\sin x) = \frac{1}{\sin x} \frac{d}{dx} \sin x = \frac{1}{\sin x} \cos x = \cot x$$

Logarithmic Differentiation

- Determine $f'(x)$ for $f(x) = \sqrt{\frac{x-1}{x^4+1}}, x > 1$

Logarithmic Differentiation

- ▶ Determine $f'(x)$ for $f(x) = \sqrt{\frac{x-1}{x^4+1}}, x > 1$
- ▶ Let $y = \sqrt{\frac{x-1}{x^4+1}}$

Logarithmic Differentiation

- ▶ Determine $f'(x)$ for $f(x) = \sqrt{\frac{x-1}{x^4+1}}, x > 1$
- ▶ Let $y = \sqrt{\frac{x-1}{x^4+1}}$
- ▶ Then

$$\ln y = \ln \sqrt{\frac{x-1}{x^4+1}} = \frac{1}{2} \ln \frac{x-1}{x^4+1} = \frac{1}{2} \ln(x-1) - \frac{1}{2} \ln(x^4+1)$$

Logarithmic Differentiation

- ▶ Determine $f'(x)$ for $f(x) = \sqrt{\frac{x-1}{x^4+1}}$, $x > 1$
- ▶ Let $y = \sqrt{\frac{x-1}{x^4+1}}$
- ▶ Then

$$\ln y = \ln \sqrt{\frac{x-1}{x^4+1}} = \frac{1}{2} \ln \frac{x-1}{x^4+1} = \frac{1}{2} \ln(x-1) - \frac{1}{2} \ln(x^4+1)$$

- ▶ Implicit differentiation and chain rule give

$$y' \frac{1}{y} = \frac{1}{2} \frac{1}{x-1} - \frac{1}{2} \frac{1}{x^4+1} 4x^3 = \frac{1}{2x-2} - \frac{2x^3}{x^4+1}$$

Logarithmic Differentiation

- ▶ Determine $f'(x)$ for $f(x) = \sqrt{\frac{x-1}{x^4+1}}$, $x > 1$
- ▶ Let $y = \sqrt{\frac{x-1}{x^4+1}}$
- ▶ Then

$$\ln y = \ln \sqrt{\frac{x-1}{x^4+1}} = \frac{1}{2} \ln \frac{x-1}{x^4+1} = \frac{1}{2} \ln(x-1) - \frac{1}{2} \ln(x^4+1)$$

- ▶ Implicit differentiation and chain rule give

$$y' \frac{1}{y} = \frac{1}{2} \frac{1}{x-1} - \frac{1}{2} \frac{1}{x^4+1} 4x^3 = \frac{1}{2x-2} - \frac{2x^3}{x^4+1}$$

- ▶ Solving for y' and back substituting

$$y' = \sqrt{\frac{x-1}{x^4+1}} \left(\frac{1}{2x-2} - \frac{2x^3}{x^4+1} \right)$$

Proving General Power Rule

- ▶ Determine $f'(x)$ for $f(x) = x^n$ for any $n \in \mathbb{R}$.

Proving General Power Rule

- ▶ Determine $f'(x)$ for $f(x) = x^n$ for any $n \in \mathbb{R}$.
- ▶ Since $f(x)$ can be negative or 0, logarithmic differentiation cannot be used directly.

Proving General Power Rule

- ▶ Determine $f'(x)$ for $f(x) = x^n$ for any $n \in \mathbb{R}$.
- ▶ Since $f(x)$ can be negative or 0, logarithmic differentiation cannot be used directly.
- ▶ $f'(0) = 0$ can be shown directly from the definition of the derivative.

Proving General Power Rule

- ▶ Determine $f'(x)$ for $f(x) = x^n$ for any $n \in \mathbb{R}$.
- ▶ Since $f(x)$ can be negative or 0, logarithmic differentiation cannot be used directly.
- ▶ $f'(0) = 0$ can be shown directly from the definition of the derivative.
- ▶ Let $y = x^n$ for $x \neq 0$. Then

$$\ln |y| = \ln |x^n| = n \ln |x|$$

Proving General Power Rule

- ▶ Determine $f'(x)$ for $f(x) = x^n$ for any $n \in \mathbb{R}$.
- ▶ Since $f(x)$ can be negative or 0, logarithmic differentiation cannot be used directly.
- ▶ $f'(0) = 0$ can be shown directly from the definition of the derivative.
- ▶ Let $y = x^n$ for $x \neq 0$. Then

$$\ln |y| = \ln |x^n| = n \ln |x|$$

- ▶ Implicit differentiation gives

$$\frac{1}{y}y' = n\frac{1}{x} \iff y' = yn\frac{1}{x} = x^n n\frac{1}{x} = nx^{n-1}$$

Linear Approximation

- ▶ Zooming in toward a point on the graph of a differentiable function, it looks more and more like its tangent line.

Linear Approximation


- ▶ Zooming in toward a point on the graph of a differentiable function, it looks more and more like its tangent line.
- ▶ Tangent lines are therefore good approximation of function values near the tangent point.

Linear Approximation

- ▶ Zooming in toward a point on the graph of a differentiable function, it looks more and more like its tangent line.
- ▶ Tangent lines are therefore good approximation of function values near the tangent point.
- ▶ Equation of the tangent line of a differentiable function at a is
$$y = f(a) + f'(a)(x - a)$$

Linear Approximation

- ▶ Zooming in toward a point on the graph of a differentiable function, it looks more and more like its tangent line.
- ▶ Tangent lines are therefore good approximation of function values near the tangent point.
- ▶ Equation of the tangent line of a differentiable function at a is $y = f(a) + f'(a)(x - a)$
- ▶ The linear function given by $L(x) = f(a) + f'(a)(x - a)$ is called the linearization of f at a .



Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$
- ▶ $f(1) = 2$ and $f'(1) = \frac{1}{4}$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$
- ▶ $f(1) = 2$ and $f'(1) = \frac{1}{4}$
- ▶ The linearization $L(x)$ of f at $a = 1$ becomes

$$L(x) = f(1) + f'(1)(x - 1) = 2 + \frac{1}{4}(x - 1) = \frac{7}{4} + \frac{x}{4}$$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$
- ▶ $f(1) = 2$ and $f'(1) = \frac{1}{4}$
- ▶ The linearization $L(x)$ of f at $a = 1$ becomes

$$L(x) = f(1) + f'(1)(x-1) = 2 + \frac{1}{4}(x-1) = \frac{7}{4} + \frac{x}{4}$$

- ▶ Suppose that we want to approximate $f(x)$ around x so that the error is < 0.5 . Which values of x will satisfy that?

$$\left| \sqrt{x+3} - \left(\frac{7}{4} + \frac{x}{4} \right) \right| < 0.5 \iff \sqrt{x+3} - 0.5 < \frac{7}{4} + \frac{x}{4} < \sqrt{x+3} + 0.5$$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$
- ▶ $f(1) = 2$ and $f'(1) = \frac{1}{4}$
- ▶ The linearization $L(x)$ of f at $a = 1$ becomes

$$L(x) = f(1) + f'(1)(x-1) = 2 + \frac{1}{4}(x-1) = \frac{7}{4} + \frac{x}{4}$$

- ▶ Suppose that we want to approximate $f(x)$ around x so that the error is < 0.5 . Which values of x will satisfy that?

$$\left| \sqrt{x+3} - \left(\frac{7}{4} + \frac{x}{4} \right) \right| < 0.5 \iff \sqrt{x+3} - 0.5 < \frac{7}{4} + \frac{x}{4} < \sqrt{x+3} + 0.5$$

- ▶ These two inequalities imply that the error is < 0.5 if $x \in]-2.66, 8.66[$

Linear Approximation - Example

- ▶ Let $f(x) = \sqrt{x+3} = (x+3)^{\frac{1}{2}}$
- ▶ $f'(x) = \frac{1}{2}(x+3)^{-\frac{1}{2}} = \frac{1}{2\sqrt{x+3}}$
- ▶ $f(1) = 2$ and $f'(1) = \frac{1}{4}$
- ▶ The linearization $L(x)$ of f at $a = 1$ becomes

$$L(x) = f(1) + f'(1)(x-1) = 2 + \frac{1}{4}(x-1) = \frac{7}{4} + \frac{x}{4}$$

- ▶ Suppose that we want to approximate $f(x)$ around x so that the error is < 0.5 . Which values of x will satisfy that?

$$\left| \sqrt{x+3} - \left(\frac{7}{4} + \frac{x}{4} \right) \right| < 0.5 \iff \sqrt{x+3} - 0.5 < \frac{7}{4} + \frac{x}{4} < \sqrt{x+3} + 0.5$$

- ▶ These two inequalities imply that the error is < 0.5 if $x \in]-2.66, 8.66[$
- ▶ If the permitted error is < 0.1 , then $x \in]-1.1, 3.9[$

Differentials

- ▶ Differentials provide another way of viewing linear approximations
- ▶ Let f be a differentiable function and let $y = f(x)$

Differentials


- ▶ Differentials provide another way of viewing linear approximations
- ▶ Let f be a differentiable function and let $y = f(x)$
- ▶ Let dx denote an independent variable that can be given the value of any real number.

Differentials

- ▶ Differentials provide another way of viewing linear approximations
- ▶ Let f be a differentiable function and let $y = f(x)$
- ▶ Let dx denote an independent variable that can be given the value of any real number.
- ▶ The differential dy is then the dependent variable given by $dy = f'(x)dx$.

Differentials

- ▶ Differentials provide another way of viewing linear approximations
- ▶ Let f be a differentiable function and let $y = f(x)$
- ▶ Let dx denote an independent variable that can be given the value of any real number.
- ▶ The differential dy is then the dependent variable given by $dy = f'(x)dx$.
- ▶ Let $P = (x, f(x))$ and $Q = (x + \Delta x, f(x + \Delta x))$ and let $dx = \Delta x$. Δy represents how much the curve falls or rises when x changes by $\Delta x = dx$ while dy how much the tangent line falls or rises by the same change of x .



Summary

You should after this lecture be familiar with:

- ▶ Chain rule
- ▶ Implicit differentiation
- ▶ Logarithmic differentiation
- ▶ Linear approximation