

Compulsory Exercise 2: Group 37

TMA4268 Statistical Learning V2019

Anders Bendiksen and Helge Bergo

31 March, 2020

Problem 1 (10p)

a) Ridge Regression (2p)

Show that the ridge regression estimator is $\hat{\beta}_{Ridge} = (X^T X + \lambda I)^{-1} X^T y$.

b) (2p)

Find the expected value and the variance-covariance matrix of $\hat{\beta}_{Ridge}$ (1P each).

c) (2P) - Multiple choice

- (i) TRUE
- (ii) FALSE
- (iii) FALSE
- (iv) TRUE

d) Forward Selection

```
library(ISLR)
set.seed(1)
train.ind = sample(1:nrow(College), 0.5*nrow(College))
college.train = College[train.ind,]
college.test = College[-train.ind,]
```

After dividing the data into a training and test set, the `regsubsets` function was used to create a forward selection model on the data, from the `leaps`-library.

```
library(leaps)
regfit.fwd = regsubsets(Outstate~., data=college.train, method="forward", nvmax = 18)
reg.summary = summary(regfit.fwd)
```

To decide on which model is best, the number of variables used in the selection was plotted against RSS, Cp, BIC and adjusted R^2 .

```
par(mfrow=c(2,2))
plot(reg.summary$rss, xlab="Number of variables", ylab="RSS", type="b")
```


```

plot(reg.summary$adjr2,xlab="Number of variables",ylab="Adjusted Rsq",type="b")
max.adj2 = which.max(reg.summary$adjr2)
points(max.adj2,reg.summary$adjr2[max.adj2], col="black",cex=2,pch=20)

plot(reg.summary$cp,xlab="Number of variables",ylab="Cp",type="b")
min.cp = which.min(reg.summary$cp)
points(min.cp,reg.summary$cp[min.cp], col="black",cex=2,pch=20)

plot(reg.summary$bic,xlab="Number of variables",ylab="BIC",type="b")
min.bic = which.min(reg.summary$bic)
points(min.bic,reg.summary$bic[min.bic], col="black",cex=2,pch=20)

```



The maximum adjusted R^2 is the one with 14 variables, with a value of 0.7706887, shown as a filled dot in the upper right plot. This is also the same number of variables as for the lowest Cp. However, all the plots are pretty flat after around 6 or 7 variables used, and it seems like using only 6 variables still gives a good adjusted R^2 value of 0.7516133, without the increased complexity of adding 7 more variables. The model is then:

```

coef(regfit.fwd,6)

```

##	(Intercept)	PrivateYes	Room.Board	Terminal	perc.alumni
##	-4726.8810613	2717.7019276	1.1032433	36.9990286	59.0863753
##	Expend	Grad.Rate			
##	0.1930814	33.8303314			

For the MSE, the following code calculates the MSE for all the variables.

```

val.errors = rep(NA,17)
x.test = model.matrix(Outstate~.,data=college.test) # notice the -index!

```

```

for (i in 1:17) {
  coefi = coef(regfit.fwd,id=i)
  pred = x.test[,names(coefi)]%*%coefi
  val.errors[i] = mean((college.test$Outstate-pred)^2)
}

# plot(sqrt(val.errors),xlab="Number of variables", ylab="Root MSE",ylim=c(1500,5000) ,pch=19,type="b")
# points(sqrt(regfit.fwd$rss[-1]/180),col="blue",pch=19,type="b")
# legend("topright",legend=c("Training","Validation"),col=c("black","blue"),pch=19)

```

The MSE of the model with 6 variables is then:

```
val.errors[6]
```

```
## [1] 3844857
```

e) (2p)


Using the Lasso method from the `glmnet`-library, a new model was selected.

To select the tuning parameter λ , cross-validation was performed, and the λ giving the lowest MSE was selected.

```

cv.out = cv.glmnet(x.train,y.train, alpha = 1)
plot(cv.out)

```



```

best.lambda = cv.out$lambda.min
best.lambda

```

```
## [1] 10.7207
```

This was used on the test set, to get the MSE for the

```

lasso.pred = predict(lasso.model,s=best.lambda ,newx=x.test)
MSE = mean((lasso.pred-y.test)^2)
MSE

```

```
## [1] 3688061
```

Finally, the coefficients of the model are shown here:

```
lasso.coef = predict(cv.out,type="coefficients",s=best.lambda)[1:18,]  
lasso.coef
```

```
##      (Intercept)      PrivateYes          Apps          Accept          Enroll  
## -1.172140e+03  2.230467e+03 -2.825215e-01  6.615811e-01 -3.778631e-01  
##      Top10perc      Top25perc      F.Undergrad      P.Undergrad      Room.Board  
##  4.589180e+01 -1.485674e+01 -5.800132e-02 -5.713770e-02  1.088115e+00  
##           Books      Personal          PhD          Terminal          S.F.Ratio  
## -9.185125e-01 -3.005419e-01  4.013410e+00  2.996744e+01 -6.936391e+01  
##      perc.alumni          Expend          Grad.Rate  
##  4.686967e+01  1.480013e-01  2.431539e+01
```

Problem 2 (9p)

a) (2p) - Multiple choice


Which of the following statements are true, which false?

- (i) A regression spline of order 3 with 4 knots has 8 basis functions.
- (ii) A regression spline with polynomials of degree $M - 1$ has continuous derivatives up to order $M - 2$, but not at the knots.
- (iii) A natural cubic spline is linear beyond the boundary knots.
- (iv) A smoothing spline is (a shrunken version of) a natural cubic spline with knots at the values of all data points x_i for $i = 1, \dots, n$.
- (v)
- (vi)
- (vii)
- (viii)

b) (2p)

Write down the basis functions for a cubic spline with knots at the quartiles q_1, q_2, q_3 of variable X .

c) (2p)




From these plots, it seems like `Room.board`, `perc.alumni` and `Grad.Rate` all have quite linear relationships with `Outstate`, while both `Terminal` and `Expend` seem to follow a non-linear relationship.

d) (3P)

- (i) Fit polynomial regression models for `Outstate` with `Terminal` as the only covariate for a range of polynomial degrees ($d = 1, \dots, 10$) and plot the results. Use the training data (`college.train`) for this task.


```
cols = rainbow(10)
deg = 1:10
polyfunc = function(d) {
  model = lm(Outstate ~ poly(Terminal,d), data=college.train)
  lines(cbind(college.train$Terminal, model$fit)[order(college.train$Terminal),], col=cols[d])
  pred = predict(model, college.train)
  mean((pred - college.train$Outstate)^2)
}
plot(college.train$Terminal, college.train$Outstate, col = "gray", pch=19,
     cex = 0.5, xlab = "Terminal", ylab = "Outstate")
MSE = sapply(deg, polyfunc)
legend("topleft", legend = paste("degree = ", deg), col = cols, cex = 0.4)
```



- (ii) Still for the training data, choose a suitable smoothing spline model to predict `Outstate` as a function of `Expend` (again as the only covariate) and plot the fitted function into the scatterplot of `Outstate` against `Expend`. How did you choose the degrees of freedom?

```
library(splines)
attach(college.train)
expend.range = range(Expend)
expend.grid = seq(from=expend.range[1],to=expend.range[2])

plot(Expend, Outstate, col = "darkgrey", pch=19, cex = 0.5)
fit.smoothspline = smooth.spline(Expend,Outstate,cv=TRUE)
lines(fit.smoothspline)
```



```
# legend("bottomright",legend=paste("DF =",round(fit.smoothspline$df,2)),cex=.8)
```

The degrees of freedom was chosen using cross-validation, and the result was 4.661.

- (iii) Report the corresponding training MSE for (i) and (ii). Did you expect that?

```
MSE.smoothspline.train = mean((predict(fit.smoothspline, college.train$Expend)$y -
                                college.train$Outstate)^2)
MSE.smoothspline.train
```

```
## [1] 6871281
```

```
# MSE.smoothspline.test = mean((predict(fit.smoothspline, college.test$Expend)$y -
                                         # college.test$Outstate)^2)
# MSE.smoothspline.test
MSE
```

```
## [1] 15075161 14330586 14249448 14247330 14231485 14230392 14153207
## [8] 14097911 13841526 13822205
```

The MSE for the polynomial regression is much higher than the MSE for the smoothing splines, but this makes a lot of sense when looking at the initial plots from 2.c). For the **Expend** variable, it seems like the data have a clearer trend than for the **Terminal** variable, and therefore the MSE is much lower.

Problem 3 (9p)

a) (2P) - Multiple choice

Which of the following statements are true, which false?

- (i) Regression trees cannot handle categorical predictors.
- (ii) Regression and classification trees are easy to interpret.
- (iii) The random forest approaches improves bagging, because it reduces the variance of the predictor function by decorrelating the trees.
- (iv) The number of trees B in bagging and random forests is a tuning parameter.

b) (4P)

Select one method from Module 8 (tree-based methods) in order to build a good model to predict **Outstate** in the **College** dataset that we used in problems 1 and 2. Explain your choice (pros/cons?) and how you chose the tuning parameter(s). Train the model using the training data and report the MSE for the test data.

c) (2p)

Compare the results (tests MSEs) among all the methods you used in Problems 1-3. Which method perform best in terms of prediction error? Which method would you choose if the aim is to develop an interpretable model?

Problem 4 (12P)

We will use the classical data set of *diabetes* from a population of women of Pima Indian heritage in the US, available in the R **MASS** package. The following information is available for each woman:

- diabetes: 0= not present, 1= present
- npreg: number of pregnancies
- glu: plasma glucose concentration in an oral glucose tolerance test
- bp: diastolic blood pressure (mmHg)

- skin: triceps skin fold thickness (mm)
- bmi: body mass index (weight in kg/(height in m)²)
- ped: diabetes pedigree function.
- age: age in years

We will use a training set (called `d.train`) with 300 observations (200 non-diabetes and 100 diabetes cases) and a test set (called `d.test`) with 232 observations (155 non-diabetes and 77 diabetes cases). Our aim is to make a classification rule for the presence of diabetes (yes/no) based on the available data. You can load the data as follows:

a) (2P) - Multiple choice

```
summary(d.train)
```


```
##      diabetes      npreg      glu      bp
## Min.   :0.0000   Min.    : 0.000   Min.    : 56.00   Min.    : 30.00
## 1st Qu.:0.0000   1st Qu.: 1.000   1st Qu.: 96.75   1st Qu.: 64.00
## Median :0.0000   Median : 2.000   Median :114.00   Median : 71.00
## Mean   :0.3333   Mean    : 3.467   Mean    :120.13   Mean    : 71.56
## 3rd Qu.:1.0000   3rd Qu.: 5.250   3rd Qu.:140.25   3rd Qu.: 80.00
## Max.   :1.0000   Max.    :17.000   Max.    :199.00   Max.    :110.00
##      skin      bmi      ped      age
## Min.   : 7.00   Min.   :18.20   Min.   :0.0850   Min.   :21.00
## 1st Qu.:22.00   1st Qu.:27.98   1st Qu.:0.2567   1st Qu.:23.00
## Median :29.00   Median :32.80   Median :0.4150   Median :27.00
## Mean   :29.14   Mean    :33.03   Mean    :0.5004   Mean    :31.55
## 3rd Qu.:36.00   3rd Qu.:37.12   3rd Qu.:0.6210   3rd Qu.:37.25
## Max.   :99.00   Max.    :67.10   Max.    :2.4200   Max.    :81.00
```

```
str(d.train)
```


```
## 'data.frame':   300 obs. of  8 variables:
## $ diabetes: num  1 1 1 1 1 1 1 1 1 1 ...
## $ npreg   : int  7 7 9 9 8 2 12 2 1 0 ...
## $ glu     : int 109 152 119 112 151 90 92 197 181 124 ...
## $ bp      : int  80 88 80 82 78 68 62 70 78 70 ...
## $ skin    : int  31 44 35 32 32 42 7 45 42 20 ...
## $ bmi     : num 35.9 50 29 34.2 42.9 38.2 27.6 30.5 40 27.4 ...
## $ ped     : num 1.127 0.337 0.263 0.26 0.516 ...
## $ age     : int  43 36 29 36 36 27 44 53 22 36 ...
```

```
plot(glu,diabetes,col="red")
```


```
points(bmi,diabetes,col="blue")
```

```
plot(npreg)
```



```
plot(log(npreg))
```



```
max(npreg)
```

```
## [1] 17
```

Start by getting to know the *training data*, by producing summaries and plots. Which of the following statements are true, which false?

- (i) Females with high glucose levels and higher bmi seem to have a higher risk for diabetes.
- (ii) Some women had up to 17 pregnancies.
- (iii) BMI and triceps skin fold thickness seem to be positively correlated.
- (iv) The distribution of the number of pregnancies per woman seems to be a bit skewed and a transformation of this variable could therefore be appropriate.
- (v)
- (vi) TRUE
- (vii) TRUE
- (viii)

b) (4P)

Fit a support vector classifier (linear boundary) and a support vector machine (radial boundary) to find good functions that predict the diabetes status of a patient. Use cross-validation to find a good cost parameter (for the linear boundary) and a good combination of cost *and* γ parameters (for the radial boundary). Report the confusion tables and misclassification error rates for the test set in both cases. Which classifier do you prefer and why? (Do not use any variable transformations or standardizations to facilitate correction).

R-hints: The response variable must be converted into a factor variable before you continue.

```
d.train$diabetes <- as.factor(d.train$diabetes)
d.test$diabetes <- as.factor(d.test$diabetes)
library(e1071)
```

To run cross-validation over a grid of two tuning parameters, you can use the `tune()` function where `ranges` defines the grid points as follows:

```
tune(..., formula, kernel=..., ranges=list(cost=c(...), gamma=c(...)))
```

c) (2P)

Compare the performance of the two classifiers from b) to *one other classification method* that you have learned about in the course. Explain your choice and report the confusion table and misclassification error rate on the test set for your chosen method and interpret what you see. What are advantages/disadvantages of your chosen method with respect to SVMs?

d) (2P) - Multiple choice

Which of the following statements are true, which false?

- (i) Under standard conditions, the maximal margin hyperplane approach is equivalent to a linear discriminant analysis.
- (ii) Under standard conditions, the support vector classifier is equivalent to quadratic discriminant analysis.
- (iii) Logistic regression, LDA and support vector machines tend to perform similar when decision boundaries are linear, unless classes are linearly separable.
- (iv) An advantage of logistic regression over SVMs is that it is easier to do feature selection and to interpret the results.

e) (2P) Link to logistic regression and hinge loss.

Look at slides 71-73 of Module 9. Show that the loss function

$$\log(1 + \exp(-y_i f(\mathbf{x}_i)))$$

is the deviance for the $y = -1, 1$ encoding in a logistic regression model.

Hint: $f(\mathbf{x}_i)$ corresponds to the linear predictor in the logistic regression approach.

Problem 5 (10P)

The following dataset consists of 40 tissue samples with measurements of 1,000 genes. The first 20 tissues come from healthy patients and the remaining 20 come from a diseased patient group. The following code loads the dataset into your session with row names describing if the tissue comes from a diseased or healthy person.

```
# id <- "1VfVCQvWt121UN39NXZ4aR9Dmsbj-p90U" # google file ID
# GeneData <- read.csv(sprintf("https://docs.google.com/uc?id=%s&export=download", id), header=F)
# colnames(GeneData)[1:20] = paste(rep("H", 20), c(1:20), sep = "")
# colnames(GeneData)[21:40] = paste(rep("D", 20), c(1:20), sep = "")
# row.names(GeneData) = paste(rep("G", 1000), c(1:1000), sep = "")
```

a) (2P)

Perform hierarchical clustering with complete, single and average linkage using **both** Euclidean distance and correlation-based distance on the dataset. Plot the dendrograms. Hint: You can use `par(mfrow=c(1,3))` to plot all three dendrograms on one line or `par(mfrow=c(2,3))` to plot all six together.

b) (2P)

Use these dendrograms to cluster the tissues into two groups. Compare the groups with respect to the patient group the tissue comes from. Which linkage and distance measure performs best when we know the true state of the tissue?

c) (1P)

With Principal Component Analysis, the first principal component loading vector solves the following optimization problem,

$$\max_{\phi_{11}, \dots, \phi_{p1}} \left\{ \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \phi_{j1} x_{ij} \right)^2 \right\} \quad \text{subject to} \quad \sum_{j=1}^p \phi_{j1}^2 = 1.$$

Explain what ϕ , p , n and x are in this optimization problem and write down the formula for the first principal component scores.

d) (2P)

- (i) (1P) Use PCA to plot the samples in two dimensions. Color the samples based on the tissues group of patients.
- (ii) (1P) How much variance is explained by the first 5 PCs?

e) (1P)

Use your results from PCA to find which genes that vary the most across the two groups.

f) (2P)

Use K-means to separate the tissue samples into two groups. Plot the values in a two-dimensional space with PCA. What is the error rate of K-means?