

ACUTE EFFECT OF DIFFERENT HALF SQUAT SET CONFIGURATIONS ON JUMP POTENTIATION

Running Title

Greater potentiation after cluster set

Authors

Daniel A. Boullosa¹; Laurinda Abreu²; Luis G.N. Beltrame¹, and David G. Behm³

Affiliations

¹ Post-Graduate Program in Physical Education, Catholic University of Brasilia, Brazil.

² Lavadores, Vigo, Spain.

³ School of Human Kinetics and Recreation; Memorial University of Newfoundland, St. John's Newfoundland, Canada.

Number of tables: 4

Number of figures: 0

Word count: 3604

Abstract word count: 287

Address for correspondence:

Prof. Dr. Daniel A. Boullosa
Pós-Graduação Stricto Sensu em Educação Física
Universidade Católica de Brasília - UCB
QS 07, LT1 S/N - Sala 111 - Bloco G
71966-700 Águas Claras - DF - Brasil
Phone number/FAX: +55 61 3356-9489
E-mail: d_boullosa@yahoo.es

ABSTRACT

The aim of this study was to evaluate the acute effect of a half squat exercise performed with different set configurations on jump potentiation. Twelve resistance-trained men were evaluated on three occasions separated by 48-96 hrs. First, they performed a 5 repetitions maximum test (5RM). Subsequently, they performed in a randomized order two sessions: one session with 5RM until failure, and another session with the same workload but with 30 s rest intervals between repetitions (i.e. cluster set; CS). Countermovement jump performance was examined during the second and third sessions for jump height and force-time parameters using a force platform at the following time intervals: before and at 1, 3, 6, 9, and 12 min. Separate comparisons for each variable at the different time intervals were analyzed using ANOVA, effect size, and qualitative inferences. The majority of the parameters improved independently of the time they occur, except peak force and vertical stiffness after a set until failure. For peak power, it appears the cluster treatment resulted in superior potentiation at 1 min whereas the 5RM treatment resulted in greater potentiation at 9 min. Effect size analysis and qualitative outcomes revealed an improvement in vertical stiffness and a lowering in the depth of the countermovement in CS. There were significant correlations between participants' 5RM relative performance and various force-time parameters only in CS. It appears that a CS induces greater peak power than a 5RM set at 1 min, although the reverse occurs at 9 min. Delayed potentiation associated with the 5RM may be attributed to greater fatigue versus the CS approach. Therefore, it follows that the optimal method for inducing peak power potentiation is dependent on the available time between heavy half-squat exercise and subsequent jump performance.

Keywords: post-activation potentiation, fatigue, complex training, warm-up, resistance training

INTRODUCTION

Post-activation potentiation (PAP) refers to the phenomena by which muscular performance characteristics are acutely enhanced as a result of their contractile history (25). Although PAP can provide physiological enhancement, it is not always related to functional performance improvements (13). For instance, increases in peak power (physiological parameter) during the push-off of a jump after squat exercises have been previously observed without any increment in jump height (functional performance) (22, 16, 4). These and other previous reports suggest that potentiation could be evident depending on the method of evaluation and/or the parameters selected for analysis, with the fatigue-potentiation relationship modulating the acute effect of the conditioning exercise during recovery (24, 1, 3). Interestingly, in the recent paper of Chaouachi et al. (4), the potentiation of selected parameters during jumping after various workloads of half squat exercises were quite variable between individuals with a significant deterioration of the jump capacity observed (4). This fact could explain the conflicting results in previous literature (25). Therefore, since the individual response to conditioning stimuli is important for the fatigue-potentiation interaction and its influence on subsequent performance, it is necessary to explore new strategies for designing conditioning protocols that concurrently enhance potentiation and limit the fatiguing response for optimal performance. This could help also for clarifying the inconsistency in PAP responses previously reported that could be related to the masking effect of fatigue on potentiation (1).

Set configuration is one of the variables that could be modified when designing resistance (RE) and jump exercises (9, 10). Recently, various studies have suggested that introducing rest intervals between exercise repetitions instead of between sets (i.e. cluster training), could be an

effective method for improving mechanical performance (7) with a lower loss of velocity and power over the whole training session (17, 11), allowing also a greater training volume (7, 15). Regarding RE, this better performance could be explained by the fact that with cluster training it is the reduced time under tension, which has been suggested to be a key factor for muscular fatigue development in RE (26). Moreover, Hardee et al. (12) have recently reported a lower rating of perceived exertion (RPE) with the introduction of rest intervals between repetitions during clean pulls when compared to the same load with traditional set configuration. Thus, it may be suggested that cluster training is an effective strategy for improving mechanical performance with a lower development of fatigue in RE. This could be an interesting approach for conditioning stimuli when looking for potentiation responses during complex training sessions or during competitions. However, to the best of our knowledge, there are no studies comparing the acute responses of cluster training with traditional set configuration. This information would be valuable for designing more effective and efficient conditioning activities in those sports requiring power performances such as jumping, sprinting or even team sports.

Therefore, the aim of this study was to compare the acute effect on jump capacity of different set configuration with the same workload (i.e. cluster set [CS] vs. set until failure) of half squats. These exercises were selected as most PAP studies have utilized them and they are closed kinetic chain activities (greater sport specificity). It was hypothesized that a CS could favor a greater magnitude and a more rapid potentiation response while jumping because of the lower fatigue development when compared to a set until failure.

METHODS

Experimental Approach to the problem

After determination of the load corresponding to 5 repetition maximum (5RM) in half squat, participants were evaluated in random order on two separate occasions for comparing the acute effect (i.e. at 1, 3, 6, 9, and 12 min of recovery) of this workload with different set configurations (i.e. set until failure vs. cluster set) on jump performance and the associated force-time parameters. The independent variable was set configuration, while jump height and kinetic parameters over the whole recovery were the dependent variables.

Participants

Twelve aspiring fire-fighters volunteered for participation on this study. Their characteristics are presented on Table 1. All of them had a minimum of 6 months of experience in RT and countermovement jumps (CMJ) evaluations. Their training routines included resistance training, power exercises, sprinting, aerobic running, swimming, and agility as other specific fire-fighting skills. Their performance level was very heterogeneous with mean best performances of ~100 kg in bench press RM; ~20 pull-ups; ~12 s in 100 m; ~3 min in 1000 m; and ~1 min 20 s in 100 m freestyle. After receiving a detailed explanation of the procedures of the study, participants provided informed written consent. The study received the approval of the local Ethics Committee for experimentation with human participants.

Place Table 1 about here

Experimental design and Procedures

This experiment was performed in February in Spain. During these days, participants executed low-to-medium intense activities for fitness maintenance. Half squat was selected because it is the exercise most utilized with leg power conditioning and for obtaining potentiation responses (18). The load of 5 RM was also selected as it usually represents > 80% RM which may be a sufficient intensity for inducing potentiation while jumping (25). Participants were fully familiarized with the procedures.

After a brief warm up of 10 min of easy running on a treadmill, 5 min of calisthenics, and 20 repetitions of abdominal crunches, participants were instructed to perform a half squat 5 RM test. All the procedures were supervised by the primary investigator who was also the physical coach for the participants at that time. The half squat was performed with free weights until 90° of knee flexion was achieved. The progression of the test was as follows, with 3-5 min of rest between sets: 10 repetitions of an estimated 50% RM; 5 repetitions of an estimated 70% RM; and subsequent attempts of 3 repetitions of estimated 5RM until the maximum load was determined and fully executed without technical difficulties. All participants achieved their 5RM with no more than 6 sets over the whole test. We decided not to include additional evaluations for determining testing reproducibility as they were habitually training with this exercise load. Therefore, a high reproducibility and reliability could be expected. Additionally, more testing sessions could have an influence on performance changes thus biasing results. The highest load was selected for the next two experimental sessions.

After determination of the best 5RM, participants performed in a randomized order the experimental protocols with a minimum of 48 hrs and a maximum of 96 hrs between sessions. All the sessions were performed during the morning at the same time of the day for every participant. They were asked not to drink caffeine beverages 1 hr before evaluations. They were allowed to drink water “ad libitum” during the whole experiment. Both experimental sessions included the previously described warm up plus a set of 5 repetitions at 50% of 5RM, and 2 maximal CMJ separated by 15 s. The best CMJ was selected for further analysis. After 3 min of rest, they performed a set of 5RM (i.e. traditional set configuration), or the same load but with 30 s recovery intervals (15) between half squat repetitions (i.e. CS).

After the end of the half squat exercise, participants performed 2 CMJs separated by ~15 s, at 1, 3, 6, 9, and 12 minutes of recovery. They were instructed to jump as high as possible. The CMJs were performed on a force plate (Quattro Jump, Kistler, Switzerland) where vertical forces were recorded with a sampling rate of 500 Hz. Participants freely chose the depth of the countermovement. Based on a previous study (3), the kinetic parameters selected for analysis were: peak power ($\text{W} \cdot \text{kg}^{-1}$), maximum force relative to body weight (BW), normalized stiffness ($\text{N} \cdot \text{m}^{-1} \cdot \text{kg}^{-1}$), and vertical displacement of center of mass during countermovement (cm). Peak power and stiffness were normalized to provide a better comparison between participants of differing body mass (see Table 1).

Statistical Analysis

Descriptive data are presented as mean $\pm$ SD. The normality distribution of variables was examined with the Kolmogorov-Smirnov test. Effects of each conditioning protocol on jump-

related parameters at different times were assessed using analysis of variance (ANOVA) for repeated measurements (2 conditioning protocols $\times$ 6 time points) with pairwise Bonferroni correction. Baseline values were further compared with the peak of each dependent variable, regardless at which time the peak occurred (4), by means of paired Student's t-test. Comparisons between measurements at peak height time and baseline were also performed. The magnitude of baseline-to-peak changes was assessed using effect size (ES) and percentage of change. Threshold values for ES were 0.2 (small), 0.6 (moderate), 1.2 (large), and 2.0 (very large) (14). Confidence intervals (90%) for the true mean change were also estimated. Magnitude-based inferences about the true change were made with reference to the smallest worthwhile change calculated as: 0.2 multiplied by the between-subject standard deviation expressed as coefficient of variation (CV %). Quantitative chances of substantial positive, trivial or negative changes were assessed qualitatively as follows: <0.5%, almost certainly not; 0.5–5%, very unlikely; 5–25%, unlikely; 25–75%, possibly; 75–95%, likely; 95–99.5%, very likely; >99.5% almost certainly. If the chance of having positive or negative changes were both >5%, the true difference was deemed unclear (14). Relationships between parameters were identified using Pearson moment product correlation coefficient (r). Confidence intervals (90%) for coefficients of correlation were also estimated. The following criteria were adopted for interpreting the magnitude of correlations between parameters: <0.1, trivial; 0.1–0.3 small; 0.3–0.5 moderate; 0.5–0.7, large; 0.7–0.9, very large; 0.9–1.0, almost perfect. If the 90% confidence intervals overlapped the thresholds for substantially positive or negative values, the magnitude was considered unclear. Inferences about correlations were made with respect to a smallest worthwhile correlation of 0.1. Differences in time to peak for each jump-related parameter was

analyzed by paired Student's t-test with a complementary ES calculation. Statistical significant level was set at a p value of <0.05 . Statistical analysis was conducted with the statistical package for the social sciences software (v17.0, SPSS Inc., Chicago, WI, USA).

ACCEPTED

RESULTS

The ANOVA revealed no significant main effects for jump-related parameters. Peak power showed a tendency for the time factor ($p=0.051$), and a significant interaction between the protocol and time factors ($p=0.048$). Simple effect analyses revealed that peak power was higher at 1 min after CS when compared to the traditional set configuration (51.8 ± 5.5 vs. 50.9 ± 4.6 $\text{W} \cdot \text{kg}^{-1}$; $p=0.046$). Conversely, peak power was lower at 9 min after CS when compared to the traditional set configuration (50.5 ± 5.9 vs. 51.8 ± 5.5 $\text{W} \cdot \text{kg}^{-1}$; $p=0.039$). No other significant interactions were observed.

Peak values of jump height and force-time parameters during CMJ, independently of the time at which they occurred are presented on Table 2A. Interestingly, the majority of the parameters improved significantly except peak force and vertical stiffness after traditional set configuration.

Place Table 2 about here

Force-time parameters associated with the best jump height are presented in Table 2B. Despite any significant change detected, it is interesting to note an opposite trend in vertical stiffness and vertical displacement of the center of mass between conditions as inferred from ES and qualitative outcomes. Thus, cluster set configuration demonstrated an improvement in vertical stiffness with a lowering in the depth of the countermovement, whereas traditional set configuration showed an impairment in vertical stiffness with an increment in the depth of the countermovement.

Correlations among parameters at jump peak height are shown in Table 3. Surprisingly, Δ peak power was not correlated with Δ jump height in any condition but exhibited a correlation with Δ vertical stiffness in both conditions. Moreover, Δ vertical stiffness was correlated with Δ peak force only in the cluster condition. Additionally, an inverse relationship between Δ vertical stiffness and the Δ vertical displacement of the center of mass in both conditions was observed.

Place Table 3 about here

The time to peak of all jump parameters are presented on Table 4. Interestingly, a more rapid increment in jump height was observed in the cluster condition when compared to the traditional set configuration ($p=0.042$), with a tendency detected ($p=0.052$) for the vertical displacement of the center of mass. Additionally, various correlations were detected among strength and peak force-time parameters but only in the cluster condition (5RM/body mass - Δ vertical stiffness, $r = -0.839$; $p=0.005$; 5RM/body mass - Δ peak Force, $r = -0.645$; $p=0.061$; 5RM - time to peak value of Vertical displacement of center of mass, $r = -0.602$; $p=0.039$).

Place Table 4 about here

DISCUSSION

The main finding of the current study was that different set configurations have different acute effects on potentiation while jumping. Thus, introducing recovery intervals between half squat repetitions (i.e. cluster training) seems to allow a more rapid improvement of jump height, and a greater potentiation of various force-time parameters when compared to a set until failure, probably as a consequence of the better fatigue-potentiation relationship in the cluster condition.

These findings are of great interest for athletes and coaches that would like to optimize their training routines as well their competitive warm-ups on an individual basis.

In line with the recent work of Chaouachi et al. (4), the potentiation of force-time parameters in the current study was also variable among individuals. In contrast, our protocols could induce significant improvements in peak jump height while these authors (4) did not observe any significant change in peak jump height independently of the time considered. Moreover, while the ANOVA of this previous study (4) showed an impairment of jump capacity after all protocols, we did not find any significant change in jump height with the ANOVA. Previous studies have suggested that factors like training experience and exercise load could modulate the potentiation-fatigue interaction for subsequent performance (22, 2). Thus, it is interesting to note that a similar load (i.e. 5RM ~ 85%RM) as employed in the previous study (4) (i.e. 70 to 90% RM) elicited better performance in the current study. This intriguing result is in line with previous literature in which the potentiation response after similar protocols was not consistent which could be related to participants' characteristics and training background (29, 5, 19). In this respect, our participants could not be considered elite level athletes since they trained concurrently antagonistic physical capabilities such as endurance and explosiveness. Hence, a lower potentiation response could be expected in the current study (5, 19). While we do not know the exact reason for such a difference, it is tempting to speculate that our protocols may induce a better fatigue-potentiation relationship as the half squats were performed until 90° knee flexion, while the participants of the previous study performed parallel squats (4). This hypothesis could be supported by the previous report of Place et al. (20) that showed a greater post-twitch potentiation after fatiguing contractions at short quadriceps muscle lengths when

compared to long muscle lengths but with a similar level of fatigue between conditions. Similarly, it may be suggested that participants of our study could experience a greater potentiation after squats because of the lower level of knee flexion. Therefore, further studies should elaborate on the potentiation-fatigue response to various exercises and workloads, and more specifically with regard to the level of knee flexion during squats.

Interestingly, the different levels of potentiation on force-time parameters after both protocols (failure vs. cluster) allowed a similar magnitude of peak height increments. However, the difference between conditions was evident on the significantly different time to peak height (3.6 ± 2.9 vs. 6.1 ± 3.3 min, for cluster and failure set, respectively; $p=0.042$). Furthermore, the timing of the appearance of peak values of selected parameters was protocol dependent with the traditional set configuration exhibiting a lower synchronization among parameters (see Table 5). This could reflect different jump coordination between conditions (23, 4). For instance, the peak values for the vertical displacement of the center of mass occurred at very different time periods (8.7 ± 3.5 vs. 5.8 ± 2.4 min; $p=0.052$, for traditional and cluster set configuration, respectively), with the direction on such changes being opposite. Moreover, the absence of correlations between Δ peak power and Δ jump height is surprising and may indicate an altered motor control after both protocols as these parameters were highly correlated at baseline (i.e. 0.754 and 0.797, $p<0.01$; for traditional and cluster set configuration, respectively). In this regard, the differences revealed by ANOVA in peak power at 1 and 9 min of recovery between conditions reinforce this assumption. Since the intensity of both protocols could be of similar magnitude, the inclusion of rest intervals between repetitions in the cluster condition may be considered the major factor for such differences between protocols. Thus, the fatigue effects possibly caused by metabolic

byproducts as well as greater neural fatigue may have delayed this potentiation response (1) after the traditional set configuration as similar volumes resulted in similar net potentiation of peak power in both conditions. Additionally, it should be noted that the completion time between the first and the last repetition in each condition was quite different (e.g. ~2 min 30 s vs. ~15 s). This consideration is important when performing multiple sets sessions. Further studies should elaborate on this observation of a delayed appearance of potentiation after traditional set configuration with different training volumes (e.g. number of sets), exercise intensities (e.g. 3RM), and other exercises (e.g. bench press).

These differences in correlations among parameters between conditions may be also influenced by the fact that force-time parameters like vertical stiffness and peak force are recorded at the end of the eccentric action, while peak power is recorded near the end of the take off (6), thus suggesting an elastic energy transfer (28) that could be also influenced by the potentiation-fatigue interaction (3). This consideration is important as the interaction of these phenomena could be depending upon the muscular regime and/or the muscle length (21). The existence of an elastic energy transfer could be supported by the high correlations observed (see Table 4) between Δ peak power and Δ vertical stiffness in both conditions. In this regard, Vuorimaa et al. (27) previously observed a lowering in EMG activity with a concurrent increment in mean power during half-squat exercises thus suggesting a possible increase in elastic energy transfer after various running-conditioning protocols. More recently, Moir et al. (18) reported a greater stiffness increment after high intensity when compared to high volume back squats but with no augmentation of the jump height. Therefore, further studies should be conducted for the assessment of EMG concurrently with kinetic parameters for evaluating the

influence of the fatigue-potential relationship on the elastic energy transfer and jump coordination after different conditioning protocols.

Of further interest are the correlations found among half squat performance and force-time parameters only after the cluster protocol. This is a novel finding with our results suggesting that the stronger the athlete, the lower and the more anticipated was the increase of force-time parameters over the eccentric movement of the push-off phase. Since these findings are related only to the cluster condition, it may be speculated that a different response could occur with different loads (e.g. heavier loads). Thus, it may be suggested that a heavier load could induce a greater response in the stronger athletes as the load of 5RM may not be the optimal stimulus for them with such a load. Conversely, the absence of correlations among these parameters after the traditional set configuration may suggest that the possible influence of strength level on such relationships could be masked by other factors such as fatigue resistance (4). In this respect, it should be considered that the training background of our participants do not favor maximum power performances but a greater endurance capacity. **Therefore, it is possible that our specific participants (i.e. aspiring fire-fighters) may possess a greater recovery capacity when compared to other athletes with greater power but lower endurance capacity.** Further studies could explore the influence of strength levels concurrently with fatigue resistance with respect to the load and set configuration of conditioning activities.

In summary, a set with the load equivalent to 5RM of half-squats performed with inter-repetition rest intervals of 30 s favors a more rapid appearance, and a greater potentiation of various kinetic parameters during vertical jumping, when compared to a set of 5RM until failure (without rest intervals). Further studies are needed for testing the effectiveness of cluster training

in other exercises, for competition, as well as the chronic effect of complex training performed with CS on performance (8). Furthermore, additional variations of CS (e.g. inclusion of dynamic stretching and/or altering pause duration) should be examined regarding their relative potentiation effects on subsequent exercises by individuals with designated training backgrounds. Additionally, more studies are needed with greater training volumes and with other intensities for determining the best dose-response in during conditioning activities.

ACCEPTED

PRACTICAL APPLICATIONS

The optimal method for inducing peak power potentiation is dependent on the available time between heavy half-squat exercise and subsequent jump performance. It appears that CS may be a superior approach to a 5RM approach prior to jump performance if the time available is close to 1 min, whereas the 5RM approach may be superior if close to 9 min is available for individuals trained in both power and endurance capacities who have been engaged in heavy-resistance training for no less than the previous 6 months. Additionally, it may be suggested that the employment of force plates is recommended for a more precise evaluation of jump potentiation since the potentiation of various kinetics parameters at different times could be expected with no observable increment in jump height. This approach could be interesting for determining the appropriate workloads for warming up in different sports as for designing complex training sessions on an individual basis.

REFERENCES

1. Behm, DG, Button, DC, Barbour, G, Butt, JC, and Young, WB. Conflicting effects of fatigue and potentiation on voluntary force. *J Strength Cond Res* 18(2):365-372, 2004.
2. Boullosa, DA, and Tuimil, JL. Postactivation potentiation in distance runners after two different field running protocols. *J Strength Cond Res* 23(5):1560-1565, 2009.
3. Boullosa, DA, Tuimil, JL, Alegre, LM, Iglesias, E, and Lusquiños, F. Concurrent fatigue and potentiation in endurance athletes. *Int J Sports Physiol Perform* 6(1):82-93, 2011.
4. Chaouachi, A, Poulos, N, Abed, F, Turki, O, Brughelli, M, Chamari, K, Drinkwater, EJ, and Behm, DG. Volume, intensity, and timing of muscle power potentiation are variable. *Appl Physiol Nutr Metab* 36(5):736-747, 2011.
5. Chiu, LZ, Fry, AC, Weiss, LW, Schilling, BK, Brown, LE, and Smith, SL. Postactivation potentiation response in athletic and recreationally trained individuals. *J Strength Cond Res* 17(4):671-677, 2003.
6. Cormie, P, McBride, JM, and McCaulley, GO. Power-time, force-time, and velocity-time curve analysis of the countermovement jump: impact of training. *J Strength Cond Res* 23(1):177-186, 2009.
7. Denton, J, and Cronin, JB. Kinematic, kinetic and blood lactate profiles of continuous and intraset rest loading schemes. *J Strength Cond Res* 20(3):528-534, 2006.
8. Docherty, D, and Hodgson, MJ. The application of postactivation potentiation to elite sport. *Int J Sports Physiol Perform* 2(4):439-444, 2007.

9. Haff, GG, Whitley, A, McCoy, LB, O'Bryant HS, Kilgore JL, Haff EE, Pierce K, and Stone MH. Effects of different set configurations on barbell velocity and displacement during a clean pull. *J Strength Cond Res* 17(1):95-103, 2003.
10. Hansen, KT, Cronin, JB, and Newton, MJ. The effect of cluster loading on force, velocity, and power during ballistic jump squat training. *Int J Sports Physiol Perform* 6(4):455-468, 2011.
11. Hardee, JP, Triplett, NT, Utter, AC, Zwetsloot KA, and McBride JM. Effect of inter-repetition rest on power output in the power clean. *J Strength Cond Res* 26(4):883-889, 2012.
12. Hardee, JP, Lawrence, MM, Utter, AC, Triplett, NT, Zwetsloot, KA, and McBride, JM. Effect of inter-repetition rest on ratings of perceived exertion during multiple sets of the power clean. *Eur J Appl Physiol* 112(8):3141-3147, 2012.
13. Hodgson, M, Docherty, D, and Robbins, D. Post-activation potentiation: underlying physiology and implications for motor performance. *Sports Med* 35(7):585-595, 2005.
14. Hopkins, WG, Marshall, SW, Batterham, AM, and Hanin J. Progressive Statistics for Studies in Sports Medicine and Exercise Science. *Med Sci Sports Exerc* 41 (1): 3-13, 2009.
15. Iglesias, E, Boulosa, DA, Dopico, X, and Carballeira E. Analysis of factors that influence the maximum number of repetitions in two upper-body resistance exercises: curl biceps and bench press. *J Strength Cond Res* 24(6):1566-1572, 2010.

16. Khamoui, AV, Brown, LE, Coburn, JW, Judelson, DA, Uribe, BP, Nguyen, D, Tran, T, Eurich, AD, and Noffal, GJ. Effect of potentiating exercise volume on vertical jump parameters in recreationally trained men. *J Strength Cond Res* 23(5):1465-1469, 2009.
17. Lawton, TW, Cronin, JB, and Lindsell, RP. Effect of interrepetition rest intervals on weight training repetition power output. *J Strength Cond Res* 20(1):172-176, 2006.
18. Moir, GL, Mergy, D, Witmer, C, and Davis, SE. The acute effects of manipulating volume and load of back squats on countermovement vertical jump performance. *J Strength Cond Res* 25(6):1486-1491, 2011.
19. Morana, C, and Perrey, S. Time course of postactivation potentiation during intermittent submaximal fatiguing contractions in endurance- and power-trained athletes. *J Strength Cond Res* 23(5):1456-1464, 2009.
20. Place, N, Maffiuletti, NA, Ballay, Y, and Lepers, R. Twitch potentiation is greater after a fatiguing submaximal isometric contraction performed at short vs. long quadriceps muscle length. *J Appl Physiol* 98(2):429-436, 2005.
21. Rijkkelikhuisen, JM, de Ruiter, CJ, Huijng, PA, and Haan, A. Low-frequency fatigue, post-tetanic potentiation and their interaction at different muscle lengths following eccentric exercise. *J Exp Biol* 208(Pt 1):55-63, 2005.
22. Rixon, KP, Lamont, HS, and Bemben, MG. Influence of type of muscle contraction, gender, and lifting experience on postactivation potentiation performance. *J Strength Cond Res* 21(2):500-505, 2007.

23. Rodacki, AL, Fowler, NE, and Bennett, SJ. Vertical jump coordination: fatigue effects. *Med Sci Sports Exerc* 34(1):105-116, 2002.
24. Sale, DG. Postactivation potentiation: role in human performance. *Exerc Sport Sci Rev* 30(3):138-143, 2002.
25. Tillin, NA, and Bishop, D. Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sports Med* 39(2):147-166, 2009.
26. Tran, QT, Docherty, D, and Behm, D. The effects of varying time under tension and volume load on acute neuromuscular responses. *Eur J Appl Physiol* 98(4):402-410, 2006.
27. Vuorimaa, T, Virlander, R, Kurkilahti, P, Vasankari T, and Häkkinen K. Acute changes in muscle activation and leg extension performance after different running exercises in elite long distance runners. *Eur J Appl Physiol* 96(3):282-291, 2006.
28. Wilson, JM, and Flanagan, EP. The role of elastic energy in activities with high force and power requirements: a brief review. *J Strength Cond Res* 22(5):1705-1715, 2008.
29. Young, WB, Jenner, A, and Griffiths, K. Acute enhancement of power performance from heavy load squats. *J Strength Cond Res* 12(2): 82-84, 1998.

Acknowledgements

We would like to thank Dr. Miguel Fernández-del-Olmo for his support for data collection. We would like to recognize the facilities provided by Gimnasio Athenas C.B. during the experiment. We also thank the athletes for their participation. The results of the present study do not constitute endorsement by the authors or the NSCA.

ACCEPTED

TABLE 1. Participants' characteristics.

	Mean $\pm$ SD
Age (years)	25.5 $\pm$ 4.9
Height (cm)	178.2 $\pm$ 8.3
Body mass (kg)	77.7 $\pm$ 7.8
Body fat (%)	9.6 $\pm$ 2.2
5 RM half squat (kg)	181.4 $\pm$ 24.1
5 RM half squat/Body Mass	2.4 $\pm$ 0.3

RM, repetitions maximum.

TABLE 2. Force-time parameters of the CMJ after isolating the peak of each dependent variable and comparing it with the baseline, regardless of at which time point the peak occurred (A), and force-time parameters associated with the best jump height (B).

A	Condition	Baseline	Peak	p	%Δ; ±90% CL	ES	Qualitative inference
CMJ (cm)							
	Traditional	44.68 ± 4.47	45.85 ± 4.44	0.030	2.70; ±1.9	0.26	possibly
	Cluster	45.12 ± 4.60	46.29 ± 5.02	0.019	2.57; ±1.7	0.24	possibly
Peak power (W·kg⁻¹)							
	Traditional	52.12 ± 5.02	53.39 ± 5.83	0.031	2.36; ±1.7	0.23	possibly
	Cluster	51.27 ± 5.87	52.45 ± 5.58	0.025	2.41; ±1.8	0.20	possibly
Peak force (BW)							
	Traditional	2.37 ± 0.21	2.46 ± 0.25	0.121	3.99; ±4.3	0.39	possibly
	Cluster	2.31 ± 0.28	2.48 ± 0.26	0.004	7.48; ±3.6	0.61	very likely
Vertical displacement of center of mass (cm)							
	Traditional	29.91 ± 4.11	33.27 ± 4.71	0.003	11.67; ±5.5	0.76	very likely
	Cluster	31.64 ± 4.74	34.18 ± 4.26	0.003	8.69; ±4.1	0.56	very likely
Vertical stiffness (N·m⁻¹·kg⁻¹)							
	Traditional	81.99 ± 10.93	88.36 ± 17.61	0.133	7.70; ±8.5	0.43	likely positive
	Cluster	76.07 ± 14.35	88.39 ± 19.69	0.002	16.02; ±7.1	0.72	very likely

B	Condition	Baseline	Values at peak height	p	%Δ; ±90% CL	ES	Qualitative inference
Peak power (W·kg⁻¹)							
	Traditional	52.12 ± 5.02	52.53 ± 6.08	0.544	0.69; ±2.0	0.07	likely trivial
	Cluster	51.27 ± 5.87	51.38 ± 5.71	0.752	0.27; ±1.5	0.02	very likely trivial
Peak force (BW)							
	Traditional	2.37 ± 0.21	2.37 ± 0.23	0.937	-0.014; ±0.32	0.02	most likely trivial
	Cluster	2.31 ± 0.28	2.35 ± 0.27	0.534	2.02; ±5.7	0.14	unclear
Vertical displacement of center of mass (cm)							
	Traditional	29.91 ± 4.11	31.32 ± 5.26	0.226	4.99; ±7.0	0.30	possibly
	Cluster	31.64 ± 4.74	30.64 ± 4.50	0.184	-2.87; ±3.6	0.22	possibly trivial
Vertical stiffness (N·m⁻¹·kg⁻¹)							
	Traditional	81.99 ± 10.93	79.07 ± 14.40	0.430	-3.33; ±7.3	0.23	unclear
	Cluster	76.07 ± 14.35	79.72 ± 14.75	0.220	5.58; ±7.7	0.25	possibly

Values are mean ± SD. Magnitudes of differences are expressed as mean percentage change (%Δ) and 90% confidence limits (±90% CL). ES, effect size.

TABLE 3. Correlations among parameters at jump peak height. Pearson's r (with 90% confidence intervals), p-values and qualitative inferences are shown.

CLUSTER condition	Δ Peak Power	Δ Peak Force	Δ Vertical Stiffness	Δ vertical displacement of center of mass
Δ CMJ	r = 0.21 (□0.32;0.64)	0.45 (□0.06;0.78)	0.10 (□0.42;0.57)	0.41 (□0.12;0.75)
	p =0.530	0.161	0.771	0.215
	unclear	likely	unclear	unclear
Δ Peak Power		0.51 (0.02;0.81)	0.69 (0.29;0.88)	□0.48 (□0.79;0.03)
		0.108	0.020	0.135
		likely	very likely	likely
Δ Peak Force			0.79 (0.48;0.92)	□0.03 (□0.52;0.48)
			0.004	0.940
			most likely	unclear
Δ Vertical Stiffness				□0.63 (□0.86;□0.19)
				0.038
				very likely

TRADITIONAL condition	Δ Peak Power	Δ Peak Force	Δ Vertical Stiffness	Δ vertical displacement of center of mass
Δ CMJ	r = □0.09 (□0.56;0.43) p =0.793 unclear	0.31 (□0.23;0.70) unclear	□0.26 (□0.67;0.27) unclear	0.43 (□0.10;0.78) unclear
Δ Peak Power		0.51 (0.01;0.80) 0.112 likely	0.94 (0.83;0.98) 0.000 most likely	□0.82 (□0.94;□0.55) 0.002 most likely
Δ Peak force			0.51 (0.01;0.80) 0.112 likely	□0.07 (□0.55;0.44) 0.826 unclear
Δ Vertical Stiffness				□0.89 (□0.96;□0.71) 0.000 most likely

CMJ, countermovement jump; Δ, change from baseline to the value associated with the best jump height.

TABLE 4. Time (min) to peak data of jump height and force-time parameters, and statistical differences (p values and effect size) among these parameters between conditions and during every condition.

	Minute of peak value (mean $\pm$ SD)		p	ES
	Traditional protocol	Cluster protocol		
CMJ height	6.1 $\pm$ 3.3	3.6 $\pm$ 2.9	0.042	0.84
Peak Power	4.4 $\pm$ 2.5	4.4 $\pm$ 4.7	0.951	0.03
Peak Force	4.9 $\pm$ 4.2	4.4 $\pm$ 4.0	0.537	0.20
Vertical stiffness	4.1 $\pm$ 3.5	5.7 $\pm$ 4.5	0.249	0.32
Vertical displacement of center of mass	8.7 $\pm$ 3.5	5.8 $\pm$ 2.4	0.052	0.83

Traditional protocol [p value (ES)]				
	Peak Power	Peak Force	Vertical stiffness	Vertical displacement of center of mass
CMJ height	0.054 (0.61)	0.279 (0.32)	0.205 (0.60)	0.049 (0.75)
Peak Power		0.550 (0.16)	0.833 (0.09)	0.020 (1.44)
Peak Force			0.547 (0.24)	0.060 (0.98)
Vertical stiffness				0.034 (1.31)

Cluster protocol [p value (ES)]				
	Peak Power	Peak Force	Vertical stiffness	Vertical displacement of center of mass
CMJ height	0.701 (0.19)	0.683 (0.16)	0.077 (0.71)	0.154 (0.67)
Peak Power		0.892 (0.04)	0.276 (0.44)	0.433 (0.31)
Peak Force			0.065 (0.51)	0.368 (0.39)
Vertical stiffness				0.665 (0.23)

CMJ, countermovement jump; ES, effect size.