

training loads and injury risk,³ and so we have also been considering the best way to model this relationship. We share Dr Menaspà's concerns regarding the use of rolling averages for the calculation of 'acute' and 'chronic' loads. Namely, that they fail to account for the decaying nature of fitness and fatigue effects over time⁴ and do not accurately represent variations in the manner in which loads are accumulated (as demonstrated in the example data presented by Dr Menaspà²). To mitigate some of these issues, we propose the use of 'exponentially weighted moving averages (EWMA)'⁵ for the calculation of acute and chronic loads, which assign a decreasing weighting for each older load value. Specifically, the EWMA for a given day is given by

$$\text{EWMA}_{\text{today}} = \text{Load}_{\text{today}} \times \lambda_a + ((1 - \lambda_a) \times \text{EWMA}_{\text{yesterday}})$$

where λ_a is a value between 0 and 1 that represents the degree of decay, with higher values discounting older observations at a faster rate. The λ_a is given by

$$\lambda_a = 2/(N + 1)$$


where N is the chosen time decay constant, typically 7 and 28 days for acute ('fatigue') and chronic ('fitness') loads, respectively. One-week and 4-week time frames appear to align well with the periodisation strategies used in many team sports, although alternative time constants may be more appropriate in different settings. Applying this method to the example data presented by Dr Menaspà² produces a different acute:chronic workload on day 28 for each of the three fictitious athletes (1.25, 1.41 and 1.55, respectively), whereas the use of rolling averages produces three identical values (1.43). Thus, in the case of athlete 1, the two approaches differ with regard to whether the athlete is considered to be within or beyond the 'sweet spot' region of 0.8–1.3.⁶ Using athlete 3 as an illustrative example ([figure 1](#)), in comparison to rolling averages the EWMA approach gives more weighting to the high loads undertaken towards the end of the 28 day period (when estimating 'fatigue') and so produces a higher (and we propose, more appropriate) acute:chronic workload ratio on day 28. Thus, the EWMA approach may be better suited to the modelling of training loads than rolling averages, and so we believe this method warrants consideration in future research and practice.

CORRESPONDENCE

Better way to determine the acute:chronic workload ratio?

We read with great interest the recent letter, "Time to bin the term 'overuse' injury: is 'training load error' a more accurate term?"¹ and in particular its associated PostScript correspondence, "Are rolling averages a good way to assess training load for injury prevention?"² We are currently investigating the association between

Figure 1 A demonstration of the differing acute:chronic workload ratio values produced when using the EWMA and rolling average methods. EWMA values were initialised with the load value for day 1 and used time decay constants of 7 and 28 days for acute and chronic loads, respectively. EWMA, exponentially weighted moving averages.



Sean Williams,¹ Stephen West,¹
Matthew J Cross,² Keith A Stokes¹

¹Department for Health, University of Bath, Bath, UK
²Rugby Football Union, Twickenham, UK

Correspondence to Dr Sean Williams, Department for Health, University of Bath, Bath BA2 7AY, UK; S.Williams@bath.ac.uk

Twitter Follow Sean Williams at @statman_sean, Stephen West at @westy160991 and Keith Stokes at @drkeithstokes

Contributors SWi and SWe came up with the idea for this correspondence piece. SWi prepared the first draft manuscript. All authors provided feedback and helped to revise the manuscript.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.



CrossMark

To cite Williams S, West S, Cross M J, et al. *Br J Sports Med* 2017;**51**:209–210.

Accepted 30 August 2016
Published Online First 20 September 2016

Br J Sports Med 2017;**51**:209–210.
doi:10.1136/bjsports-2016-096589

REFERENCES

- 1 Drew MK, Purdam C. Letter: Time to bin the term 'overuse' injury: is 'training load error' a

more accurate term? *Br J Sports Med* 2016;50:1423–4.

- 2 Menaspà P. Correspondence: Are rolling averages a good way to assess training load for injury prevention? *Br J Sports Med* Published Online First: 24 May 2016 doi:10.1136/bjsports-2016-096131
- 3 Cross MJ, Williams S, Trewartha G, et al. The influence of in-season training loads on injury risk in professional Rugby Union. *Int J Sports Physiol Perform* 2016;11:350–5.
- 4 Hawley JA. Adaptations of skeletal muscle to prolonged, intense endurance training. *Clin Exp Pharmacol Physiol* 2002;29:218–22.
- 5 Hunter JS. The exponentially weighted moving average. *J Quality Technol* 1986;18:203–10.
- 6 Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50:273–80.