

Oxytocin Nasal Spray

Neuropeptide hormone · Intranasal · Bonding / mood / sexual function

Nasal Spray

PRN or Daily

Refrigerated

Off-Label Use

COMMON INDICATIONS (OFF-LABEL)

- ▶ Anxiety / social anxiety
- ▶ Sexual dysfunction (M/F)
- ▶ Autism social adjunct
- ▶ Mood support
- ▶ Libido / arousal support
- ▶ PTSD adjunct
- ▶ Bonding / relational health
- ▶ Orgasmic dysfunction
- ▶ Postpartum bonding

STANDARD DOSING PROTOCOLS

STARTING

10 IU

1 spray PRN

STANDARD

20–40 IU

Daily or PRN

SEXUAL USE

20–40 IU

30–60 min before

MAXIMUM

80 IU

Per day

DOSING NOTES

Typical concentration is 40 IU/mL with a metered spray delivering ~10 IU per spray (verify on dispensed label). For social/anxiety/bonding use, dose 30–60 min before the social context or daily AM. For sexual function, 30–60 min before activity. Effects last ~30–60 minutes. Prime nasal sprays before first use; alternate nostrils. Can be paired with PT-141 for sexual dysfunction protocols.

DOSING BY INDICATION

INDICATION	TYPICAL DOSE	FREQUENCY / NOTES
Anxiety / social context	10–20 IU	30–60 min before situation; PRN
Daily mood / bonding	20–40 IU	Daily AM; assess at 4 weeks
Sexual function (M/F)	20–40 IU	30–60 min before activity
PTSD / autism adjunct	20–40 IU	Twice daily, specialist supervision
Maximum daily	Up to 80 IU	Do not exceed without specialist input

STORAGE & COUNSELING

STORAGE

- ▶ Refrigerate at 36–46°F (2–8°C) before opening
- ▶ After opening, may store at room temp per BUD
- ▶ Protect from heat and light
- ▶ Discard if cloudy or discolored

COUNSELING PEARLS

- ▶ Prime spray before first use; tilt head slightly forward
- ▶ Avoid sniffing forcefully — let absorb in nasal mucosa
- ▶ Common: nasal irritation, mild headache, dry mouth
- ▶ Caution: pregnancy, hyponatremia, fluid retention