

# Tesamorelin

GHRH analog · 5 mg/mL injectable · Visceral fat reduction / GH support

SubQ Injection

Daily QHS

5 mg/mL

Visceral Fat Focus

## COMMON INDICATIONS

- ▶ HIV-related lipodystrophy (FDA)
- ▶ Visceral adiposity
- ▶ Abdominal fat reduction
- ▶ Adult GH decline
- ▶ Body composition support
- ▶ Metabolic syndrome adjunct
- ▶ NAFLD / hepatic fat
- ▶ Cognitive support
- ▶ Athletic / longevity

## STANDARD DOSING PROTOCOL (5 MG/ML)

STANDARD

**1 mg**

Daily SubQ QHS

HIGHER DOSE

**2 mg**

Daily SubQ QHS

CYCLE LENGTH

**12–24 wks**

Then reassess

MAINTENANCE

**1–2 mg**

Daily ongoing

## DOSING NOTES

Tesamorelin is a stabilized GHRH analog — stronger and longer-acting than sermorelin, with more pronounced effects on visceral fat. FDA-approved dose is 2 mg SubQ daily for HIV-associated lipodystrophy; off-label use commonly starts at 1 mg daily and titrates as tolerated. Inject SubQ in abdomen at bedtime, on empty stomach (food blunts GH release). Visceral fat reduction is typically measurable at 12–26 weeks. Monitor IGF-1 to keep within physiologic range.

## VOLUME REFERENCE (5 MG/ML)

| TARGET DOSE | VOLUME TO DRAW | U-100 SYRINGE | NOTES                         |
|-------------|----------------|---------------|-------------------------------|
| 0.5 mg      | 0.10 mL        | 10 units      | Sensitive patients, starting  |
| 1.0 mg      | 0.20 mL        | 20 units      | Standard starting / off-label |
| 1.4 mg      | 0.28 mL        | 28 units      | Intermediate dose             |
| 2.0 mg      | 0.40 mL        | 40 units      | FDA-approved standard dose    |

## MONITORING & CAUTIONS

### MONITORING

- ▶ Baseline IGF-1; recheck at 3 months, then q6 mo
- ▶ Fasting glucose / HbA1c (may ↑ insulin resistance)
- ▶ Lipid panel at baseline and 6 months
- ▶ Waist circumference / body composition q3 months

### CAUTIONS

- ▶ Contraindicated: active malignancy, hypopituitarism, pregnancy
- ▶ Caution: diabetes (may worsen glycemic control)
- ▶ Common: injection site reaction, joint pain, fluid retention
- ▶ Hold if IGF-1 rises >3 SD above age-adjusted mean