

5-Amino-1MQ

Selective NNMT inhibitor · Oral capsule · Metabolic / cellular optimization

Oral Capsule

Once Daily AM

Cycle Therapy

Off-Label / Investigational

COMMON INDICATIONS (OFF-LABEL)

- ▶ Metabolic optimization
- ▶ Adipose tissue reduction
- ▶ Body recomposition
- ▶ NAD+ pathway support
- ▶ Mitochondrial function
- ▶ Cellular energy / longevity
- ▶ Adjunct to GLP-1 therapy
- ▶ Plateaued weight loss
- ▶ Insulin sensitivity support

STANDARD DOSING & CYCLE

WEEK 1

50 mg

Once daily AM

WEEKS 2–4

50–100 mg

Once daily AM

CYCLE LENGTH

8–12 wks

Then break

OFF CYCLE

2–4 wks

Reset NNMT

DOSING NOTES

Take in the morning or early afternoon — late dosing may interfere with sleep due to metabolic activation. Capsules can be taken with or without food. Many protocols split daily dose (50 mg AM + 50 mg early PM) for steadier exposure. Reassess at end of cycle; intermittent off-cycle periods are recommended given NNMT enzymatic adaptation. Limited human clinical trial data — dosing is based on observational protocols.

DOSING BY GOAL

GOAL	TYPICAL DOSE	NOTES
Metabolic / fat loss support	50–100 mg daily AM	8–12 week cycles, 4-week break
Adjunct to GLP-1 plateau	50 mg daily AM	Lower dose; monitor response 4–6 wks
NAD+ / cellular health	50 mg daily AM	Often paired with NAD+ injection protocol
Sensitive patients / starting	50 mg every other day	Build up over 1–2 weeks if tolerated

AVAILABLE FORMULATION

CAPSULE STRENGTHS

- ▶ **50 mg capsules** — standard maintenance dose
- ▶ **100 mg capsules** — established users / higher-dose protocols
- ▶ Custom strengths available on request

CAUTIONS

- ▶ Not for use in pregnancy or lactation
- ▶ Caution with stimulants, BP and diabetes meds
- ▶ Avoid in patients under 21 or with eating disorder hx
- ▶ Investigational — limited human safety data