

Selank Acetate (TP-7)

Heptapeptide tuftsin analog · Nasal spray (7,500 mcg/mL) · Anxiolytic / nootropic

Nasal Spray

Daily Cycle

7,500 mcg/mL

Investigational

COMMON INDICATIONS (INVESTIGATIONAL)

- ▶ Generalized anxiety
- ▶ Situational anxiety
- ▶ Stress / overwhelm
- ▶ Cognitive enhancement
- ▶ Focus / mental clarity
- ▶ Brain fog adjunct
- ▶ Asthenia / low energy
- ▶ Mild depression adjunct
- ▶ Benzodiazepine alternative

STANDARD DOSING PROTOCOL (7,500 MCG/ML)

STARTING

300 mcg

Daily ×3–7 days

STANDARD

750 mcg

Daily, divided

HIGHER DOSE

Up to 900 mcg

Daily, divided

CYCLE

10–14 days

Then 1–2 wk break

DOSING NOTES

At 7,500 mcg/mL concentration, a 100-mL metered spray delivers ~750 mcg per spray. Most common protocol: 1 spray (~750 mcg) per nostril once daily, OR split into 2–3 smaller doses for steadier effect. Russian clinical trials supporting use are intranasal — bioavailability via nasal mucosa bypasses GI breakdown. Cycles of 10–14 days are typical; longer cycles up to 4 weeks are reported. **Investigational in the US.**

DOSING BY INDICATION

INDICATION	TYPICAL DAILY DOSE	NOTES
Mild anxiety / stress	300–500 mcg/day	1 spray AM, often sufficient
Standard anxiety / GAD	750 mcg/day	1 spray each nostril daily
Cognitive / nootropic	500–900 mcg/day	Split AM/midday
Acute / situational	300–750 mcg PRN	30–60 min before stressor
Maximum reported	~1,000 mcg/day	Reserve for specialist guidance

Verify spray volume per actuation on dispensed bottle label; spray output may vary by device.

STORAGE & COUNSELING

STORAGE

- ▶ Refrigerate at 36–46°F (2–8°C)
- ▶ Protect from light
- ▶ BUD per pharmacy label

CAUTIONS

- ▶ Mild nasal irritation, occasional headache
- ▶ Caution with SSRIs / SNRIs (limited data)
- ▶ Avoid in pregnancy / lactation
- ▶ Investigational; not FDA-approved