

# Botox Dosing by Zone

Standard unit ranges, injection points, depth, and key precautions by treatment area

UPPER FACE — STANDARD DOSING

| Zone                 | Units (Female) | Units (Male) | Points      | Depth                             | Key Precaution                                                                                               |
|----------------------|----------------|--------------|-------------|-----------------------------------|--------------------------------------------------------------------------------------------------------------|
| Forehead (frontalis) | 10–15u         | 15–20u       | 4–8         | Intramuscular mid-frontalis       | Stay ≥3cm above orbital rim medially. Never treat lateral without treating brow depressors — Spock brow risk |
| Glabella (11s)       | 20–25u         | 25–30u       | 5           | Deep intramuscular — corrugator   | Palpate corrugator during frown. Always include procerus point. Risk: ptosis if too inferior                 |
| Procerus             | 4–6u           | 5–8u         | 1           | Superficial intramuscular midline | Part of glabella complex. Single midline point between brows                                                 |
| Brow lift            | 2–4u per side  | Not typical  | 1–2 lateral | Superficial subcut — lateral brow | Target lateral orbicularis depressor. Never inject frontalis to lift brow                                    |

PERIORBITAL — STANDARD DOSING

| Zone                     | Units (Female) | Units (Male)   | Points | Depth                               | Key Precaution                                                     |
|--------------------------|----------------|----------------|--------|-------------------------------------|--------------------------------------------------------------------|
| Crow's feet              | 6–12u/side     | 10–15u/side    | 3–4    | Superficial subcut                  | Stay ≥1cm from orbital rim. Never treat pretarsal — ectropion risk |
| Lower lid (infraorbital) | 2–4u/side      | Rarely treated | 2      | Very superficial — intradermal      | Advanced technique only. Risk: ectropion, swelling, cheek descent  |
| Bunny lines              | 2–4u/side      | 3–5u/side      | 2      | Superficial intramuscular — nasalis | Stay lateral on nasal wall. Avoid angular artery zone inferiorly   |

| LOWER FACE – STANDARD DOSING |                |              |                  |                                     |                                                                                        |
|------------------------------|----------------|--------------|------------------|-------------------------------------|----------------------------------------------------------------------------------------|
| Zone                         | Units (Female) | Units (Male) | Points           | Depth                               | Key Precaution                                                                         |
| Lip flip (upper)             | 4–6u           | Rarely done  | 4–6 along border | Superficial orbicularis             | Conservative — over-treatment: speech difficulty, can't use straw                      |
| DAO (corners)                | 3–5u/side      | 4–6u/side    | 1–2 per side     | Intramuscular at mandibular border  | Inject AT mandibular border — not modiolus. Risk: smile asymmetry if zygomaticus hit   |
| Mentalis                     | 4–8u           | 6–10u        | 1–2 midline      | Intramuscular chin bulk             | Avoid too lateral — DLI risk causes lower lip droop                                    |
| Masseter (bruxism)           | 20–30u/side    | 25–40u/side  | 3–5              | Intramuscular — have patient clench | Avoid parotid (anterior border). Avoid superior (zygomaticus). Full effect at 3 months |
| Depressor labii              | 2–4u/side      | 3–5u/side    | 1 per side       | Intramuscular                       | Lower lip asymmetry correction. Careful — impairs lip closure                          |

| BOTOX CONVERSION GUIDE   |                           |                          |                                                                       |
|--------------------------|---------------------------|--------------------------|-----------------------------------------------------------------------|
| Product                  | Conversion Ratio to Botox | Reconstitution           | Notes                                                                 |
| Botox (onabotulinumA)    | 1:1 baseline              | 100u vial → 2–4mL saline | Standard reference product                                            |
| Dysport (abobotulinumA)  | 2.5:1 (some use 3:1)      | 300u vial — check label  | Faster onset 2–5 days. Slightly wider diffusion                       |
| Xeomin (incobotulinumA)  | 1:1 with Botox            | 100u vial                | No accessory proteins. Room temp stable. Good for antibody resistance |
| Jeuveau (prabotulinumA)  | 1:1 with Botox            | 100u vial                | Similar efficacy to Botox. Strong rewards program                     |
| Daxxify (daxibotulinumA) | Similar to Botox          | Per package insert       | 6–9 month duration. Different pricing model                           |

| NECK & SPECIALTY – STANDARD DOSING |              |                  |                                                                                |
|------------------------------------|--------------|------------------|--------------------------------------------------------------------------------|
| Zone                               | Units        | Points           | Key Notes                                                                      |
| Platysmal bands (Nefertiti)        | 25–50u total | 4–6 per band     | Pinch band, inject intramuscularly. NEVER deep — carotid, jugular underneath   |
| Hyperhidrosis — axilla             | 50u/side     | 10–15 per axilla | Intradermal grid pattern 1–2cm apart. Works 6–12 months                        |
| Hyperhidrosis — palms              | 100u/side    | Grid pattern     | Painful — requires nerve block or ice. Effects 4–6 months                      |
| Gummy smile                        | 2–4u/side    | 2 per side       | LLSAN at nasolabial junction. Start conservative — 2u per side first treatment |
| Nasal tip droop                    | 2–4u total   | 1–2 midline      | Depressor septi nasi at columellar base. Lifts tip during animation            |

| KEY RULES – NEVER FORGET |                                                                                                                          |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------|
| ★                        | Never inject forehead within 2–3cm of orbital rim medially or 1.5cm laterally — ptosis risk                              |
| ★                        | Always include procerus when treating glabella — incomplete treatment causes persistent horizontal lines                 |
| ★                        | DAO must be injected at mandibular border — NOT at the modiolus — smile asymmetry is the #1 DAO complication             |
| ★                        | Masseter: have patient clench to identify muscle bulk. Never inject anterior to the masseter's anterior border — parotid |
| ★                        | Assess at 14 days. Never judge Botox results before 14 days — results not complete until then                            |

