

Treatment Room Setup Checklist

Complete room setup by service type — injectables, esthetic, IV therapy, and laser

INJECTABLE TREATMENT ROOM — STANDARD SETUP

Patient chair/table	Adjustable treatment chair or table. Patient positioned appropriately for zone being treated (reclined for forehead, seated for lower face).
Lighting	Bright, even, overhead lighting + magnification lamp or ring light for assessment. Natural-looking lighting for post-treatment review.
Sterile field	Sterile drape on instrument tray. All supplies within reach before patient enters room. Never reach over patient mid-treatment.
Injectable supplies	Product (stored appropriately — Botox refrigerated), syringes (27–32g for toxin, 25–30g for filler), alcohol wipes, gauze, topical numbing, cannulas if using.
Emergency supplies	Hyaluronidase IN THE ROOM (not a cabinet, not down the hall). Epinephrine auto-injector if allergic reaction possible. Emergency protocol posted visibly.
Camera	Clean iPhone or camera. Ring light. Photo backdrop or designated photo spot. Photo consent confirmed before any photos taken.
Documentation	Tablet or laptop for charting. Pre-printed consent forms if not digital. Patient chart pulled up before patient enters.

ESTHETIC TREATMENT ROOM — STANDARD SETUP

Table setup	Fresh linens for every patient. Head rest with clean cover. Blanket available. Patient comfort is a reflection of your practice.
Steam/warmth	Facial steamer (if using) turned on and warmed 10 minutes before patient arrives. Warm towels if no steamer.
Products staged	All products for the specific treatment laid out in order of use before patient arrives. Never "hunting" for products mid-treatment.
Tools sanitized	All reusable tools (extractors, brushes, spoons) sanitized and in covered clean container. Disposable items in sealed packaging.
Microneedling specific	Microneedling device charged and set to correct depth for today's protocol. Topical anesthetic out. Serums for post-needling ready.
Post-care supplies	Cool compress ready for post-treatment. Patient-specific post-care printed and ready at end of treatment.

IV THERAPY ROOM – STANDARD SETUP

IV stand	IV pole with bag hooks. Drip chamber and IV line primed with saline before patient enters. No air in the line.
IV supplies	18–22g angiocatheter (based on patient and vein), tourniquet, alcohol wipes, tegaderm, gauze, IV tape, flush syringe (10mL normal saline).
IV bag / medications	Confirm product, concentration, and patient against order before preparation. Label bag with patient name, date, contents, time prepared.
Monitoring supplies	Blood pressure cuff. Pulse oximeter. These should be checked pre-infusion and post-infusion for all patients.
Emergency supplies	Epinephrine auto-injector in the room. Benadryl IM available. Emergency protocol posted. Staff trained in anaphylaxis response.
Patient comfort	Comfortable reclining chair. Warm blanket. Entertainment (TV, tablet). IV therapy takes 1–4 hours — comfort is non-negotiable.

UNIVERSAL ROOM STANDARDS – EVERY ROOM EVERY DAY

- ✓ Surfaces wiped with hospital-grade disinfectant between every patient — treatment table, chair, equipment touchpoints
- ✓ Fresh linens / covers for every patient — never reuse. Cross-contamination risk.
- ✓ Sharps container within arm's reach during any procedure involving needles. Never recap needles.
- ✓ No food or drink in treatment rooms — infection control + professionalism
- ✓ Check expiration dates on all products and medications at the start of each week. Expired products are a liability.
- ✓ Document any adverse reaction or incident in the patient chart the same day. Never wait.

LASER / IPL ROOM – STANDARD SETUP

Protective eyewear	BOTH patient and provider eyewear BEFORE powering on the device. Wavelength-specific eyewear — confirm it matches the device being used.
Device calibration	Calibrate device per manufacturer protocol before first patient of the day. Check energy output, contact cooling, and spot size.
Skin prep	Clean skin — no lotions, sunscreen, or makeup in treatment area. Shave hair removal areas day before (not day of). Confirm Fitzpatrick type.
Test patch	For new patients or sensitive areas — test patch 24 hours before or same day (per protocol). Document settings and response.
Cooling	Contact cooling (built into device), cold gel, or chilled air cooling. Cooling reduces pain and protects epidermis — especially in darker skin.
Post-laser supplies	Cool gel or cold compress ready immediately post-treatment. Aloe vera or calming serum. SPF 50+ for patient to apply before leaving.