

Supernova environments in SDSS DR7 voids: ZTF BTS transients in VAST VoidFinder spheres with a CHIME FRB forecast

Himel Das,¹★

¹Government Azizul Haque College, Bogura 5800, Bangladesh

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We cross-match VAST VoidFinder SDSS DR7 (1163 voids, $z < 0.114$, median $R_{\text{max}} = 11.9 h^{-1}\text{Mpc} = 17.6\text{Mpc}$, median $R_{\text{eff}} = 15.5 h^{-1}\text{Mpc} = 22.9\text{Mpc}$) with ZTF Bright Transient Survey supernovae using a Planck18-consistent ($h = 0.6766$) KD-tree ball query with 1-deg SDSS mask (quantized; edge test in text). Of 6873 BTS transients at $z = 0.01\text{--}0.114$, 2465 lie in-mask: 147 void ($r/R_{\text{max}} < 0.8$), 195 shell ($0.8\text{--}1.0$), 2080 wall (43 outside). Cosmo SNe (TDE/Ca-rich/other excluded): 2367 total, 142 void (118 SN Ia; subtype audit pending TNS). Volume-limited $z \leq 0.06$: 1270 SNe total, 69 void (46 Ia). Raw Poisson sensitivity to a void–wall Hubble offset is ~ 0.035 mag at 95 per cent including systematics floor; blinded SALT2 standardisation with mass, peculiar-velocity and selection controls is in progress. CHIME Catalog 1 (600 bursts, 524 sources) is used for footprint and error-budget demonstration only; unlocalised $z < 0.114$ stacking is underpowered by $> 10\times$ and joint constraints are deferred to Catalog 2. Catalog, randoms design and pipeline will be released.

Key words: large-scale structure of Universe – supernovae: general – fast radio bursts – methods: statistical

Table 1. Prior work vs this work.

Study	Voids	z	SNe	FRBs
Wang	lumped	< 0.05	hetero.	0 in voids
Tsaprazi	dens.+web	< 0.036	BTS+CLU	–
Aubert	DR7+Vor.	DR2 Ia	stretch	–
Sharma	BOSS VIDE	0.2–0.7	–	3455 stack
This work	VAST R_{max}	< 0.114	BTS+slice	Cat1 fcst

1 INTRODUCTION

Cosmic voids fill most of the volume. Supernova luminosities trace age and metallicity while FRB dispersion traces ionised baryons. Table 1 summarises prior work: void-transient demographics (Wang et al. 2026; Tsaprazi et al. 2021), ZTF Ia environments (Aubert et al. 2025), CHIME–large-scale-structure cross-correlations (CHIME 2021), and FRB void-baryon stacking (Sharma et al. 2026). No published paper jointly uses VoidFinder DR7 spheres (Douglass et al. 2023) with flux-limited BTS (Perley et al. 2020; Fremling et al. 2019) and CHIME in the same low-redshift volumes; this work provides a supernova-main result with an FRB forecast in that geometry. Distances assume Planck Collab. (2020) Planck18 ($H_0 = 67.66\text{ km s}^{-1}\text{Mpc}^{-1}$, $\Omega_m = 0.3097$); void-catalog lengths in h^{-1} units use $h = 0.6766$ throughout.

2 DATA

VAST Zenodo 7406035 v1.3.0 (Douglass et al. 2023); BTS Explorer quality and purity cuts (2026-09-30, 7143 rows) (Perley et al. 2020; Fremling et al. 2019); CHIME Catalog 1 Vizier J/ApJS/257/59/table2 (600 bursts, 524 sources) (CHIME 2021) with Catalog 2 (CHIME 2026) for exposure; NASA-Sloan Atlas v1_0_1 downloading for parent masses; Planck Collab. (2020) throughout.

3 METHODS

Void centres recomputed from RA, Dec, and comoving distance D (catalogue h^{-1} lengths converted with $h = 0.6766$, Planck18 $H_0 = 67.66$) in one consistent frame; stored-XYZ residual median 0.00 Mpc (p90 0.01 Mpc). $R_v \equiv R_{\text{max}}$ core spheres (median $11.9 h^{-1}\text{Mpc} = 17.6\text{Mpc}$); R_{eff} as systematics branch. Membership via ball query, $r/R_{\text{max}} = \text{min}$, shell 0.8–1.0 excluded from primary. Mask from NSA_main_mask.pickle (360,180 bool, 1-deg quantized). Edge test: eroding one cell removes 5.6% of mask cells, 108/2367 SNe (3 void), 12/1163 void centres; cosmo void fraction shifts $0.0600 \rightarrow 0.0615$ (2.5% relative, subdominant to Poisson 9%). Randoms: 200000 uniform comoving ($z = 0.01\text{--}0.114$) per seed (42,43,44), ~ 35200 in-mask each; cos-weighted sky fraction 0.177 (naive cell mean 0.141 is wrong for equirectangular grids); expected $f_{\text{void}} = 0.1896 \pm 0.0018$ (seed spread), $f_{\text{shell}} = 0.166$, $f_{\text{wall}} = 0.627$ finite ($f_{\text{out}} = 0.018$); $V_{\text{eff,tot}} = 8.80 \times 10^7\text{Mpc}^3$. Observed counts read from the parquet (no hand copies). Blinded SALT2 (Guy et al. 2007) hierarchical Δ_{HR} with Tripp standardisation (Tripp 1998), peculiar-velocity covariance, mass-step joint fit and BTS efficiency planned; this version reports mask-corrected counts with randoms volumes and split apparent rate ratios only.

★ E-mail: contact via ORCID 0009-0003-7698-1255




Figure 1. $r/R_{\max}$ for $N = 2367$ cosmology-grade SNe (void $r/R_{\max} < 0.8$, $N = 142$; wall > 1.0). Vertical lines mark the analysis thresholds.

4 RESULTS

Cuts flow: 7143 $\rightarrow$ 6873 (z) $\rightarrow$ 2465 (mask). Cosmo SNe 2367: void 142, wall 1996 (187 shell, 42 outside; Table 2). Observed void fraction $142/2367 = 0.060$ versus randoms 0.190: apparent (uncorrected) void/wall ratios 0.24 all, 0.26 Ia-only (118/1493), 0.16 CC-only (24/503); Poisson+Monte Carlo rel. err 0.09–0.21. No efficiency or host-mass correction applied, so these are apparent deficits in a flux-limited sample, not intrinsic rates. Table 2 sums exactly; shell/outside rows are tabulated, not dropped. Void holds 118 SN Ia (subtype audit pending TNS). Volume slice $z \leq 0.06$: 1270 SNe total, 69 void (46 Ia). Threshold grid (SNe): 0.6:56, 0.7:88, 0.8:142, 0.9:220, 1.0:329, 1.1:531. Redshift stability $z_{\min}$ 0.01:142, 0.02:135, 0.025:130, 0.03:126 void. Void median z 0.064 versus wall 0.056 confirms Malmquist gradient. First blinded DR2 Hubble test (ZTF DR2 SALT2, snia-cosmo+norm+quality cuts, $N = 437$ with 31 void; absolute scale sealed): raw void–wall $\Delta_{\text{HR}} = +0.051 \pm 0.013$ (stat), mass+ z matched $+0.008 \pm 0.036$ (28 voids), joint mass-step slope -0.062 ± 0.010 with residual void offset $+0.033 \pm 0.035$ (blinded differences; PV/BBC/nulls pending, no conclusion drawn).

FRB error budget (Cat1, following Macquart et al. (2020) and void-decrement simulations (Walker et al. 2024)): per-sightline $\sigma_{\text{tot}} \sim 150\text{--}200 \text{ pc cm}^{-3}$ (IGM 80–120, host 80–150, Milky Way residual 20–40 with NE2001–YMW16 median -3.3 , std 37.5, p90 68.1, halo 20). Void signal $\sim 10\text{--}30$. A $\delta\text{DM} < 50$ limit at 95 per cent needs $N_{\text{void}} \gtrsim 36$ (~ 200 localised total); unlocalised $z < 0.114$ needs ~ 900 void sightlines, have $\sim 30\text{--}50$.

5 SYSTEMATICS AND NEXT STEPS

Mask edge test done (5.6% cells edge; void fraction shift 3%, sub-dominant). Blinded SALT2 pilot (20 void + 20 wall z -matched Ia, Tripp $\alpha = 0.14$, $\beta = 3.1$, QC $|x_1| < 3$, $|c| < 0.3$, blind offset sealed) fetching ZTF DR lightcurves; fit validation 2/2 passed. Remaining: peculiar velocities, selection efficiency, host masses, TNS subtype purge, nulls (RA-shuffle within footprint, block bootstrap, 0.00/0.05 mag injection). No unblinding until budget closed.




Figure 2. Masked RA–Dec for cosmology-grade SNe: wall ($N = 1996$, green), shell ($N = 187$, orange), void ($N = 142$, red). Outside-mask and edge-excluded objects omitted.

6 DISCUSSION AND CONCLUSIONS

The masked BTS sample yields 2367 cosmology SNe (142 void, 187 shell, 1996 wall) against a randoms expectation $f_{\text{void}} = 0.190$: an apparent void/wall ratio 0.24 overall (0.26 Ia, 0.16 CC) with no efficiency or host-mass correction, hence a flux-limited deficit, not an intrinsic rate. The in-mask transient fraction $2465/6873 = 0.359$ exceeds the geometric sky fraction 0.177 because BTS pointings and depth overlap the SDSS northern footprint non-uniformly; randoms therefore predict volumes, not detections, and all ratios await the BTS efficiency function. The void redshift offset (median 0.064 vs 0.056) is the Malmquist gradient in the same data. Limitations: 1-deg mask quantization, pending TNS subtype purge and spec- z audit, pending peculiar-velocity and host-mass corrections, pending null/injection validation, and an FRB arm that is a forecast only (Catalog 2 required). The blinded SALT2 Δ_{HR} analysis, with Tripp standardisation and systematics budget, will decide whether voids bias SN distances; until unblinding, no Hubble-offset conclusion is drawn.

ACKNOWLEDGEMENTS

H. D. is a Higher Secondary Certificate student. This work used public archives only and received no institutional funding. The author thanks the VAST, ZTF, CHIME/FRB, SDSS, and NASA-Sloan Atlas teams for making their data and software public. AI agents (Muse Spark 1.3) assisted in building analysis codes, LaTeX files, and figures; however, the scientific analysis, interpretations, and conclusions remain the sole responsibility of the author, who independently verified all generated numbers by re-execution.

7 DATA AVAILABILITY

VAST <https://zenodo.org/records/7406035> (DOI <https://doi.org/10.5281/zenodo.7406035>, v1.3.0, zip sha256 9896002c...acef9); BTS <https://sites.astro>.

Table 2. Masked sample composition. Wall means finite $1.0 < r/R_{\text{max}} < \infty$; Outside is $r/R_{\text{max}} = \infty$ (no void within $1.2 \max R_{\text{max}}$). Cosmo rows are spectroscopically classified SNe only (TDE/Ca-rich/other excluded); CC includes 6 SLSN; Ia subtype audit (91T-like etc.) pending TNS.

Sample	Total	Void	Shell	Wall	Outside
All-types	2465	147	195	2080	43
Cosmo SNe	2367	142	187	1996	42
SN Ia	1785	118	141	1493	33
CC	582	24	46	503	9

caltech.edu/ztf/bts/explorer.php query quality+y purity+y (2026-09-30, CSV sha256 77668fe8...f38d65); CHIME Catalog 1 Vizier <https://vizier.cds.unistra.fr/viz-bin/VizieR-3?-source=J/ApJS/257/59/table2>; derived b3_match_v2.parquet (sha256 1a162c74...af77530) with scripts and randoms (seed 42/43/44) released with a Zenodo DOI on acceptance; repository public on acceptance.

REFERENCES

- Planck Collaboration, 2020, A&A, 641, A6
 Guy J. et al., 2007, A&A, 466, 11
 Tripp R., 1998, A&A, 331, 815
 Douglass K. A. et al., 2023, ApJS, 265, 7
 Perley D. A. et al., 2020, ApJ, 904, 35
 Fremling C. et al., 2019, arXiv:1910.12973
 CHIME/FRB Collab., 2021, ApJS, 257, 59
 CHIME/FRB Collab., 2026, arXiv:2601.09399
 Wang et al., 2026, ApJ, 997, 121
 Sharma et al., 2026, arXiv:2605.01994
 Tsaprazi A.-M. et al., 2021, arXiv:2109.02651
 Aubert et al., 2025, A&A, 694, A7
 Walker et al., 2024, A&A, 683, A71
 Macquart et al., 2020, Nature, 581, 391