

# 1 General info

## 1.1 Sum Square Error Formula(SSE)

$$E(\theta) = \sum_i^n (\hat{y}_i - y_i)^2$$

Since the formula for the predicted value is

$$\hat{y} = \theta_1 * x_i + \theta_2$$

the SSE formula can be summarized into

$$E(\theta) = \sum_i^n (\theta_1 * x_i + \theta_2 - y_i)^2$$

## 1.2 Canonical machine learning optimization problem

?? Formula not understandable. Lecture 3 deck, slide 10

## 1.3 vocabulary

- 1.Hypothesis function: The function used to predict values
- 2.Objective function: The function is used to identify the difference between the predicted values from the hypothesis function and the real values
- 3.Optimization problem: Is used to optimize the hypothesis parameters to reduce the objective function

### 1.3.1 Signs

- 1.Input features:  $x^i$
- 2.Target features:  $y^i$
- 3.Model parameters:  $\theta$
- 4.Hypothesis function:  $h_\theta = h_\theta(x) = \sum_{j=1}^n \theta_j * x_j$
- 5.Objective function:  $\ell$

## 1.4 Regression with canonical formulation and matrix

- 1.Hypothesis function  
 $h_\theta(x) = \sum_{j=1}^n \theta_j * x_j$   
 $\theta_j$  is the intercept for  $x = 0$ , for  $x \neq 0$  it is the slope
- 2.Objective function  
 $\arg \min_\theta$ : Means that  $\theta$  (the parameter vector) should be optimized regarding min error  $\theta = l(\hat{y}, y) = 1/2(\hat{y} - y)^2$
- 3.Optimization problem  
 $\theta$ : Vector of parameters that need to be optimized (Slope)  
 $n$ : number of features (Of each instance)  
 $m$ : number of instances (Number of data points in the dataset)

$$\theta = \arg \min_{\theta} \sum_{i=1}^m \left( \sum_{j=1}^n \theta_j x_j^i - y^i \right)^2$$

### 1.4.1 Optimization problem formula explained

$\theta = \arg \min_{\theta}$ : The to be optimized value is all parameter vector  $\theta$ , so the Slope, to minimize errors

$\sum_{i=1}^m$ : The sum of errors for all data points in the dataset

$\theta_j x_j^i$ :  $\theta$  is the slope for a specific feature  $x_j$  at position j at the i'th instance

$\sum_{j=1}^n \theta_j x_j^i$ : Sums up the slope j multiplied with the feature j for the data point i, so  $\hat{y}^i$  is calculated

$(\sum_{j=1}^n \theta_j x_j^i - y^i)^2$ : Calculates the squared difference between the predicted value  $\hat{y}^i = \sum_{j=1}^n \theta_j x_j^i$  and the actual value  $y^i$  for the data point i

## 2 Minimizing Loss function

$$l(\theta) = \sum_{i=1}^n (\theta_j x_j^i - y^i)^2$$

## 3 To repeat

Deck 3 Slide 15

## 4 Gradient descent

Negative Derivated error function and step for step insert x-values to find the lowest y-value

The stepsize indicates the difference between the current and the next entered x-value. If its to high, the lowest point might be skipepd.

The optimal  $\theta$  is to be found by this formula:

$$\theta_j := \theta_j - \alpha \sum_{i=1}^m (\theta^\top x^{(i)} - y^{(i)}) x_j^{(i)} ==$$

$$\theta = (X^T X)^{-1} X^T y$$

## 5 Bias and Variance

Bias: Describes how the model performs on training data. Low Bias = Good fit for training data

Variance: Describes how the model performs on test data. High variance = Bad fit for new data (Overfitting)

## 6 Underfit and overfit

Underfit: Low variance and high bias

Overfit: High variance and low bias With increasing model complexity, the training loss decreases. The generalization/prediction starts to increase because of overfit

## 7 Absolute Regression Test Metrics

Signs:

e: Error

n: Amount of data points

### 7.1 Mean Square Error/Deviation

$$\frac{1}{n} \sum_{i=1}^n e_i^2$$

Is the average of the squares of the differences between the actual values and the predicted values. Use: Larger errors are penalized more heavily. Therefore it is good for an overview, but might be skewed by outliers. Lower errors are better

### 7.2 Root Mean Square Error

Term

$$\sqrt{\frac{1}{n} \sum_{i=1}^n e_i^2}$$

Is the root of the mean square error, but because of the root, the units stay the same

### 7.3 Mean Absolute Error/Deviation

Term

$$\frac{1}{n} \sum_{i=1}^n |e_i|$$

It is more robust to outliers and gives a straightforward interpretation of the average error magnitude.

### 7.4 Average Error

Term

$$\frac{1}{n} \sum_{i=1}^n e_i$$

Indicates whether the predictions are on average over- or underpredicting the target response

## 8 Relative Regression Test Metrics

Are used, measure error or performance in ratios or percent. More usefull when comparing datasets with different units and scales.

### 8.1 r-Squared

Returns a value between  $r^2=0$  and  $r^2=1$

1: The model has a perfect fit

0: The model has no fit

$r^2 < 0$ : Worse then a horizontal line

## 9 Model training cycle

1.Divide data into training, validation, test set

2.Train model on training set

3.Use model on validation data to see performance and adjust hyperparameters (polynomial degree)

4.Retrain system on training and validation dataset

5.Evaluate performance on test set

### 9.1 Test data leakage

Test data leakage describes the usage of test data to adjust the model (a.e. adjusting hyperparameters). Therefore the test data set should be isolated.

#### 9.1.1 Solving test data leakage

1: Recollect data if overfitting to test set is suspected

2: Dont look at test set

## 10 Regularization

The degree of polynomial can be seen as complexity of the model

Regularization is used to prevent overfitting by penalizing large coefficient values.

## **10.1 L2 Regularization/Ridge regression**

## **11 PROBLEMS**

Lecion 4 slide 7 R Squared Leciton 4 slide 24