

Modeling of weak polyelectrolyte hydrogels under compression.

Implications for water desalination.

Oleg V. Rud

June 21, 2021

Department of Physical and Macromolecular Chemistry, Faculty of Science, Charles University in Prague, Hlavova 8, Praha 2 128 00, Czech Republic

Intro

Introduction

figures/step2.png

- Hydrogels for desalination
- Forward osmosis
- Various stimuli: thermo-, pH-, electric-, magnetic-, light-induced gel collapse
- Manfred Wilhelm and Yu Chi experiment

Fengler, C., Arens, L., Horn, H., Wilhelm, M. (2020). **Desalination of Seawater Using Cationic Poly(acrylamide) Hydrogels and Mechanical Forces for Separation.** Macromolecular Materials and Engineering

Yu, C., Wang, Y., Lang, X., Fan, S. (2016). **A Method for Seawater Desalination via Squeezing**

The model of a polyelectrolyte gel.

Langevin dynamics.




Figure 1: The snapshot of the hydrogel model for Langevin dynamics

- Diamond network of point particles
- Lennard–Jones interaction

$$V_{LJ}(r) = \begin{cases} 4\epsilon \left(\left(\frac{\sigma}{r-r_c} \right)^{12} - \left(\frac{\sigma}{r-r_c} \right)^6 \right) & , r < r_c \\ 0 & , r > r_c \end{cases}$$

- FENE potential

$$V_{FENE}(r) = -\frac{1}{2}\Theta\Delta r_{max}^2 \ln \left[1 - \left(\frac{r-r_0}{\Delta r_{max}} \right)^2 \right]$$

- Electrostatic interaction

$$V_{EL} = l_B k_B T \cdot \frac{q_1 q_2}{r}$$


Grand-reaction ensemble.




$$K = \mu_{\text{H}^+}^{\ominus} + \mu_{\text{A}^-}^{\ominus} - \mu_{\text{HA}}^{\ominus}$$



$$K = \mu_{\text{Na}^+} + \mu_{\text{Cl}^-}$$



$$K = 2\mu_{\text{Ca}^{2+}} + \mu_{\text{Cl}^-}$$



$$\Delta\Omega = k_B T \ln \left(K^\xi \prod_i V^{\nu_i \xi} \frac{N_i!}{(N_i + \nu_i \xi)!} \right) + \Delta E$$

Simulation protocol.

1. Choose randomly: LMD or RE.
2. Simulate the chosen, collecting 50 samples of:

LMD: pressure, P ,
and $\{R_e\}$

RE: number of ion-
ized segments, N_{A-}


RE: number of salt ions,
 N_{Na^+} , N_{Cl^-} , $N_{Ca^{2+}}$

Check the autocorrelation of each samples array.
Pearson coefficient must be < 0.2 .

3. Repeat collecting at least 100 averages from each process.

Results

Seawater model solution



$$c_{\text{Cl}^-} = 0.54 \text{ mol/l}$$

of negative ions

$$c_{\text{Na}^+} = 0.47 \text{ mol/l}$$

of positive ions

$$c_{\text{Ca}^{2+}} = 0.063 \text{ mol/l}$$

of positive divalent ions

$$c_{\text{Na}^+} \simeq 0.87 \cdot c_{\text{Cl}^-}$$
$$c_{\text{Ca}^{2+}} \simeq 0.117 \cdot c_{\text{Cl}^-}$$

$$\mu_{\text{Na}^+} = \mu_{\text{Cl}^-} - 0.139kT$$
$$\mu_{\text{Ca}^{2+}} = \mu_{\text{Cl}^-} - 2.03kT$$

Compression of hydrogel.






Figure 2: Compression of hydrogel in equilibrium with the bath of salinity $c_s = c_{\text{Cl}^-} = 0.007 \text{ mol/l}$

Desalination effect.



- The gel is compressed from the initial volume, V_0 to a volume V^{gel} .
- And exchanges small ions with a reservoir solution.

Desalination and/or ion exchange.



Mean field model of hydrophobic weak polyelectrolyte gel.

- The gel is **regular network**.
big enough to treat it is
homogeneous medium
- Affine deformation
approximation $R \sim V^{1/3}$
- weak hydrophobic polyacid,
 $pK = 6$, $\chi = 2.0$

Ionization equilibrium

figures/grand_reaction_model_noCa.png




Rud, O., Borisov, O., Kosovan, P. (2018).

Thermodynamic model for a reversible desalination cycle


using weak polyelectrolyte hydrogels. Desalination, 442, 32–43.

The desalination cycle.



(a) step 1

(b) step 2



The desalination cycle.



figures/cycle_PV_chi0.pdf

(a) $\chi = 0$



figures/cycle_PV_chi2.pdf

(b) $\chi = 2$

Questions?


Monovalent salt. “No electrostatics” vs “Mean field theory”.

$$\frac{\alpha}{1-\alpha} 10^{\text{pK}-\text{pH}} = \sqrt{1 + \left(\frac{\alpha c_p}{2c_s} \right)^2} - \frac{\alpha c_p}{2c_s}$$


Together with electroneutrality condition it translates to

$$-\frac{\alpha^3 c_p}{c_s} + \alpha^2 \left(\frac{c_p}{c_s} + \Theta - \frac{1}{\Theta} \right) + \frac{2\alpha}{\Theta} - \frac{1}{\Theta} = 0$$

where $\Theta = 10^{\text{pK}-\text{pH}}$.



(e) low salinity, $c_s = 0.007$ mol/l



(f) high salinity, $c_s = 0.209$ mol/l