

Standard Treatment Guidelines: Hypertension Quick Reference Guide

Ministry of Health and Family Welfare (MoHFW) | National Health Mission India

Section 1: Primary Health Centre Hypertension Management

Screen adult patients for hypertension during primary clinic visits. Persistent blood pressure $\geq 140/90$ mmHg requires lifestyle modification (dietary sodium restriction $< 2\text{g/day}$, regular aerobic physical activity for 30 minutes daily) and structured clinical follow-up. First-line pharmacological management for essential hypertension includes Amlodipine 5mg once daily or Telmisartan 40mg once daily under clinical supervision.

Section 2: Red-Flag Emergency Triage Referral Criteria

Healthcare workers and AI triage systems must identify predefined red-flag symptoms for immediate emergency transfer: 1. Severe chest pain radiating to jaw or left arm (suspected acute coronary syndrome). 2. Sudden onset of severe dyspnea (shortness of breath) or acute respiratory failure. 3. Altered mental status, confusion, focal neurological deficit, or loss of consciousness. 4. Severe abdominal rigidity or uncontrollable acute hemorrhage. 5. Persistent severe hypertension BP $\geq$ 180/110 mmHg with acute target organ damage. When red-flag symptoms are identified, immediately issue an emergency alert and arrange transfer to the nearest Community Health Centre (CHC) or District Hospital.