



CS6109 – GRAPH THEORY

Module – 2

Presented By


Dr. S. Muthurajkumar,
Assistant Professor,
Dept. of CT, MIT Campus,
Anna University, Chennai.

Module - 2

- Edge Graphs and Traversability
- Eccentricity Sequences and Sets
- Isometry


Edge Graphs

- Definition: Let $G(V, E)$ be a graph with $V = \{v_1, v_2, \dots, v_n\}$ and $E = \{e_1, e_2, \dots, e_m\}$.
- The edge graph $L(G)$ of G has the vertex set E and two vertices e_i and e_j are adjacent in $L(G)$ if and only if the corresponding edges e_i and e_j of G are adjacent in G .
- $L(G)$ is the edge graph of G . A graph G is an edge graph if it is isomorphic to the edge graph $L(H)$ of some graph H .



Observations Edge Graphs

1. A graph G is connected if and only if $L(G)$ is connected.
2. If H is a subgraph of G , then $L(H)$ is a subgraph of $L(G)$.
3. The edges incident at a vertex of G form a maximal complete subgraph of $L(G)$.
4. In G , if $e = uv$ is an edge, then the degree of e in $L(G)$ is the number of edges of G adjacent to e in G . Clearly, $d_{L(G)}(e) = d_G(u) + d_G(v) - 2$.
5. For $n > 1$, $L^n(G) = L(L^{n-1}(G))$ and $L^0(G) = G$.



Theorem 9.1 The number of edges m' in $L(G)$ when G has degree sequence $[d_i]_1^n$ is given by

$$m' = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m.$$

Proof Let $[d_i]_1^n$ be the degree sequence of the graph G and let $L(G)$ the edge graph of G , have m' edges.

As the degree of the vertex v_i in G is d_i , there are d_i edges incident on v_i . From these d_i edges, any two are adjacent at v_i in G . Hence the number of edges contributed by v_i to $L(G)$ is $\binom{d_i}{2}$.

$$\begin{aligned} \text{Thus, } m' &= \sum_{i=1}^n \binom{d_i}{2} = \sum_{i=1}^n \frac{d_i(d_i - 1)}{2} = \frac{1}{2} \sum_{i=1}^n (d_i^2 - d_i) \\ &= \frac{1}{2} \sum_{i=1}^n d_i^2 - \frac{1}{2} \sum_{i=1}^n d_i \\ &= \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m. \end{aligned}$$


Theorem 9.1 The number of edges m' in $L(G)$ when G has degree sequence $[d_i]_1^n$ is given by

$$m' = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m.$$

Proof:

We know that $m = \frac{1}{2} \left(\sum_{i=1}^n d_i \right)$

Therefore $m' = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - \frac{1}{2} \left(\sum_{i=1}^n d_i \right)$



$$m' = \frac{1}{2}(1^2 + 2^2 + 2^2 + 2^2 + 3^2) - \frac{1}{2}(1 + 2 + 2 + 2 + 3)$$




$$= 11 - 5$$

$$= 6 \text{ (Number of edges in } L(G)\text{).}$$

Theorem 9.2 The edge graph of a graph G is a path if and only if G is a path.

Proof Let G be a graph with n vertices. Assume G is a path P_n . Then $L(G)$ is the path P_{n-1} with $n - 1$ vertices.

Conversely, let $L(G)$ be a path. Then no vertex of G has degree greater than two. For, if G has a vertex v of degree greater than two, the edges incident to v form a complete subgraph of $L(G)$ with at least three vertices. Therefore G is either a cycle or a path. But G cannot be a cycle, since the edge graph of a cycle is a cycle.



Theorem 9.3 A connected graph is isomorphic to its edge graph if and only if it is a cycle.

Proof Let G be a connected graph with n vertices, m edges and with degree sequence $[d_i]_1^n$. Let $L(G)$ be the edge graph of G . The number of vertices in $L(G)$ is m . The number of edges m' in $L(G)$ is given by

$$m' = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m.$$

Clearly, $L(G)$ is connected and $L(C_n) = C_n$.

Conversely, let $G \cong L(G)$.

Then G and $L(G)$ have the same number of vertices and edges.

$$\text{So, } n = m \text{ and } m = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m. \quad \longrightarrow \quad m = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right) - m.$$

$$\text{Therefore, } n = m \text{ and } \sum_{i=1}^n d_i^2 = 4m. \quad \longleftarrow \quad m + m = \frac{1}{2} \left(\sum_{i=1}^n d_i^2 \right)$$

Thus, variance

$$\left(\sum_{i=1}^n d_i^2 \right) = 4m$$

$$\{[d_i]\} = \frac{1}{n} \sum_{i=1}^n d_i^2 - \left(\frac{1}{n} \sum_{i=1}^n d_i \right)^2$$

$$\left[\text{Because } Var = \frac{1}{N} \sum_i f_i x_i^2 - \left(\frac{1}{N} \sum_i f_i x_i \right)^2 \text{ and we have } f_i = 1 \right]$$


$$= \frac{1}{n} 4m - \frac{1}{n^2} (2m)^2 = \frac{4m}{m} - \frac{4m^2}{m^2} = 4 - 4 = 0.$$

$$\left(\sum_{i=1}^n d_i^2 \right) = 4m$$

Therefore the d_i 's are equal and G is regular of degree d , say.

$$\text{So } nd = 2m \text{ implies that } d = \frac{2m}{n} = \frac{2m}{m} = 2.$$

Thus G is a 2-regular connected graph, that is, C_n .



$$\left(\frac{1}{n} \sum_{i=1}^n d_i \right)^2 = \frac{1}{n^2} \left(\sum_{i=1}^n d_i \right)^2$$


$$m = \frac{1}{2} \left(\sum_{i=1}^n d_i \right)$$

$$\left(\sum_{i=1}^n d_i \right) = 2m$$


Edge Graphs and Traversability

Theorem 9.10 If G is Eulerian, then $L(G)$ is both Eulerian and Hamiltonian.


Proof Let G be Eulerian and let $\{e_1, e_2, \dots, e_m\}$ be the edge sequence of an Euler line in G . Let the edge e_i in G be represented by the vertex v_i in $L(G)$, $1 \leq i \leq m$. Then $v_1v_2 \dots v_mv_1$ is a Hamiltonian cycle of $L(G)$. Now, if $e = u_iu_j \in E(G)$ and the vertex v in $L(G)$ represents the edge e , then $d_{L(G)}(v) = d_G(u_i) + d_G(u_j) - 2$, which is obviously even and greater than or equal to two, since both $d_G(u_i)$ and $d_G(u_j)$ are even (and ≥ 2). Thus in $L(G)$ every vertex is of even degree (≥ 2). Hence $L(G)$ is Eulerian.



Edge Graphs and Traversability



G




$L(G)$

$L(G)$ is both Eulerian and Hamiltonian, but G is not Eulerian.

Dominating walk

- A dominating walk of a graph G is a closed walk W in G (which can be just a single vertex) such that every edge of G not in W is incident with W . For example, the walk $v_1 v_2 v_3 v_4$ in the graph is the dominating walk.



Corollary 9.1 The edge graph of a Hamiltonian graph is Hamiltonian.

Proof Let G be a Hamiltonian graph with Hamiltonian cycle C . Then C is a dominating walk of G , and hence, $L(G)$ is Hamiltonian.

We note that the converse of Corollary 9.1 is not true in general. To see this, consider the graph G as shown in Figure 9.16. Clearly $L(G)$ is Hamiltonian but G is not.




Fig. 9.16

Eccentricity Sequences and Sets

Eccentricity of a vertex: Let G be a connected graph. The eccentricity of a vertex v in G is the distance of the vertex u farthest from v . It is denoted by $e(v)$. That is, $e(v) = \max\{d(u, v) : u \in V\}$.

The minimum eccentricity is called the *radius* of G and the maximum eccentricity is called the *diameter* of G . The radius is denoted by r and diameter by d .

Therefore, $r = \min\{e(v) : v \in V\}$ and $d = \max\{e(v) : v \in V\}$.

Definition: A positive sequence $[e_i]_1^n$ is called an *eccentricity sequence* if it is an eccentricity sequence of some graph. The graph is said to realise the sequence. A set of positive integers is called an *eccentricity set* if it is an eccentricity set of some graph. The graph is said to realise the set. (The set of distinct eccentricities in a graph is called the eccentricity set of that graph.)

Theorem 9.18 If uv is an edge of a connected graph G , then $|e(u) - e(v)| \leq 1$.

Proof Let w be an eccentric vertex of u (i.e., w is the farthest vertex from u). Then by the triangle inequality for the metric d (distance), we have

$$d(u, w) \leq d(u, v) + d(v, w)$$

so that $e(u) \leq d(u, v) + d(v, w)$. (9.18.1)

But u and v are adjacent, therefore $d(u, v) = 1$.

Also, $e(v) \geq d(v, w)$ so that $d(v, w) \leq e(v)$.

Thus, from (9.18.1) we have

$$e(u) \leq 1 + d(v, w) \text{ so that } e(u) \leq 1 + e(v).$$

Therefore, $e(u) - e(v) \leq 1$. (9.18.2)

Similarly, by considering an eccentric vertex of v , we have

$$e(v) - e(u) \leq 1. \tag{9.18.3}$$

From (9.18.2) and (9.18.3) it follows that





$$|e(u) - e(v)| \leq 1.$$

Note The above result shows that the eccentricities of two adjacent vertices are either equal or differ by 1 as $|e(u) - e(v)| \leq 1$ gives $|e(u) - e(v)| = 0$ or $|e(u) - e(v)| = 1$.

Isometry

Let G_1 and G_2 be connected graphs with vertex sets V_1 and V_2 respectively. Then G_2 is said to be *isometric* from G_1 if for each $v \in V_1$, there is a one-one map $\phi_v : V_1 \rightarrow V_2$ such that ϕ_v preserves distances from v , that is $d_{G_2}(u, v) = d_{G_1}(\phi_v(v), \phi_v(u))$ for every $u \in V_1$.

Two graphs G_1 and G_2 are said to be *isometric* if they are isometric from each other.



Theorem 9.23 If G_1 and G_2 are k -regular graphs of order n , where $k \geq n - 1/2$, then G_1 and G_2 are isometric.

Proof Since G_1 is a k -regular graph with $k \geq n - 1/2$, $d(G_1) \leq 2$.
Let $u \in V(G_1)$ and $v \in V(G_2)$ be any two vertices and define

$$\phi_u : V(G_1) \rightarrow V(G_2) \text{ by } \phi_u(u) = v.$$

For $i = 1, 2, \dots, k$, let $u_i \in N_1(u)$ and $v_i \in N_1(v)$ and define $\phi_u(u_i) = v_i$.

For $i = k + 1, \dots, n - 1$, let $u_i \in N_2(u)$ and $v_i \in N_2(v)$ and again let $\phi_u(u_i) = v_i$.


The neighbourhoods are in the appropriate graphs. Then ϕ_u is an isometry of G_2 from G_1 at u . Since u and v are arbitrary, it is easily seen that G_2 is isometric from G_1 , and G_1 is isometric from G_2 .

Theorem 9.24 A necessary condition for two graphs to be isometric is that they have the same degree set and the same eccentricity set.

Proof Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be isometric graphs. As G_2 is isometric from G_1 , let ϕ_v be the one-one mapping from $V(G_1)$ to $V(G_2)$. Therefore, $d(v|G_1) = d(\phi_v(v)|G_2)$. Also ϕ_v has the property of preserving distance, therefore $e(v|G_1) = e(\phi_v(v)|G_2)$. So the eccentricity set of G_1 is included in the eccentricity set of G_2 .

Again, as G_1 is isometric from G_2 , therefore, the degree set and eccentricity set of G_2 are included respectively in the degree set and eccentricity set of G_1 .

Hence the degree sets are equal in G_1 and G_2 and the eccentricity sets are equal in G_1 and G_2 .



REFERENCES:

1. Narsingh Deo, "Graph Theory with Applications to Engineering and Computer Science", Prentice-Hall of India Pvt. Ltd, 2003.
2. S. Pirzada, "An Introduction to Graph theory", University Press, 2012.
3. Frank Harary, "Graph Theory", Narosa Publishing House, 2001.
4. West D. B., "Introduction to Graph Theory", 2nd Edition, Pearson Education, 2001.
5. Diestel R, "Graph Theory", 5th Edition, Springer, 2017.

Thank you.