



# **CS6109 – GRAPH THEORY**

## **Module – 6**

### **Presented By**



Dr. S. Muthurajkumar,  
Assistant Professor,  
Dept. of CT, MIT Campus,  
Anna University, Chennai.

# Module - 6

- Digraph
- Properties
- Euler Digraph
- Tournament graph
- Applications

# Digraphs (Directed graphs)


- A digraph  $D$  is a pair  $(V, A)$ , where  $V$  is a nonempty set whose elements are called the vertices and  $A$  is the subset of the set of ordered pairs of distinct elements of  $V$ .



Directed graph with 5 vertices and 10 edges.

# In-degree/Out-degree

- The number of edges incident out of a vertex  $v_i$  is called the *out-degree* (or *out-valence* or *outward demidegree*) of  $v_i$  and is written  $d^+(v_i)$ .
- The number of edges incident into  $v_i$  is called the *in-degree* (or *invalence* or *inward demidegree*) of  $v_i$  and is written as  $d^-(v_i)$ .




$$\begin{array}{ll} d^+(v_1) = 3, & d^-(v_1) = 1, \\ d^+(v_2) = 1, & d^-(v_2) = 2, \\ d^+(v_5) = 4, & d^-(v_5) = 0. \end{array}$$

Directed graph with 5 vertices and 10 edges.


# Multidigraphs

- A multidigraph  $D$  is a pair  $(V, A)$ , where  $V$  is a nonempty set of vertices and  $A$  is a multiset of arcs, which is a multisubset of the set of ordered pairs of distinct elements of  $V$ .
- The number of times an arc occurs in  $D$  is called its multiplicity and arcs with multiplicity greater than one are called multiple arcs of  $D$ .




# General Digraphs

- A general digraph  $D$  is a pair  $(V, A)$ , where  $V$  is a nonempty set of vertices, and  $A$  is a multiset of arcs, which is a multisubset of the cartesian product of  $V$  with itself.
- An arc of the form  $uu$  is called a loop of  $D$  and arcs which are not loops are called proper arcs of  $D$ .
- The number of times an arc occurs is called its multiplicity. A loop with multiplicity greater than one is called a multiple loop.





# Oriented Graph

- A digraph containing no symmetric pair of arcs is called an oriented graph.



# Isomorphic Digraphs

- Isomorphic graphs were defined such that they have identical behavior in terms of graph properties.
- In other words, if their labels are removed, two isomorphic graphs are indistinguishable.




Two nonisomorphic digraphs.




# Simple Digraphs


- A digraph that has no self-loop or parallel edges is called a simple digraph.




Simple Digraph



Simple Digraph




Simple Digraph




Not Simple Digraph

# Asymmetric Digraphs

- Digraphs that have at most one directed edge between a pair of vertices, but are allowed to have self-loops, are called *asymmetric* or *antisymmetric*.




*Asymmetric Graph*



*Not Asymmetric Graph*

# Symmetric Digraphs


- Digraphs in which for every edge  $(a, b)$  (i.e., from vertex  $a$  to  $b$ ) there is also an edge  $(b, a)$ .




**Symmetric Digraphs**

# Simple Symmetric/Asymmetric Digraphs

- A digraph that is both simple and symmetric is called a *simple symmetric digraph*.
- Similarly, a digraph that is both simple and asymmetric is *simple asymmetric*.




*Simple Asymmetric Digraph*




*Simple Symmetric Digraph*

# Complete Digraphs

- Digraphs in which for every edge  $(a, b)$  (i.e., from vertex  $a$  to  $b$ ) there is also an edge  $(b, a)$ .




Complete symmetric digraph of four vertices.




Complete symmetric digraph

# Complete Symmetric/Asymmetric Digraphs

- A *complete symmetric digraph* is a simple digraph in which there is exactly one edge directed from every vertex to every other vertex.
- A *complete asymmetric digraph* is an asymmetric digraph in which there is exactly one edge between every pair of vertices.




Complete symmetric digraph



Complete asymmetric digraph

# Tournament


- A complete antisymmetric digraph, or a complete oriented graph is called a tournament. Clearly, a tournament is an orientation of  $K_n$ .



Tournament

# Euler Digraphs

- A digraph  $D$  is said to be Eulerian if it contains a closed walk which traverses every arc of  $D$  exactly once. Such a walk is called an Euler walk.
- A digraph  $G$  a closed directed walk (i.e., a directed walk that starts and ends at the same vertex) which traverses every edge of  $G$  exactly once is called a *directed Euler line*. A digraph containing a directed Euler line is called an *Euler digraph*.




The walk  $a b c d e f$  is an Euler line.



**THEOREM 9-1:** A digraph  $G$  is an Euler digraph if and only if  $G$  is connected and is balanced [i.e.,  $d^-(v) = d^+(v)$  for every vertex  $v$  in  $G$ ].


- $d^+(1) = 2,$
- $d^-(1) = 2,$
- $d^+(2) = 2,$
- $d^-(2) = 2,$
- $d^+(3) = 1,$
- $d^-(3) = 1,$
- $d^+(4) = 1.$
- $d^-(4) = 1.$



The walk  $a b c d e f$  is an Euler line.

**THEOREM 9-1:** A digraph  $G$  is an Euler digraph if and only if  $G$  is connected and is balanced [i.e.,  $d^-(v) = d^+(v)$  for every vertex  $v$  in  $G$ ].


- $d^+(r) = 1,$
- $d^-(r) = 1,$
- $d^+(d) = 2,$
- $d^-(d) = 2,$
- $d^+(c) = 2,$
- $d^-(c) = 2,$
- $d^+(a) = 2.$
- $d^-(a) = 2.$
- $d^+(b) = 1.$
- $d^-(b) = 1.$



The walk  $r \rightarrow d \rightarrow a \rightarrow b \rightarrow c \rightarrow d \rightarrow c \rightarrow a \rightarrow r$  is an Euler line.

# Hamiltonian Digraphs


- A connected digraph  $D$  is **Hamiltonian** if  $D$  contains a cycle that itself contains all of the vertices in  $D$ .
- This cycle is known as a **Hamiltonian Cycle**.



Hamiltonian cycle -  $abcda$

# Tournaments

- A tournament is an orientation of a complete graph.
- Therefore in a tournament each pair of distinct vertices  $v_i$  and  $v_j$  is joined by one and only one of the oriented arcs  $(v_i, v_j)$  or  $(v_j, v_i)$ .
- If the arc  $(v_i, v_j)$  is in  $T$ , then we say  $v_i$  dominates  $v_j$  and is denoted by  $v_i \rightarrow v_j$ . The relation of dominance thus defined is a complete, irreflexive, antisymmetric binary relation.



# REFERENCES:

1. Narsingh Deo, "Graph Theory with Applications to Engineering and Computer Science", Prentice-Hall of India Pvt. Ltd, 2003.
2. S. Pirzada, "An Introduction to Graph theory", University Press, 2012.
3. Frank Harary, "Graph Theory", Narosa Publishing House, 2001.
4. West D. B., "Introduction to Graph Theory", 2nd Edition, Pearson Education, 2001.
5. Diestel R, "Graph Theory", 5th Edition, Springer, 2017.

Thank you.