

# DATA SCIENCE WITH R

# REGRESSION ANALYSIS

Overview

Simple Linear Regression

Multiple Linear Regression



**Regression Assumptions**

Implementation in SAS



# Regression Assumptions

1. Model is *linear in parameters*



# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population



# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population
  - The errors are *statistically independent* from one another



# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population
  - The errors are *statistically independent* from one another
3. The expected value of the errors is always zero



# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population
  - The errors are *statistically independent* from one another
3. The expected value of the errors is always zero
4. The independent variables are not too strongly *collinear*



# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population
  - The errors are *statistically independent* from one another
3. The expected value of the errors is always zero
4. The independent variables are not too strongly *collinear*
5. The independent variables are measured *precisely*





# Regression Assumptions

1. Model is *linear in parameters*
2. The data are a *random sample* of the population
  - The errors are *statistically independent* from one another
3. The expected value of the errors is always zero
4. The independent variables are not too strongly *collinear*
5. The independent variables are measured *precisely*
6. The residuals have *constant variance*



# Regression Assumptions

7. The errors are normally distributed



# Regression Assumptions

- 7. The errors are normally distributed
- 8. The model is correctly specified



# Regression Assumptions

- 7. The errors are normally distributed
- 8. The model is correctly specified

If all these conditions hold, then OLS estimators are **BLUE** –



# Regression Assumptions

- 7. The errors are normally distributed
- 8. The model is correctly specified

If all these conditions hold, then OLS estimators are **BLUE** – Best Linear Unbiased Estimators



# Regression Assumptions

In real life, all the conditions may not be met.



# Regression Assumptions

In real life, all the conditions may not be met.

We need to:

1. Check if the assumptions are holding up



# Regression Assumptions

In real life, all the conditions may not be met.

We need to:

1. Check if the assumptions are holding up
2. If not, assess how to correct for violations






# Regression Results: Assumptions

## CHECKING IF ASSUMPTIONS ARE VALID


### 1. Check for linearity – plot the residuals against each IV

- If data is linearly related, we should see no pattern in the plot



# Regression Results: Assumptions

If relationship is non-linear?



# Regression Results: Assumptions

## 2. The residuals should have constant variance – homoscedasticity


- Plot the residuals against Predicted Y



# Regression Results: Assumptions

## 2. The residuals should have constant variance – homoscedasticity

- Plot the residuals against Predicted  $\hat{Y}$




Homoscedasticity




# Regression Results: Assumptions

## 2. The residuals should have constant variance – homoscedasticity

- Plot the residuals against Predicted Y



Homoscedasticity



Heteroscedasticity



# Regression Results: Assumptions

**If errors are heteroscedastic?**



# Regression Results: Assumptions

## **If errors are heteroscedastic?**

- The presence of heteroscedasticity does not imply bias in the estimates



# Regression Results: Assumptions

## If errors are heteroscedastic?

- The presence of heteroscedasticity does not imply bias in the estimates
- Heteroscedasticity leads to bias in the standard errors, leading to issues with hypothesis testing and confidence intervals





# Regression Results: Assumptions

## If errors are heteroscedastic?

- The presence of heteroscedasticity does not imply bias in the estimates
- Heteroscedasticity leads to bias in the standard errors, leading to issues with hypothesis testing and confidence intervals
  - Std error is a measure of variance, and therefore if standard errors are biased, then hypothesis test results will be biased leading to wrong inferences



# Regression Results: Assumptions


**3. The residuals are normally distributed**



# Regression Results: Assumptions

## 3. The residuals are normally distributed

- Histogram or probability plot for the residuals



# Regression Results: Assumptions

**If residuals are not normally distributed?**



# Regression Results: Assumptions

**If residuals are not normally distributed?**

Hypothesis test outcomes may be invalid, though less of an issue with large samples



# Regression Results: Assumptions

## 4. The IVs are not too correlated - multicollinearity

- The IVs should not be highly correlated to one another



# Regression Results: Assumptions

## 4. The IVs are not too correlated - multicollinearity

- The IVs should not be highly correlated to one another
- Check pairwise correlations, or generate VIF



# Regression Results: Assumptions

**4. If some of the IVs are highly correlated?**





# Regression Results: Assumptions

## 4. If some of the IVs are highly correlated?

- Estimates are not biased, but the standard errors are inflated, leading to misleading hypothesis test outcomes –



# Regression Results: Assumptions

## 4. If some of the IVs are highly correlated?

- Estimates are not biased, but the standard errors are inflated, leading to misleading hypothesis test outcomes –

Either drop the correlated variables, or combine them



# Multiple Linear Regression

## MODELING TECHNIQUES

Running a model given data is an easy task given that all the computation is done via SAS or Excel



# Multiple Linear Regression

## MODELING TECHNIQUES

Running a model given data is an easy task given that all the computation is done via SAS or Excel

**The skill of an analyst lies in generating the right model to understand and solve for the business issue at hand**



# Multiple Linear Regression

## MODELING TECHNIQUES

Running a model given data is an easy task given that all the computation is done via SAS or Excel

**The skill of an analyst lies in generating the right model to understand and solve for the business issue at hand**

The first model, or naïve model that is generated from data is usually used as a starting point



# Multiple Linear Regression

## MODELING TECHNIQUES

Running a model given data is an easy task given that all the computation is done via SAS or Excel

**The skill of an analyst lies in generating the right model to understand and solve for the business issue at hand**

The first model, or naïve model that is generated from data is usually used as a starting point

- Depending on domain understanding and modeling techniques knowledge, typically many models are run before arriving at a final model



# Multiple Linear Regression

## MODELING TECHNIQUES

Running a model given data is an easy task given that all the computation is done via SAS or Excel

**The skill of an analyst lies in generating the right model to understand and solve for the business issue at hand**

The first model, or naïve model that is generated from data is usually used as a starting point

- Depending on domain understanding and modeling techniques knowledge, typically many models are run before arriving at a final model

How can multiple models be run using the same data?



# Multiple Linear Regression

## MODELING TECHNIQUES

### Step-wise Regression

It may be useful to run models adding (or removing) one variable at a time





# Multiple Linear Regression

## MODELING TECHNIQUES

### Step-wise Regression

It may be useful to run models adding (or removing) one variable at a time

Two types of step-wise regressions:



# Multiple Linear Regression

## MODELING TECHNIQUES

### Step-wise Regression

It may be useful to run models adding (or removing) one variable at a time

Two types of step-wise regressions:

- Forward – Add one variable at a time



# Multiple Linear Regression

## MODELING TECHNIQUES

### Step-wise Regression

It may be useful to run models adding (or removing) one variable at a time

Two types of step-wise regressions:

- Forward – Add one variable at a time
- Backward – Remove one variable at a time



# Multiple Linear Regression

## MODELING TECHNIQUES

### **Forward Step-wise regression**

Variables are added one at a time until the model cannot be significantly improved by adding another variable

- Note that the variable order we use to add has an impact, so multiple step-wise forward regression models could be run before arriving at a best model

### **Backward Step-wise regression**

This approach is the reverse, where we start with a model that has all explanatory variables, and variables are dropped one by one based on p-value (highest p-value dropped first).

- Re-run model without the variable dropped, and then drop next variable with highest p-value. Continue till no other variables can be dropped based on a pre-determined cut-off value (5%, 10%)



# Multiple Linear Regression

## MODELING TECHNIQUES

Other modeling techniques could include:



# Multiple Linear Regression

## MODELING TECHNIQUES

Other modeling techniques could include:

- Transforming variables – use log transformation for example



# Multiple Linear Regression

## MODELING TECHNIQUES

Other modeling techniques could include:

- Transforming variables – use log transformation for example
- Creating interaction variables – trying to capture impact of variable A and Variable B together



# Multiple Linear Regression

## MODELING TECHNIQUES

Other modeling techniques could include:

- Transforming variables – use log transformation for example
- Creating interaction variables – trying to capture impact of variable A and Variable B together
- Aggregating or disaggregating variables – Adding up marketing, or disaggregating promotions





# Multiple Linear Regression

## MODELING TECHNIQUES

Other modeling techniques could include:

- Transforming variables – use log transformation for example
- Creating interaction variables – trying to capture impact of variable A and Variable B together
- Aggregating or disaggregating variables – Adding up marketing, or disaggregating promotions
- Creating stock variables



# Recap

- Simple Linear Regression
- Multiple Linear Regression
- Assumptions



# Coming Up

## Regression Analysis

### Case Study



# THANK YOU

