1

GATE-CS.51

EE23BTECH11046 - Poluri Hemanth*

Question: Consider the following recurrence:

$$f(1) = 1;$$

 $f(2n) = 2f(n) - 1$, for $n \ge 1$;
 $f(2n+1) = 2f(n) + 1$, for $n \ge 1$.

Then, which of thefollowing is/are TRUE?

- (A) $f(2^n 1) = 2^n 1$
- $(B) f(2^n) = 1$
- (C) $f(5 \cdot 2^n) = 2^{n+1} + 1$
- (D) $f(2^n + 1) = 2^n + 1$
- [GATE-CS.51 2022]

Solution:

(A)

For k=1.

$$f(2^{k} - 1) = 1$$

$$= 2^{k} - 1$$

$$f(2^{k+1} - 1) = f(2(2^{k} - 1) + 1)$$

$$= 2f(2^{k} - 1) + 1$$

$$= 2(2^{k} - 1) + 1$$

$$= 2^{k+1} - 1$$
(1)
(3)
(4)
(5)
(6)

Hence statement A is TRUE (B)

Let $f(2^n) = 1$ for any random $n \ge 1$

$$f(2^{n+1}) = 2f(2^n) - 1 (7)$$

= 1 (8)

Therefore B will be proved

$$f(2^0) = f(1)$$
 (9)
= 1 (10)

Hence $f(2^n) = 1$ for every $n \ge 1$ value. So statement B is TRUE.

(C) Let, $f(5 \cdot 2^n) = 2^{n+1} + 1$ is true for any n value,

$$f(5 \cdot 2^{n+1}) = f(2(5 \cdot 2^n))$$
(11)
= 2 f(5 \cdot 2^n) + 1 (12)

$$=2(2^{n+1}+1)-1$$

(13)

$$=2^{n+2}+1$$
 (14)

Therefore C will be proved

$$f(5) = f(2 \cdot 2 + 1) \tag{15}$$

$$= 2f(2) + 1 \tag{16}$$

$$=3 \tag{17}$$

Hence statement C is TRUE.

(D)

For any $n \ge 1$

$$f(2^{n} + 1) = 2f(2^{n-1}) + 1 (18)$$

From (B)

$$f(2^n + 1) = 3 (19)$$

Hence statement D is FALSE.