

AREA

THEME

Title



Henrik Andrews

Norwegian university of Science and Technology

November 15, 2023

-abstract -sammendrag -acknowledgments -list of figure -list of tables

1 Introduction

References

- Ajello, M., Costamante, L., Sambruna, R. M., Gehrels, N., Chiang, J., Rau, A., Escala, A., Greiner, J., Tueller, J., Wall, J. V., and Mushotzky, R. F. (2009). The evolution of swift/bat blazars and the origin of the mev background. *The Astrophysical Journal*, 699(1):603.
- Hogg, D. W. (2000). Distance measures in cosmology.
- Jacobsen, I. B., Wu, K., On, A. Y. L., and Saxton, C. J. (2015). High-energy neutrino fluxes from AGN populations inferred from X-ray surveys. *Mon. Not. Roy. Astron. Soc.*, 451(4):3649–3663.
- Netzer, H. (2015). Revisiting the unified model of active galactic nuclei. *Annual Review of Astronomy and Astrophysics*, 53(1):365–408.
- Shields, G. A. (1999). A brief history of active galactic nuclei. *Publications of the Astronomical Society of the Pacific*, 111(760):661–678.
- Ueda, Y., Akiyama, M., Ohta, K., and Miyaji, T. (2003). Cosmological evolution of the hard x-ray active galactic nucleus luminosity function and the origin of the hard x-ray background. *The Astrophysical Journal*, 598(2):886.

2 Cosmological distances

In order to investigate sources very far away from an observer it is important to understand the influence this distance has on your desired observable. Therefore in astrophysics and astronomy in general there are distances created to take into account the effects of an expanding universe.

2.1 cosmographic parameters

The most notorious parameter is the Hubble constant H_0 . This parameter sets the recession speed of a point at proper distance d and our current position via this equation.

$$v = H_0 d \quad (1)$$

The subscript 0 refers to the present epoch since in general H_0 changes with time. The value of H_0 is quite debated so I will follow the nomenclature of D. Hogg and write it as a parameterized equation.

$$H_0 = 100 \frac{km}{s} \frac{1}{Mpc} h$$

where h is a dimensionless number that according to current knowledge can take the value between 0.5 to 0.8

The Hubble constant has units of inverse time and therefore one can define the Hubble time as

$$t_H = \frac{1}{H_0}$$

The constant also has units of speed, and therefore we can define the Hubble distance.

$$D_H = \frac{c}{H_0}$$

2.2 Components of the universe

In this paper and in most articles one refers to the flat lambda CDM model to parameterize the contents of the universe and by extension the properties of its expansion. Here two important parameters to define are the mass density of the universe ρ_0 and the cosmological constant Λ . These variables which change with time also define the metric tensor in general relativity and allow us to model the curvature of the universe given an initial configuration. njaaa, skriv på nytt.

One can write these into dimensionless variables as such

$$\Omega_m = \frac{8\pi G\rho_0}{3H_0^2}$$

$$\Omega_\Lambda = \frac{\Lambda c^2}{3H_0^2}$$

In general one has a third density parameter Ω_k which defines the curvature of spacetime and is defined from

$$\Omega_m + \Omega_\Lambda + \Omega_k = 1$$

The flat lambda CDM has three components, the dark energy density, the dark matter density, and the ordinary matter parameter. In the rest of this paper and other articles used by this dissertation the values of these components are represented by their Ω parameter and one has a universe with $\Omega_\Lambda = 0.7$, and $\Omega_m = 0.3$ also known as the flat lambda since the curvature parameter is 0

2.3 Redshift

Redshift is defined as the fractional Doppler shift of emitting light. The Doppler effect is a known effect on different observables in our universe where the relative motion of sources to observers will impact the observable. The redshift is quantified for a light source as

$$z = \frac{\nu_e}{\nu_o} - 1 = \frac{\lambda_o}{\lambda_e} - 1 \quad (2)$$

Here o refers to the observed quantity and e the emitted. If one want to connect this redshift to the velocity of the observed object one needs to go to general relativity. There is an analog in special but one omits it here since one does not use it. The important factor is that redshift is an observable and can help us determine distances of objects. especially if they are far away the relative velocity becomes negligible and only the effect of the expansion of the universe become important.

2.4 Comoving distance

The comoving distance or more clearly the line of sight distance for an observer located here at earth is a foundational distance measure in cosmography. All other distance measures can be derived from it. One derives it by defining the small co-moving distance δD_c . This quantity defines the distance between two objects that remains constant when both objects expand with the Hubble flow. One can think of it as a proper distance from relativity since it is constant in all "time-frames". If one wants the total comoving distance one integrates all δD_c in the line of sight from $z = 0$ to the object. In order to do this one need to take into account the different densities in the universe, from ? one defines the function

$$E(z) = \sqrt{\Omega_m(z+1)^3 + \Omega_k(1+z)^2 + \Omega_\Lambda} \quad (3)$$

This function is defined by the density parameters defined above and also the redshift z . One can also relate this to the measured Hubble constant as measured by an hypothetical observer at redshift z via $H(z) = H_0 E(z)$.

One then recives the comoving distance D_c from

$$D_c = D_H \int_0^z \frac{dz}{E(z)} \quad (4)$$

2.5 Comoving distance part 2

D_c is the line of sight of an object and its observer but given different space-time geometry that line is distorted. If one looks a two objects, both at redshift z the distance between them will be given as as a function of the angle between them. If they are sperated by an angle $d\theta$ then the distance between them will be $d\theta D_m$ where D_m is the comoving distance

$$D_m = \begin{cases} D_H \frac{1}{\sqrt{\Omega_k}} \sinh\left(\frac{\sqrt{\Omega_k} D_c}{D_H}\right) & \text{if } \Omega_k > 0 \\ D_c & \text{if } \Omega_k = 0 \\ D_H \frac{1}{\sqrt{|\Omega_k|}} \sin\left(\frac{\sqrt{|\Omega_k|} D_c}{D_H}\right) & \text{if } \Omega_k < 0 \end{cases}$$

The different cases are dependent on the curvature of the universe, and one can see that one enters hyperbolic geometry or spherical geometry based on the different curvatures. The true curvature of the universe is still a mystery but recent studies suggest that it is flat, as expected.

2.6 Angular distance

The anglul

2.7 Luminosity distance

The luminosity distance D_l is defined through the relation between the bolometrix flux F of a source and its bolometric luminosity L . The flux is the amount of energy per unit time per unit area and the luminosity is the total amount of energy per unit time. The luminosity distance is defined as

$$D_l = \sqrt{\frac{L}{4\pi F}} \quad (5)$$

In more clearer words it describes the total loss in energy due to lights travel across an expanding universe. The total observed flux an observer will see will be different from different distances. the intrinsic luminosity emmited at the source

It is related to the transverse comoving distance via

$$D_l = (1+z)D_m \quad (6)$$

This of course is for bolometric quantities, but if one wants to calculate the spectral flux/ differential flux one need to take into account a correction. This correction comes from the fact that one is viewing a redshifted object. The

object is emitting in a different band than observed. The spectrum of the differential flux F_ν is related to the spectral luminosity via

$$F_\nu = (1+z) \frac{L_{(1+z)\nu}}{L_\nu} \frac{L_\nu}{4\pi D_l^2} \quad (7)$$

If one wishes to translate an observed spectral flux of particles to a density distribution one needs to take into account this redshift effect

3 Particle theory

3.1 Emission lines

When a particle is ionized by the continuum radiation of a source it will emit a photon when it returns to its ground state. This photon will have a specific wavelength that is determined by the energy difference between the two states. This wavelength is called the emission line of the particle. When looking at dynamical systems with high velocities the doppler shift of these emission lines becomes important since it will affect the observed spectra of the source.

<https://lweb.cfa.harvard.edu/~pberlind/whipple/agn.html>

3.1.1 Broad emission lines

Broad emission lines in the case of AGN are formed from the high density gas clouds located close to the central black hole. The high density parameter is inferred from the fact that one only sees broad emission from permitted line transitions (e.g. hydrogen Lyman and Balmer, iron II, and magnesium II). High densities allow for collisional de-excitation. And the broadening is an indication that these gas clouds are moving at huge velocities around the massive objects. This implies that they are located close to the black hole.

3.1.2 Narrow emission lines

Narrow emission lines are on the other hand formed in low density gas clouds. The low densities are inferred from the fact that one sees both permitted and forbidden line transitions. They are narrow lines due to their velocities being substantially lower than the inner most gas clouds.

4 Neutrino production

We will be discussing two types of neutrino generation. The pp chain and $p\gamma$ production

5 Types of AGNs

5.1 Active galactic nuclei

Active Galactic Nuclei (AGNs) is an interesting field in astrophysical studies. Since their discovery, there has been rapid advancement in understanding these phenomena. Today, AGNs are known to be among the brightest entities in the night sky, but they only gained significant attention in the 1950s. This shift occurred with the arrival of new radio observations, which revealed a new type of quasi-stellar object through the discovery of Quasars.

Initially, these luminous objects, characterized by broad, unidentifiable spectral lines, were enigmatic to scientists in the early 1960s. However, with the identification of more sources and their optical parts, it became clear that these were not stars but a distinct class of celestial objects. Research done by M. Schmidt on of the emission lines from the Quasar 3C 273 opened the interpretation of these celestial objects. He found that the emission lines of quasars were similar to hydrogen, but were redshifted by a factor of 0.158, an exceptionally high value at the time. Observations at the same time also revealed significant variability in quasar luminosity, suggesting that these objects were no larger than one light year across.

These observations lead to the speculation of super luminous objects located very far away from earth. The problem was that such objects had no reasonable explanation at the time. It was not until the mid 1960 early 1970s when modern cosmology was afoot that more of these issues were resolved.

Observation of the surrounding galaxy of AGNs with matching redshift and observation of gravitational lensing cemented the distances of these objects. In addition the modern view of black holes which had only been a theory in the 1950s came to fruition and the modern model of a AGN was born. This modern perspective views AGNs as supermassive black holes that accrete matter from surrounding accretion disks. This accretion releases large amounts of energy and has also according to processes such as the Blandford-Znajek process (1977), been shown to produce relativistic jets, when the black hole is rotating.

In the most recent times a landmark achievement was achieved in March 2021, when scientists associated with the Event Horizon Telescope project presented the first image of the supermassive black hole at the center of the Messier 87 galaxy, located 55 million light-years away. This image, showing a bright ring surrounding a dark central region, aligns with predictions for an accreting supermassive black hole, reinforcing our understanding of these powerful cosmic sources.

5.2 AGN structure and classification

The modern view of AGNs is unified model that combined different categories of powerful luminous objects visible in the night sky. These distinctions that astronomers made still have value, but to understand an AGN it is important to get a picture of the modern structure of an AGN

An active galactic nucleus is defined as a galaxy containing a massive accreting black hole. This mass according to ? is defined as $M_{BH} > 10^5 M_{\odot}$. AGNs also contain an Eddington ratio exceeding the limit of $\frac{L_{AGN}}{L_{Edd}} = 10^{-5}$, where L_{AGN} is the bolometric luminosity, and L_{Edd} is the Eddington luminosity for a solar composition gas. These definitions are help constrain what galaxies might contain an AGN, for example it excludes the Milky Way by these criteria, but it fails to capture the full structure definition of an AGN. Therefore the structure of most AGNs will include several of the following components.

- A close rotational dominated accretion disc around the SMBH. The thickness defining this accretion flow will distinguish different AGNs. One example is an optically thin accretion disks that sometimes become advection dominated. These flows will be referred to as radiation inefficient accretion flows (RIAF) due to the special nature of the disk.
- high density gas clouds that are said to be dust free moving at high velocities close to the black hole, in the so called broad line region (BLR)
- Low density gas clouds that move at lower velocities further away from the black hole in the so called narrow line region (NLR)
- A axisymmetric structure of dust that is responsible for the obscuration of the central region of the AGN. This is called the torus. It lies at a luminosity depended distance from the SMBH, but according to ? this is around 0.1 - 10 pc depending on the luminosity.
- A relativistic jet that is powered by the accretion disk. This is not always present but is a common feature of AGNs.

The reader is directed to image ?? for a visual representation of the different components.




Figure 1: AGN unification

5.3 Types of AGNs

<https://astrobites.org/guides/galaxy-and-agn-types/>

Before the unification of the AGNs astronomers named the puzzling objects based on their observational properties. These names are still used to this day and are somewhat useful since their observational properties are important parameters for further study. The different classifications are important in understanding which objects could have the potential to produce the different observables one looks for in the night sky. There is a lot of talk around AGNs being possible sources for ultra high energy cosmic rays (UHECRs) and neutrinos. This is yet to be confirmed, but the theoretical framework for the necessary particle acceleration is there. Therefore it seems appropriate to discuss the different types of AGNs and their observational properties.

5.3.1 Type I and II AGNs

One distinguishes type I and type II AGNs based on the presence of broad emission lines. In other words this distinction is a matter of a visible nucleus or not. Type I refers to sources whose nucleus is exposed to the observer and whose spectrum has both narrow and broad emission lines. Type II refers to sources whose nucleus is obscured by a torus and therefore only has narrow emission lines.

5.3.2 Blazars

The most extreme class of AGN. These sources are distinguished by their relativistic jets that are pointed towards the observer. This jet produces both synchrotron and Inverse Compton gamma rays and are extremely variable over short timescales. The emission is also highly polarized. Often and including in this paper one divides blazars into subgroups based on the emission lines. The two most common are BL Lacs and Flat spectrum radio quasars (FSRQs). The difference between these two is the presence of broad emission lines. BL Lacs have no broad emission lines while FSRQs do.

5.3.3 Radio galaxies

As the name suggests these sources are very bright in the radio band. They usually refer to AGN viewed edge on, where the torus might block the emissions from the accretion disk. The orientation of radio galaxies gives way for strong synchrotron radiation, and therefore they are often used to study the jet structure of AGNs.

Model	A	L_{star}	γ_1	γ_2	v_1	v_2	z_c	L_c	α	$corr-fac$
RG-SLDDE	8.375e-07	2.138e+44	2.15	1.10	4.00	-1.50	1.90	3.981e+44	0.317	1
AMPLE-Blazar	1.379e-07	1.810e+44	-0.87	2.73	3.45	-0.25				400
AMPLE-FSRQ	1.750e-08	2.420e+44	-50.00	2.49	3.67	-0.30				400
APLE-BLlac	8.300e-08	1.000e+44	2.61	-0.79						400

Table 1: X-ray LF parameters

5.3.4 Seyfert galaxies

Spiral galaxies the have a birght nucleus. They are bright in the optical band and have a smaller active region than radio galaxies. They are often divided into two groups Seyfert I and Seyfert II where the distinction comes from type I and II. They also show quite high variablilty indicating a small emmiting region.

<https://iopscience.iop.org/article/10.1086/305813/fulltext/37493.text.html>: :text=,line

6 AGN geometry

7 luminosity functions

*what is a LF * X-ray LF *

a luminosity function is a function that describes the distribution of objects for a population of celestial sources, such as galaxies or quasars. It is a powerful tool for understanding the properties and evolution of these objects, as well as the larger-scale structure of the universe. The function describes how a population varies based on luminosity but also crucially on its comoving volume element. We usually talk about the differential luminosity function given as

$$\frac{d\Psi(L, z)}{dL} = \frac{d^2 N(L, z)}{dL dV_c} \quad (8)$$

The quantity of interest is now a number density which can be very useful in deriving observed flux of different objects here on earth.

Several articles express the luminosity function in abase 10 logarithm and we note the conversion between the two.

$$\frac{d\Psi(Lx, z)}{d\text{Log}(Lx)} = \ln(10) Lx \frac{d\Psi(Lx, z)}{d(Lx)} \quad (9)$$

7.1 X-ray LF

One way of calculating the neutrino flux of AGNs is based on their connecting with x-ray radiation. Therefore in some literature, it is of interest to define the x-ray luminosity function for AGNs.

In both ? and ? they calculate the luminsity function for different types of AGNs

This is over a particular band of wavelengths

This equation does not have a simple equation to describe all types of light sources, but for different intervals of wavelengths and redshifts (z) one can observe different trends for different light sources.

In the paper ? the author uses the xray luminosity function. The X-ray luminosity function is used to describe the distribution of X-ray luminosities of objects in a specific population, such as galaxies, galaxy clusters, or active

galactic nuclei (AGNs). It provides information about the number density of objects at various X-ray luminosity levels within a given volume of the universe.

In ? there is a mention of several different models for different populations of AGNs. There she highlights two, ? and ?. They are used for different populations.

The present day XLF is presented in ? and is given by a simple power law.

$$\frac{d\Psi(Lx, 0)}{d\text{Log}(Lx)} = A \ln(10) \left(\frac{Lx}{Lc}\right)^{(1-\gamma_2)} \quad (10)$$

However there is a break with high enough score count and this break can be better fitted with a double power law