

Epidemics in Networks

Part 2 — Compartmental Disease Models

Joel C. Miller & Tom Hladish

17–19 July 2017

Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References

Recall our key questions

For SIR:

- ▶ \mathcal{P} , the probability of an epidemic.
- ▶ \mathcal{A} , the “attack rate”: the fraction infected if an epidemic happens (better named the attack ratio).
- ▶ \mathcal{R}_0 , the average number of infections caused by those infected early in the epidemic.
- ▶ $I(t)$, the time course of the epidemic.

For SIS:

- ▶ \mathcal{P}
- ▶ $I(\infty)$, the equilibrium level of infection
- ▶ \mathcal{R}_0
- ▶ $I(t)$

Simple Compartmental Models

The most common models are compartmental models.

- ▶ Continuous time or Discrete time
- ▶ SIR or SIS

Simple Compartmental Models

The most common models are compartmental models.

- ▶ Continuous time or Discrete time
- ▶ SIR or SIS

The major assumptions:

- ▶ Every individual is average.
- ▶ Every interaction of u is with a randomly chosen other individual.

Simple Compartmental Models

The most common models are compartmental models.

- ▶ Continuous time or Discrete time
- ▶ SIR or SIS

The major assumptions:



- ▶ Every individual is average.
- ▶ Every interaction of u is with a randomly chosen other individual.

Throughout:

$$S + I + R = N$$



[That is, we look at absolute number rather than proportions of the population. Unfortunately this is standard across much of the field and it causes our equations and initial conditions to be littered with N s that do nothing to help us understand what is happening. I haven't used this convention in past years' notes, so there may be typos occasionally. I've given up fighting this.]

Continuous time: Kermack–McKendrick



Assumptions:



Continuous time: Kermack–McKendrick



Assumptions:

- ▶ Individuals **recover** with rate γ .



Continuous time: Kermack–McKendrick



Assumptions:

- ▶ Individuals **recover** with rate γ .
- ▶ Infected individuals **transmit to others** at rate $\beta \langle K \rangle$ (usually we combine these into a single parameter).
 - ▶ β represents the transmission rate of the disease per partnership.
 - ▶ $\langle K \rangle$ represents the typical number of partners of an infected individual at any time.



Continuous time: Kermack–McKendrick



Assumptions:

- ▶ Individuals **recover** with rate γ .
- ▶ Infected individuals **transmit to others** at rate $\beta \langle K \rangle$ (usually we combine these into a single parameter).
 - ▶ β represents the transmission rate of the disease per partnership.
 - ▶ $\langle K \rangle$ represents the typical number of partners of an infected individual at any time.
- ▶ The proportion of transmissions that go to susceptible individuals is S/N .

Continuous time: Kermack–McKendrick




Assumptions:

- ▶ Individuals **recover** with rate γ .
- ▶ Infected individuals **transmit to others** at rate $\beta \langle K \rangle$ (usually we combine these into a single parameter).
 - ▶ β represents the transmission rate of the disease per partnership.
 - ▶ $\langle K \rangle$ represents the typical number of partners of an infected individual at any time.
- ▶ The proportion of transmissions that go to susceptible individuals is S/N .
- ▶ An implicit assumption is that each interaction is with a new randomly infected individual.

Stochastic simulation — SIR case


What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?

Cumulative infections



Stochastic simulation — SIR case


What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?



Stochastic simulation — SIR case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?

Cumulative infections




$$N = 1250$$

Stochastic simulation — SIR case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?


Cumulative infections



Stochastic simulation — SIR case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?

Cumulative infections




The distinction between small and large outbreaks becomes clear as N increases.

Stochastic simulation — SIR case

What does the final size distribution look like?


Stochastic simulation — SIR case

What does the final size distribution look like?



Stochastic simulation — SIR case

What does the final size distribution look like?





In large populations:

Small outbreaks affect the same **number** of individuals.

Stochastic simulation — SIR case

What does the final size distribution look like?



In large populations:


Small outbreaks affect the same number of individuals.

Epidemics affect approximately the same proportion of the population

Stochastic simulation — SIS case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?


Infections



$$N = 50$$

Stochastic simulation — SIS case


What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?



Stochastic simulation — SIS case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?


Infections



$$N = 1250$$

Stochastic simulation — SIS case



What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?



Stochastic simulation — SIS case

What behavior do we see with $\beta \langle K \rangle = 2$, $\gamma = 1$?

Infections




Typically extinction occurs either early or after exponentially long time. The **proportion** infected at equilibrium is approximately the same for different population sizes.

Stochastic simulation — SIS case

What does the “equilibrium” distribution look like?


Stochastic simulation — SIS case

What does the “equilibrium” distribution look like?



Stochastic simulation — SIS case

What does the “equilibrium” distribution look like?



The equilibrium proportion infected is about the same for different population sizes.

Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References



Recall our key questions



For SIR:

- ▶ \mathcal{P} , the probability of an epidemic.
- ▶ \mathcal{A} , the “attack rate”: the fraction infected if an epidemic happens (better named the attack ratio).
- ▶ \mathcal{R}_0 , the average number of infections caused by those infected early in the epidemic.
- ▶ $I(t)$, the time course of the epidemic.

For SIS:



- ▶ \mathcal{P}
- ▶ $I(\infty)$, the equilibrium level of infection
- ▶ \mathcal{R}_0
- ▶ $I(t)$







- ▶ SIR equations are:

$$\dot{S} =$$



- ▶ SIR equations are:



$$\dot{S} = -\beta \langle K \rangle IS/N$$



- ▶ SIR equations are:

$$\dot{S} = -\beta \langle K \rangle IS/N$$



$$\dot{I} =$$



- ▶ SIR equations are:

$$\dot{S} = -\beta \langle K \rangle IS/N$$

$$\dot{I} = \beta \langle K \rangle IS/N - \gamma I$$





- ▶ SIR equations are:

$$\dot{S} = -\beta \langle K \rangle IS/N$$

$$\dot{I} = \beta \langle K \rangle IS/N - \gamma I$$

$$\dot{R} =$$





- ▶ SIR equations are:

$$\dot{S} = -\beta \langle K \rangle IS/N$$

$$\dot{I} = \beta \langle K \rangle IS/N - \gamma I$$

$$\dot{R} = \gamma I$$



- ▶ SIR equations are:

$$\dot{S} = -\beta \langle K \rangle IS/N$$

$$\dot{I} = \beta \langle K \rangle IS/N - \gamma I$$

$$\dot{R} = \gamma I$$



- ▶ SIS equations are

$$\dot{S} = -\beta \langle K \rangle IS/N + \gamma I$$


$$\dot{I} = \beta \langle K \rangle IS/N - \gamma I$$

Comparison

SIR:



SIS:



Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References

Recall our key questions

For SIR:

- ▶ \mathcal{P} , the probability of an epidemic.
- ▶ \mathcal{A} , the “attack rate”: the fraction infected if an epidemic happens (better named the attack ratio).
- ▶ \mathcal{R}_0 , the average number of infections caused by those infected early in the epidemic.
- ▶ $I(t)$, the time course of the epidemic.

For SIS:

- ▶ \mathcal{P}
- ▶ $I(\infty)$, the equilibrium level of infection
- ▶ \mathcal{R}_0
- ▶ $I(t)$

\mathcal{R}_0

An important quantity in disease modeling is \mathcal{R}_0 .

\mathcal{R}_0

An important quantity in disease modeling is \mathcal{R}_0 .

- ▶ \mathcal{R}_0 is the expected number of infections caused by an individual infected early in the epidemic.

\mathcal{R}_0

An important quantity in disease modeling is \mathcal{R}_0 .

- ▶ \mathcal{R}_0 is the expected number of infections caused by an individual infected early in the epidemic.
- ▶ If $\mathcal{R}_0 < 1$ epidemics cannot occur.
- ▶ If $\mathcal{R}_0 > 1$ epidemics can occur, **but are not guaranteed**.

\mathcal{R}_0

An important quantity in disease modeling is \mathcal{R}_0 .

- ▶ \mathcal{R}_0 is the expected number of infections caused by an individual infected early in the epidemic.
- ▶ If $\mathcal{R}_0 < 1$ epidemics cannot occur.
- ▶ If $\mathcal{R}_0 > 1$ epidemics can occur, **but are not guaranteed**.

So it's an important threshold parameter, but it doesn't address the probability of epidemics.

\mathcal{R}_0 calculation

The calculation of \mathcal{R}_0 is the same for SIR and SIS:

- ▶ Under our assumptions, every interaction an infected individual has is with a new randomly chosen individual.
- ▶ Early in the epidemic, the probability it is with a susceptible individual is $S/N \approx 1$.
- ▶ The typical infection duration is $1/\gamma$.
- ▶ The transmission rate during infection is $\beta \langle K \rangle$.
- ▶ So the number of new infections is $\mathcal{R}_0 = \beta \langle K \rangle / \gamma$.

Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References

Recall our key questions


For SIR:

- ▶ \mathcal{P} , the probability of an epidemic.
- ▶ \mathcal{A} , the “attack rate”: the fraction infected if an epidemic happens (better named the attack ratio).
- ▶ \mathcal{R}_0 , the average number of infections caused by those infected early in the epidemic.
- ▶ $I(t)$, the time course of the epidemic.

For SIS:


- ▶ \mathcal{P}
- ▶ $I(\infty)$, the equilibrium level of infection
- ▶ \mathcal{R}_0
- ▶ $I(t)$

SIS epidemic probability on annealed networks



Consider an individual u with degree k who becomes infected at time $t = t_0$. Define $i_m(t)$ and $r_m(t)$ to be the probability u has transmitted to m individuals and is infectious or recovered.


SIS epidemic probability on annealed networks



Consider an individual u with degree k who becomes infected at time $t = t_0$. Define $i_m(t)$ and $r_m(t)$ to be the probability u has transmitted to m individuals and is infectious or recovered.

- ▶ The probability of transmitting at least once before recovering is $\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma}$.


SIS epidemic probability on annealed networks



Consider an individual u with degree k who becomes infected at time $t = t_0$. Define $i_m(t)$ and $r_m(t)$ to be the probability u has transmitted to m individuals and is infectious or recovered.

- ▶ The probability of transmitting at least once before recovering is $\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma}$.
- ▶ The probability the first m events are transmissions is $\left[\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right]^m$.

SIS epidemic probability on annealed networks



Consider an individual u with degree k who becomes infected at time $t = t_0$. Define $i_m(t)$ and $r_m(t)$ to be the probability u has transmitted to m individuals and is infectious or recovered.

- ▶ The probability of transmitting at least once before recovering is $\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma}$.
- ▶ The probability the first m events are transmissions is $\left[\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right]^m$.
- ▶ For exactly m transmissions before recovery it is

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

For reasons that will be clear later, we define

$$f(x) = \sum_m r_m(\infty) x^m$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

For reasons that will be clear later, we define

$$f(x) = \sum_m r_m(\infty) x^m = \frac{\gamma}{(\langle K \rangle \beta + \gamma) \left(1 - \frac{\langle K \rangle \beta x}{\langle K \rangle \beta + \gamma} \right)}$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

For reasons that will be clear later, we define

$$f(x) = \sum_m r_m(\infty) x^m = \frac{\gamma}{(\langle K \rangle \beta + \gamma) \left(1 - \frac{\langle K \rangle \beta x}{\langle K \rangle \beta + \gamma} \right)} = \frac{\gamma}{\langle K \rangle \beta (1 - x) + \gamma}$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

For reasons that will be clear later, we define

$$f(x) = \sum_m r_m(\infty) x^m = \frac{\gamma}{(\langle K \rangle \beta + \gamma) \left(1 - \frac{\langle K \rangle \beta x}{\langle K \rangle \beta + \gamma}\right)} = \frac{\gamma}{\langle K \rangle \beta (1 - x) + \gamma}$$

- ▶ The probability of extinction after the first individual is the probability it recovers without transmitting.

$$f(0) = r_0(\infty)$$

We have

$$r_m(\infty) = \left(\frac{\langle K \rangle \beta}{\langle K \rangle \beta + \gamma} \right)^m \frac{\gamma}{\langle K \rangle \beta + \gamma}$$

For reasons that will be clear later, we define

$$f(x) = \sum_m r_m(\infty) x^m = \frac{\gamma}{(\langle K \rangle \beta + \gamma) \left(1 - \frac{\langle K \rangle \beta x}{\langle K \rangle \beta + \gamma}\right)} = \frac{\gamma}{\langle K \rangle \beta (1-x) + \gamma}$$

- ▶ The probability of extinction after the first individual is the probability it recovers without transmitting.

$$f(0) = r_0(\infty)$$

- ▶ The probability of reaching no further than the first generation of offspring is the probability that none of the first generation individuals causes a transmission.

$$\sum_m r_m(\infty) [f^{(0)}]^m = f(f(0)) = f^{(2)}(0)$$

[The superscript with parentheses denotes function iteration]


- ▶ The probability of reaching no further than generation g is the probability that none of the first generation individuals causes a transmission chain of length longer than $g - 1$.

$$\sum_m r_m(\infty) [f^{(g-1)}(0)]^m = f^{(g)}(0)$$

- ▶ The probability the outbreak goes extinct in a finite number of generations (in an infinite population) is $\lim_{g \rightarrow \infty} f^{(g)}(0)$.


Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0.




Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0.




Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0.




Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0.




Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0. A more direct way is with a “cobweb diagram”.



Cobweb diagrams

Consider $f(x) = \frac{1+x^3}{2}$. We can keep iterating f on 0. A more direct way is with a “cobweb diagram”.



Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References

Recall our key questions

For SIR:

- ▶ \mathcal{P} , the probability of an epidemic.
- ▶ \mathcal{A} , the “attack rate”: the fraction infected if an epidemic happens (better named the attack ratio).
- ▶ \mathcal{R}_0 , the average number of infections caused by those infected early in the epidemic.
- ▶ $I(t)$, the time course of the epidemic.

For SIS:

- ▶ \mathcal{P}
- ▶ $I(\infty)$, the equilibrium level of infection
- ▶ \mathcal{R}_0
- ▶ $I(t)$

SIS equilibrium size

- The SIS equations are

$$\dot{S} = -\beta \langle K \rangle \frac{IS}{N} + \gamma I$$

$$I = \beta \langle K \rangle \frac{IS}{N} - \gamma I$$

SIS equilibrium size

- ▶ The SIS equations are

$$\dot{S} = -\beta \langle K \rangle \frac{IS}{N} + \gamma I$$

$$I = \beta \langle K \rangle \frac{IS}{N} - \gamma I$$

- ▶ Setting $\dot{S} = 0$, we conclude that either $I_{\text{eq}} = 0$ (the disease has gone extinct) or $\beta \langle K \rangle S_{\text{eq}}/N = \gamma$.

SIS equilibrium size

- ▶ The SIS equations are

$$\dot{S} = -\beta \langle K \rangle \frac{IS}{N} + \gamma I$$

$$I = \beta \langle K \rangle \frac{IS}{N} - \gamma I$$

- ▶ Setting $\dot{S} = 0$, we conclude that either $I_{\text{eq}} = 0$ (the disease has gone extinct) or $\beta \langle K \rangle S_{\text{eq}}/N = \gamma$.
- ▶ Taking the second case gives $S_{\text{eq}} = \gamma N / \beta \langle K \rangle$ and $I_{\text{eq}} = N(1 - \gamma / \beta \langle K \rangle)$.

SIS equilibrium size

- ▶ The SIS equations are

$$\dot{S} = -\beta \langle K \rangle \frac{IS}{N} + \gamma I$$

$$I = \beta \langle K \rangle \frac{IS}{N} - \gamma I$$

- ▶ Setting $\dot{S} = 0$, we conclude that either $I_{\text{eq}} = 0$ (the disease has gone extinct) or $\beta \langle K \rangle S_{\text{eq}}/N = \gamma$.
- ▶ Taking the second case gives $S_{\text{eq}} = \gamma N / \beta \langle K \rangle$ and $I_{\text{eq}} = N(1 - \gamma / \beta \langle K \rangle)$.
- ▶ If $\gamma > \beta \langle K \rangle$ the disease must die out.

SIS equilibrium size

- ▶ The SIS equations are

$$\dot{S} = -\beta \langle K \rangle \frac{IS}{N} + \gamma I$$

$$I = \beta \langle K \rangle \frac{IS}{N} - \gamma I$$

- ▶ Setting $\dot{S} = 0$, we conclude that either $I_{\text{eq}} = 0$ (the disease has gone extinct) or $\beta \langle K \rangle S_{\text{eq}}/N = \gamma$.
- ▶ Taking the second case gives $S_{\text{eq}} = \gamma N / \beta \langle K \rangle$ and $I_{\text{eq}} = N(1 - \gamma / \beta \langle K \rangle)$.
- ▶ If $\gamma > \beta \langle K \rangle$ the disease must die out.
- ▶ If $\gamma < \beta \langle K \rangle$ at equilibrium the disease has died out or reaches an equilibrium having a fraction $\gamma / \beta \langle K \rangle$ susceptible and the rest infected.

Alternate equations for SIR

Before deriving the final size relation, we derive an alternate system of equations. The system has some important properties:

Alternate equations for SIR

Before deriving the final size relation, we derive an alternate system of equations. The system has some important properties:

- ▶ There is a single governing ODE.
- ▶ It will make the final size relation trivial.
- ▶ It has a useful alternate interpretation that gives insight into equations for disease on networks.

Deriving alternate equations

- ▶ Starting from

$$\dot{S} = -\beta \langle K \rangle I \frac{S}{N}$$

Deriving alternate equations

- ▶ Starting from

$$\dot{S} = -\beta \langle K \rangle I \frac{S}{N}$$

- ▶ We write

$$\dot{S} + \frac{\beta \langle K \rangle I}{N} S = 0$$

Deriving alternate equations

- ▶ Starting from

$$\dot{S} = -\beta \langle K \rangle I \frac{S}{N}$$

- ▶ We write

$$\dot{S} + \frac{\beta \langle K \rangle I}{N} S = 0$$

- ▶ Multiplying by $e^{\xi(t)}$ we have

$$\dot{S}e^\xi + \frac{\beta \langle K \rangle I}{N} Se^\xi = 0$$

Deriving alternate equations

- ▶ Starting from

$$\dot{S} = -\beta \langle K \rangle I \frac{S}{N}$$

- ▶ We write

$$\dot{S} + \frac{\beta \langle K \rangle I}{N} S = 0$$

- ▶ Multiplying by $e^{\xi(t)}$ we have

$$S e^\xi + \frac{\beta \langle K \rangle I}{N} S e^\xi = 0$$

- ▶ We choose ξ so that $\dot{\xi} = \beta \langle K \rangle I / N$. Then e^ξ is an integrating factor. We can arbitrarily assume $\xi(0) = 0$.

Deriving alternate equations

- ▶ Starting from

$$\dot{S} = -\beta \langle K \rangle I \frac{S}{N}$$

- ▶ We write

$$\dot{S} + \frac{\beta \langle K \rangle I}{N} S = 0$$

- ▶ Multiplying by $e^{\xi(t)}$ we have

$$Se^\xi + \frac{\beta \langle K \rangle I}{N} Se^\xi = 0$$



- ▶ We choose ξ so that $\dot{\xi} = \beta \langle K \rangle I / N$. Then e^ξ is an integrating factor. We can arbitrarily assume $\xi(0) = 0$.
- ▶ We have

$$\frac{d}{dt} Se^\xi = \dot{S}e^\xi + S\dot{\xi}e^\xi = 0$$

where $\xi(0) = 0$

Deriving alternate equations

- ▶ So $Se^{\xi} = S(0)$ and thus $S(t) = S(0)e^{-\xi}$.

Deriving alternate equations

- ▶ So $Se^\xi = S(0)$ and thus $S(t) = S(0)e^{-\xi}$.
- ▶ We have $\dot{\xi} = \beta \langle K \rangle I / N$ and $\dot{R} = \gamma I$. So

$$\dot{\xi} = \frac{\beta \langle K \rangle \dot{R}}{N\gamma}$$

Deriving alternate equations

- ▶ So $Se^\xi = S(0)$ and thus $S(t) = S(0)e^{-\xi}$.
- ▶ We have $\dot{\xi} = \beta \langle K \rangle I/N$ and $\dot{R} = \gamma I$. So

$$\dot{\xi} = \frac{\beta \langle K \rangle \dot{R}}{N\gamma}$$

- ▶ And so

$$\xi = \frac{\beta \langle K \rangle}{\gamma} \frac{R - R(0)}{N} = \mathcal{R}_0 \frac{R - R(0)}{N}$$

or

$$R = \frac{N\xi}{\mathcal{R}_0} + R(0)$$

Deriving alternate equations

- ▶ So $Se^\xi = S(0)$ and thus $S(t) = S(0)e^{-\xi}$.
- ▶ We have $\dot{\xi} = \beta \langle K \rangle I/N$ and $\dot{R} = \gamma I$. So

$$\dot{\xi} = \frac{\beta \langle K \rangle \dot{R}}{N\gamma}$$

- ▶ And so

$$\xi = \frac{\beta \langle K \rangle}{\gamma} \frac{R - R(0)}{N} = \mathcal{R}_0 \frac{R - R(0)}{N}$$

or

$$R = \frac{N\xi}{\mathcal{R}_0} + R(0)$$



- ▶ Finally

$$I = N - R - S = N - \frac{N\xi}{\mathcal{R}_0} - R(0) - S(0)e^{-\xi}$$

SIR final size

- Our full equations are thus

$$\dot{\xi} = \beta \langle K \rangle \frac{I}{N} = \beta \langle K \rangle \left(1 - \frac{\xi}{\mathcal{R}_0} - \frac{R(0)}{N} - \frac{S(0)e^{-\xi}}{N} \right)$$

combined with

$$S = S(0)e^{-\xi}, \quad R = \frac{N\xi}{\mathcal{R}_0} + R(0), \quad I = N - S - R$$

SIR final size

- ▶ Our full equations are thus

$$\dot{\xi} = \beta \langle K \rangle \frac{I}{N} = \beta \langle K \rangle \left(1 - \frac{\xi}{\mathcal{R}_0} - \frac{R(0)}{N} - \frac{S(0)e^{-\xi}}{N} \right)$$

combined with

$$S = S(0)e^{-\xi}, \quad R = \frac{N\xi}{\mathcal{R}_0} + R(0), \quad I = N - S - R$$

- ▶ So we find a single independent governing equation (for ξ) that depends only on parameters, initial conditions, and ξ .

SIR final size

- ▶ Our full equations are thus

$$\dot{\xi} = \beta \langle K \rangle \frac{I}{N} = \beta \langle K \rangle \left(1 - \frac{\xi}{\mathcal{R}_0} - \frac{R(0)}{N} - \frac{S(0)e^{-\xi}}{N} \right)$$

combined with

$$S = S(0)e^{-\xi}, \quad R = \frac{N\xi}{\mathcal{R}_0} + R(0), \quad I = N - S - R$$

- ▶ So we find a single independent governing equation (for ξ) that depends only on parameters, initial conditions, and ξ .
- ▶ The final size is found using $I(\infty) = 0$, so

$$S(\infty) = N - R(\infty)$$

SIR final size

- ▶ Our full equations are thus

$$\dot{\xi} = \beta \langle K \rangle \frac{I}{N} = \beta \langle K \rangle \left(1 - \frac{\xi}{\mathcal{R}_0} - \frac{R(0)}{N} - \frac{S(0)e^{-\xi}}{N} \right)$$

combined with

$$S = S(0)e^{-\xi}, \quad R = \frac{N\xi}{\mathcal{R}_0} + R(0), \quad I = N - S - R$$

- ▶ So we find a single independent governing equation (for ξ) that depends only on parameters, initial conditions, and ξ .
- ▶ The final size is found using $I(\infty) = 0$, so

$$S(\infty) = N - R(\infty)$$

- ▶ Assuming $S(0) \approx N$ we conclude

$$Ne^{-\xi(\infty)} = N - R(\infty)$$

SIR final size

- ▶ Our full equations are thus

$$\dot{\xi} = \beta \langle K \rangle \frac{I}{N} = \beta \langle K \rangle \left(1 - \frac{\xi}{\mathcal{R}_0} - \frac{R(0)}{N} - \frac{S(0)e^{-\xi}}{N} \right)$$

combined with

$$S = S(0)e^{-\xi}, \quad R = \frac{N\xi}{\mathcal{R}_0} + R(0), \quad I = N - S - R$$

- ▶ So we find a single independent governing equation (for ξ) that depends only on parameters, initial conditions, and ξ .
- ▶ The final size is found using $I(\infty) = 0$, so

$$S(\infty) = N - R(\infty)$$

- ▶ Assuming $S(0) \approx N$ we conclude

$$Ne^{-\xi(\infty)} = N - R(\infty)$$

- ▶ Assuming $R(0) \approx 0$ and rearranging gives


$$R(\infty)/N = 1 - e^{-\mathcal{R}_0 R(\infty)/N}$$

So the final fraction $r(\infty) = R(\infty)/N$ satisfies

$$r(\infty) = 1 - e^{-\mathcal{R}_0 r(\infty)}$$

So the final fraction $r(\infty) = R(\infty)/N$ satisfies


$$r(\infty) = 1 - e^{-R_0 r(\infty)}$$



We can solve this iteratively, starting from a guess $r = 1$.

So the final fraction $r(\infty) = R(\infty)/N$ satisfies

$$r(\infty) = 1 - e^{-R_0 r(\infty)}$$



The results are in good agreement with simulation.

Direct derivation of alternate equations

Any time you derive an expression for some quantity, you should think about what it means. — Peter Saeta (my freshman physics professor)

Direct derivation of alternate equations

Any time you derive an expression for some quantity, you should think about what it means. — Peter Saeta (my freshman physics professor)

We will give a direct derivation of these new equations, and later use this approach to derive SIR equations for diseases in networks.

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

- ▶ The transmission rate per infected individual is $\beta \langle K \rangle$,

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

- ▶ The transmission rate per infected individual is $\beta \langle K \rangle$,
 - ▶ The population-wide transmission rate is $\beta \langle K \rangle I$.

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

- ▶ The transmission rate per infected individual is $\beta \langle K \rangle$,
 - ▶ The population-wide transmission rate is $\beta \langle K \rangle I$.
 - ▶ Of these, a fraction $1/N$ are expected to go to any given individual.

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

- ▶ The transmission rate per infected individual is $\beta \langle K \rangle$,
 - ▶ The population-wide transmission rate is $\beta \langle K \rangle I$.
 - ▶ Of these, a fraction $1/N$ are expected to go to any given individual.
- ▶ So

$$\frac{d}{dt} \hat{\xi} = \frac{\beta \langle K \rangle}{N} I$$

Derivation

- ▶ Define $\hat{\xi}$: the expected number of transmissions a random individual has received by time t .
- ▶ Then

$$S(t) = S(0)e^{-\hat{\xi}(t)}$$

is the probability an individual was initially susceptible and has not received any further transmissions.

- ▶ The transmission rate per infected individual is $\beta \langle K \rangle$,
 - ▶ The population-wide transmission rate is $\beta \langle K \rangle I$.
 - ▶ Of these, a fraction $1/N$ are expected to go to any given individual.
- ▶ So

$$\frac{d}{dt} \hat{\xi} = \frac{\beta \langle K \rangle}{N} I$$

- ▶ Thus $\hat{\xi}$ satisfies the same relations as ξ , and we can conclude that $\hat{\xi} = \xi$.

Introduction to Compartmental Models

Dynamics

\mathcal{R}_0

Epidemic Probability

Epidemic size

References

References |

- [1] Anne Schneeberger, Catherine H Mercer, Simon AJ Gregson, Neil M Ferguson, Constance A Nyamukapa, Roy M Anderson, Anne M Johnson, and Geoff P Garnett.
Scale-free networks and sexually transmitted diseases: a description of observed patterns of sexual contacts in britain and zimbabwe.
Sexually transmitted diseases, 31(6):380–387, 2004.
- [2] Joel C. Miller and Erik M. Volz.
Model hierarchies in edge-based compartmental modeling for infectious disease spread.
Journal of Mathematical Biology, 67(4):869–899, 2013.
- [3] Joel C. Miller.
Spread of infectious disease through clustered populations.
Journal of the Royal Society Interface, 6(41):1121, 2009.