

2.10. Geometrical optics: Optical instruments



TEASER?

- ANSWER FRAGMENT

... dummy ...

geometrical optics: optical instruments

- using reflection and refraction to study lenses and optical instruments
 - covers simple devices like magnifying glasses to complex ones like telescopes and the human eye
 - introduces key concepts such as image formation, aberrations, and optical limitations
-

primer on lenses & the lensmaker's equation

- lenses have two surfaces which can be planar, convex or concave
- convex lenses are converging while concave lenses are diverging
- when a ray enters air-to-lens and then lens-to-air, it is refracted according to snell's law
- focal length is the same on both sides (assuming a thin lens)

lensmaker's equation

$$\frac{1}{f} = (n - 1) \left(\frac{1}{R_1} + \frac{1}{R_2} \right)$$

- n is the index of refraction
 - R_1 and R_2 are the radii of curvature (positive for convex, negative for concave)
-

thin lenses model

- approximates the lens as having negligible thickness
- assumes paraxial rays (small angle

approximations:

$\sin(\theta)$

$\approx \theta$

$\cos(\theta)$

≈ 1

,

$$\frac{\tan(\theta)}{\cos(\theta)} \approx 1$$

- uses a single principal plane instead of two separate refraction planes
- distinguishes between real and virtual focal points

image formation at thin lenses via ray tracing

- three principal rays determine the image location:
 - parallel ray: travels parallel to the principal axis
 - focal point ray: passes through the focal point
 - central (optical center) ray: goes through the lens center with little

deviation

- for convex lenses:
 - parallel ray refracts through the focal point on the opposite side
 - for concave lenses:
 - refracted rays appear to diverge from the focal point on the same side as the object
-

thin lens equation & lateral magnification

- object distance d_o and image distance d_i are measured along the principal axis
- for convex lenses, the thin lens equation is:

$$\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f}$$

- for concave lenses, the equation becomes:

$$\frac{1}{d_o} - \frac{1}{d_i} = \frac{1}{f}$$

- lateral magnification is defined as:

$$m = \frac{h_i}{h_o} = -\frac{d_i}{d_o}$$

- sign conventions:
 - h_o always positive
 - h_i positive if upright, negative if inverted
 - d_o positive for real objects
 - d_i positive for real images and negative for virtual images
 - f positive for convex and negative for concave lenses
-

combining lenses

- analyze the first lens using the thin lens equation:
 - calculate d_{i1} and magnification $m_1 = -\frac{d_{i1}}{d_{o1}}$
- treat the image from the first lens as the object for the second lens

- determine the second object distance d_{o2} based on the lens separation and image location
- apply the thin lens equation for the second lens:
 - calculate d_{i2} and magnification $m_2 = -\frac{d_{i2}}{d_{o2}}$
- total magnification is:

$$m_{eff} = m_1 \times m_2$$

- careful use of sign conventions is crucial
-

pinhole cameras

- a lensless camera where a tiny pinhole focuses light onto a light-sensitive surface
- image formation is based on similar triangles:

$$\frac{h_i}{d_i} = \frac{h_o}{d_o}$$

- optimal pinhole diameter is given by:

$$\frac{d_{pin}}{\lambda f} \approx \sqrt{2.44}$$

$\sqrt{2.44}$

λf

- λ is the wavelength of light
 - f is the distance from the pinhole to the image plane
 - advantages:
 - simple, inexpensive, infinite depth of field
 - disadvantages:
 - low light sensitivity and sharpness limited by diffraction
-

the human eye

- functions similarly to a camera with additional biological sophistication
- key components:
 - cornea ($n \approx 1.376$): primary site of light bending

- iris and pupil: control light entry
- lens (
 $n_{\text{lens}} \approx 1.385$
 $n_{\text{aqueous}} \approx 1.405$
): adjusts shape (accommodation) to focus light
- retina and fovea: convert light into neural signals; fovea is the sharpest point
- aqueous humor (
 $n_{\text{aqueous}} \approx 1.336$
) and vitreous humor (
 $n_{\text{vitreous}} \approx 1.337$
): transparent media filling the eye

- the eye has a fixed image distance
- corrective lenses are used to adjust the focus for nearsightedness, farsightedness, and astigmatism

corrective lenses

- nearsightedness (myopia):
 - image forms in front of the retina

- corrected using diverging (negative) lenses
 - farsightedness (hyperopia):
 - image forms behind the retina
 - corrected using converging (positive) lenses
 - astigmatism:
 - irregular cornea or lens shape causes blurred images
 - corrected with cylindrical lenses
-

magnifying glass

- a simple magnifier that uses a single converging lens to enlarge the apparent size of objects
- the apparent size is based on the angular magnification

$$M = \frac{\theta'}{\theta}$$
- with a relaxed eye (image at infinity):
 - object is placed at the focal point f
 - angular magnification is:

$$M = \frac{N}{f}$$

- where
 $N = 25$

,

text cm

is the near point

- when the eye is focused at the near point:
 - the virtual image is formed at
 $d_i = -N$
 - using the thin lens equation:

$$\frac{1}{f} = \frac{1}{d_o} - \frac{1}{N}$$

- the object distance becomes:

$$d_o = \frac{Nf}{N + f}$$

- angular magnification is then:

$$M = 1 + \frac{N}{f}$$

- a larger subtended angle leads to a larger apparent size

Telescopes

introduction

- telescopes are optical instruments designed to magnify distant objects
- they allow observation of celestial bodies and faraway phenomena in greater detail

components

- objective lens/mirror collects light from the object and forms a real image
- eyepiece lens magnifies the image formed by the objective

angular magnification

- the total angular magnification M of a telescope is defined as the ratio of the angle subtended by the image at the eye to the angle subtended by the object as seen by the unaided eye

- for an objective with focal length f_o and an eyepiece with focal length f_e , the magnification is given by

$$M = \frac{\theta'}{\theta} = \frac{h/f_e}{h/f_o} = \frac{f_o}{f_e}$$

- with the small angle approximation $\theta' \approx h/f_e$ and $\theta \approx h/f_o$

image formation

- the image of the first lens (objective) becomes the object for the second lens (eyepiece)
- objective lens forms a real, inverted image near its focal point
- eyepiece lens acts as a magnifying glass producing a virtual, inverted final image

types of telescopes

REFRACTING TELESCOPES (KEPLERIAN)

- design uses two converging lenses
- objective lens has a long focal length to collect light and form a real, inverted, diminished image at its focal plane

- eyepiece lens has a short focal length and magnifies the real image
- advantages include simple design and good image quality for small apertures
- disadvantages include heavy, expensive objective lenses and chromatic aberration

REFLECTING TELESCOPES (NEWTONIAN)

- design uses a concave mirror as the objective to collect and focus light
- light reflects off the primary mirror forming a real, inverted image at the focal point
- a secondary flat mirror reflects the image to the side for viewing through an eyepiece
- advantages include absence of chromatic aberration and easier manufacture of large mirrors
- disadvantages include diffraction effects from the secondary mirror and the need for periodic realignment

Aberration

introduction

- aberration refers to imperfections in image formation due to the breakdown of the small angle approximation
- real lenses have thickness and imperfections that can lead to blurring or distortion

types of aberrations

- monochromatic aberrations occur with a single wavelength and include spherical aberration, coma, astigmatism, curvature of field, and distortion
- chromatic aberration arises from dispersion where different wavelengths are refracted by different amounts

spherical aberration

- occurs when rays far from the optical axis focus at different points than rays close to the axis
- causes a blurred image or a circle of least confusion

- correction methods include aspherical lenses, aperture stops, or compound lenses

coma

- causes off-axis points to be imaged as asymmetrical, comet-shaped blurs
- results from off-axis rays being refracted by different amounts
- can be minimized using aspherical or carefully designed compound lenses

astigmatism

- happens when a lens or mirror has different focal lengths in two perpendicular planes
- a point object is imaged as two short perpendicular line segments
- correction involves using a combination of lenses or cylindrical lenses

curvature of field

- the image of a flat object is formed on a curved surface instead of a flat plane
- results in a sharp center with blurred edges or vice versa

- field-flattening lenses or specific lens designs can correct this aberration

distortion

- causes the shape of the image to differ from the shape of the object
- barrel distortion makes straight lines bow outwards and pincushion distortion makes them bow inwards
- symmetric compound lenses or software correction can reduce distortion

chromatic aberration

- arises because the refractive index of a lens varies with the wavelength of light
- results in colored fringes especially at high contrast edges
- achromatic doublets or apochromatic lenses are used for correction

aberrations in mirrors

- spherical aberration in mirrors occurs for the same geometric reasons as in lenses

- parabolic mirrors can correct spherical aberration for objects at infinity
- mirrors do not exhibit chromatic aberration because reflection is independent of wavelength

summary of aberrations

- spherical aberration: caused by the spherical shape of the lens/mirror; corrected by aspherical lenses, aperture stops, or compound lenses
- coma: off-axis rays cause a comet-shaped blur; minimized with aspherical or compound lenses
- astigmatism: different focal lengths in perpendicular planes; corrected by lens combinations or cylindrical lenses
- curvature of field: focal points on a curved surface lead to uneven focus; corrected with field-flattening lenses
- distortion: non-uniform magnification leads to curved lines; minimized with symmetric compound lenses or digital correction

- chromatic aberration: dispersion causes colored fringes; corrected with achromatic doublets or apochromatic lenses
-