


Specification, Implementation, and Complexity of Replicated Data Types with Composite Operations

Hengfeng Wei

State Key Laboratory for Novel Software Technology, Nanjing University, China


hfwei@nju.edu.cn

 <https://orcid.org/0000-0002-0427-9710>

Huang Yu

State Key Laboratory for Novel Software Technology, Nanjing University, China

yuhuang@nju.edu.cn

 <https://orcid.org/0000-0001-8921-036X>

Jian Lu

State Key Laboratory for Novel Software Technology, Nanjing University, China

lj@nju.edu.cn

Abstract

2012 ACM Subject Classification

Keywords and phrases Replicated set

Digital Object Identifier 10.4230/LIPIcs...

1 Introduction

2 Preliminaries

2.1 Observed-Remove Set (OR-Set)

$$\mathcal{F}_{\text{orset}}(\text{rd}, E, \text{op}, \text{vis}, \text{ar}) = \{a \mid \exists e \in E. \text{op}(e) = \text{add}(a) \quad (1)$$

$$\wedge (\forall f \in E. \text{op}(f) = \text{rm}(a) \implies \neg(e \xrightarrow{\text{vis}} f))\}. \quad (2)$$

3 Replicated Data Types with Composite Operations

3.1 Specification

We consider a composite operation of a replicated data type τ in the form of $C = A \oplus B$, where A , B , and C are different objects of type τ .

Following [?], we specify the semantics of a composite operation $A \oplus B$ of a replicated data type τ by a function \mathcal{F}_τ that determines the return value of \oplus based on prior operations performed on the two objects involved (i.e., A and B). However, \mathcal{F}_τ for a composite operation \oplus takes as parameters two, not one as in [?], *operation contexts*, one on each object involved.

Q : Generalize to different data types?

Note: Partial operation context [?]



© Hengfeng Wei, Yu Huang, and Jian Lu;

licensed under Creative Commons License CC-BY

Leibniz International Proceedings in Informatics

LIPICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

► **Definition 1** (Product of Operation Contexts). Consider two operation contexts for the same replicated data type τ :

$$\mathcal{C}_A = (E_A, \text{op}_A, \text{vis}_A, \text{ar}_A) \quad (3)$$

$$\mathcal{C}_B = (E_B, \text{op}_B, \text{vis}_B, \text{ar}_B) \quad (4)$$

The product $\mathcal{C} = \mathcal{C}_A \times \mathcal{C}_B$ of \mathcal{C}_A and \mathcal{C}_B is also an operation context defined as $\mathcal{C} = (E, \text{op}, \text{vis}, \text{ar})$, where

$$\text{--- } E = E_A \times E_B$$

$$\text{--- } \text{op} = \text{op}_A \sqcup \text{op}_B$$

$$\text{--- } \text{vis} = \text{vis}_A \times \text{vis}_B$$

$$\text{--- } \text{ar} = \text{ar}_A \times \text{ar}_B$$

► **Definition 2.**

$$\mathcal{F}_\tau(\oplus, \mathcal{C}_A, \mathcal{C}_B) = \mathcal{F}_\tau(\oplus, \mathcal{C}_A \times \mathcal{C}_B) \quad (5)$$

4 Replicated Counter with Composite Operations

We consider the replicated counter data type with composite operations including addition (+) and subtraction (−).

5 Replicated Set with Composite Operations

We consider the replicated set data type with composite operations including union (\cup), intersection (\cap), difference (\setminus), and symmetric difference (\otimes).

5.1 Specification

General method for set union, set intersection, and set difference: There are four ...

$$\mathcal{F}_{\text{orset}}(A \setminus B, \mathcal{C}_A, \mathcal{C}_B) \quad (6)$$

$$= \{a \mid \exists (e, e') \in E_A \times E_B. ((\text{op}(e) = \text{add}(a) \wedge \text{op}(e') = \text{rm}(a)) \quad (7)$$

$$\wedge \forall (f, f') \in E_A \times E_B. ((\text{op}(f) = \text{add}(a) \wedge \text{op}(f') = \text{add}(a)) \vee (\text{op}(f) = \text{rm}(a) \wedge \text{op}(f') = \text{add}(a)) \quad (8)$$

$$\implies \neg((e, e') \xrightarrow{\text{vis}} (f, f'))\}. \quad (9)$$

► **Example 3** (Specification for Composite Operations of Replicated Set).

5.2 Protocol

5.2.1 Methodology

Op-based

We always assume that $A \neq \emptyset$, $A' \neq \emptyset$, and $(\mathbb{A} \cup \mathbb{R}) \cap (\mathbb{A}' \cup \mathbb{R}') = \emptyset$. Note that for any replica state $(a, \mathbb{A}, \mathbb{R})$, we have $\mathbb{R} \subseteq \mathbb{A}$.

$Q : \emptyset$ issues The goal is to find \square and \diamond such that

$$\dots \iff \square \subseteq \diamond$$

U : universe of elements
 I : set of identifiers
 $\mathbb{A}, \mathbb{R} \in \text{SUBSET } \mathcal{I}$
 $\Sigma = \mathcal{U} \times \mathbb{A} \times \mathbb{R}$
 $\sigma_0 = \lambda r. \emptyset$

■ **Figure 1** Implementation of replicated set with composite operations.

5.2.2 Set Union (\cup)

The goal is to find \square and \diamond such that

$$(\mathbb{R} \subseteq \mathbb{A}) \vee (\mathbb{R}' \subseteq \mathbb{A}') \iff \square \subseteq \diamond.$$

► **Theorem 4** (Set Union). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

$$(\mathbb{R} \subseteq \mathbb{A}) \vee (\mathbb{R}' \subseteq \mathbb{A}') \iff (\mathbb{R} \cup \mathbb{R}') \subseteq (\mathbb{A} \cup \mathbb{A}').$$

► **Theorem 5** (Set Union (Alternative)). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

$$(\mathbb{R} \subseteq \mathbb{A}) \vee (\mathbb{R}' \subseteq \mathbb{A}') \iff (\mathbb{A} \times \mathbb{R}') \cap (\mathbb{R} \times \mathbb{A}') \subseteq (\mathbb{A} \times \mathbb{A}')$$

Q : How to choose? Another way:

5.2.3 Set Intersection (\cap)

The goal is to find \square and \diamond such that

$$(\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \subseteq \mathbb{A}') \iff \square \subseteq \diamond.$$

► **Theorem 6** (Set Intersection). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

$$(\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \subseteq \mathbb{A}') \iff (\mathbb{A} \times \mathbb{R}') \cup (\mathbb{R} \times \mathbb{A}') \subseteq (\mathbb{A} \times \mathbb{A}').$$

$$(\mathbb{R} \subseteq \mathbb{A}) \vee (\mathbb{R}' \subseteq \mathbb{A}')$$

Another way:
IF THEN ELSE

5.2.4 Set Difference (\setminus)

The goal is to find \square and \diamond such that

$$(\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \not\subseteq \mathbb{A}') \iff \square \subseteq \diamond.$$

► **Theorem 7** (Set Difference). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

$$(\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \not\subseteq \mathbb{A}') \iff (\mathbb{R} \times \mathbb{A}') \subseteq (\mathbb{A} \times \mathbb{R}').$$

5.2.5 Set Symmetric Difference (\otimes)

The goal is to find \square and \diamond such that

$$((\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \not\subseteq \mathbb{A}')) \vee ((\mathbb{R} \not\subseteq \mathbb{A}) \wedge (\mathbb{R}' \subseteq \mathbb{A}')) \iff \square \subseteq \diamond.$$

By the identity $A \otimes B = (A \cup B) \setminus (A \cap B)$ and Theorems 4, 6, and 7, we have

► **Theorem 8** (Set Symmetric Difference). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

$$\begin{aligned}
 & ((\mathbb{R} \subseteq \mathbb{A}) \wedge (\mathbb{R}' \not\subseteq \mathbb{A}')) \vee ((\mathbb{R} \not\subseteq \mathbb{A}) \wedge (\mathbb{R}' \subseteq \mathbb{A}')) \\
 & \iff (\mathbb{R} \cup \mathbb{R}') \times (\mathbb{A} \cup \mathbb{A}') \subseteq (\mathbb{A} \cup \mathbb{A}') \times ((\mathbb{A} \times \mathbb{R}') \cup (\mathbb{R} \times \mathbb{A}'))
 \end{aligned}$$

It fails to use
Theorem 4 here.

By the identity $A \otimes B = (A \setminus B) \cup (B \setminus A)$ and Theorems 5 and 7, we have

► **Theorem 9** (Set Symmetric Difference (Alternative)). *Given that $\mathbb{R} \subseteq \mathbb{A}$ and $\mathbb{R}' \subseteq \mathbb{A}'$, we have*

94

95

96

97

98

$$\begin{aligned} & ((\mathbb{R} \subset \mathbb{A}) \wedge (\mathbb{R}' \not\subset \mathbb{A}')) \vee ((\mathbb{R} \not\subset \mathbb{A}) \wedge (\mathbb{R}' \subset \mathbb{A}')) \\ & \iff (\mathbb{A} \times \mathbb{R}') \times (\mathbb{R}' \times \mathbb{A}) \cap (\mathbb{R} \times \mathbb{A}') \times (\mathbb{A}' \times \mathbb{R}) \subset (\mathbb{A} \times \mathbb{R}') \times (\mathbb{A}' \times \mathbb{R}) \end{aligned}$$

99

6 Related Work

100

[?] [?]

101

7 Conclusion and Future Work

102

References
