


Universal Virasoro Constraints for Additive Categories

Arkadij Bojko

Institute of Mathematics, Academia Sinica

August 24, 2024



Structure of the talk.

1. History and background
2. Geometric formulation of Virasoro constraints and the main claim
3. Reformulation in terms of vertex algebras
 - 3.1 Joyce's construction of VA's
 - 3.2 The conformal element
 - 3.3 Virasoro constraints make virtual fundamental classes into physical states
4. Main results for quivers and varieties

Witten's conjecture

1. The moduli space of algebraic pointed curves $\overline{\mathcal{M}}_{g,n}$ parametrizing semistable¹ (C, x_1, \dots, x_n) :




Figure: (C, x_1, \dots, x_n) where C has genus 7 here, $n = 11$, and x_1, \dots, x_{11} are represented by the black dots.

¹This means that each non-singular curve in the picture satisfies $2 - 2g - n < 0$ where n is the number of marked points and ordinary double points it shares.

Witten's conjecture

1. The moduli space of algebraic pointed curves $\overline{\mathcal{M}}_{g,n}$ parametrizing semistable¹ (C, x_1, \dots, x_n) :




Figure: (C, x_1, \dots, x_n) where C has genus 7 here, $n = 11$, and x_1, \dots, x_{11} are represented by the black dots.

2. There are line bundles $L_i \rightarrow \overline{\mathcal{M}}_{g,n}$ which at each C are given by $T_C^*|_{x_i}$:




Figure: $L_i|_{(C, x_1, \dots, x_n)} = T_C^*|_{x_i}$

¹This means that each non-singular curve in the picture satisfies $2 - 2g - n < 0$ where n is the number of marked points and ordinary double points it shares.

Witten's conjecture

1. The moduli space of algebraic pointed curves $\overline{\mathcal{M}}_{g,n}$ parametrizing semistable¹ (C, x_1, \dots, x_n) :




Figure: (C, x_1, \dots, x_n) where C has genus 7 here, $n = 11$, and x_1, \dots, x_{11} are represented by the black dots.

2. There are line bundles $L_i \rightarrow \overline{\mathcal{M}}_{g,n}$ which at each C are given by $T_C^*|_{x_i}$:




Figure: $L_i|_{(C, x_1, \dots, x_n)} = T_C^*|_{x_i}$

3. Denote the powers of the first Chern classes by $\tau_d := \psi_i^d := c_1(L_i)^d$

¹This means that each non-singular curve in the picture satisfies $2 - 2g - n < 0$ where n is the number of marked points and ordinary double points it shares.

Witten's conjecture II

1. Define the potential

$$F(\vec{t}) = \sum_{g \geq 0} \left\langle \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \right\rangle_g \lambda^{2g-2} = \int_{\sum_{g,n \geq 0} [\overline{\mathcal{M}}_{g,n}]} \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \lambda^{2g-2}$$

which collects the invariants

$$\langle \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k} \rangle_{g,n} = \int_{\overline{\mathcal{M}}_{g,n}} \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k}$$

with $\sum_i a_i = n$.

3. These constraints determine τ and F uniquely.

Witten's conjecture II

1. Define the potential

$$F(\vec{t}) = \sum_{g \geq 0} \left\langle \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \right\rangle_g \lambda^{2g-2} = \int_{\sum_{g,n \geq 0} [\overline{\mathcal{M}}_{g,n}]} \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \lambda^{2g-2}$$

which collects the invariants

$$\langle \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k} \rangle_{g,n} = \int_{\overline{\mathcal{M}}_{g,n}} \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k}$$

with $\sum_i a_i = n$.

3. These constraints determine τ and F uniquely.

Witten's conjecture II

1. Define the **potential**

$$F(\vec{t}) = \sum_{g \geq 0} \left\langle \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \right\rangle_g \lambda^{2g-2} = \int_{\sum_{g,n \geq 0} [\mathcal{M}_{g,n}]} \exp \left[\sum_{d \geq 0} \tau_d t_d \right] \lambda^{2g-2}$$

which collects the invariants

$$\langle \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k} \rangle_{g,n} = \int_{\mathcal{M}_{g,n}} \tau_0^{a_0} \tau_1^{a_1} \cdots \tau_k^{a_k}$$

with $\sum_i a_i = n$.

2. The interpretation of **Witten's conjecture** by **Dijkgraaf, Verlinde, Verlinde (90')** states that the τ -function

$$\tau = \exp \left[-\frac{1}{2} F(x, \vec{t}) \right]$$

satisfies **Virasoro constraints**.

3. These constraints determine τ and F **uniquely**.

Virasoro constraints for GW

1. Dijkgraaf, Verlinde, and Verlinde (90') defined a sequence of second order differential operators L_k on $\mathbb{C}[[t_0, t_1, \dots, t_n, \dots]]$ by

$$L_k = T_k + R_k,$$

where

Virasoro constraints for GW

1. Dijkgraaf, Verlinde, and Verlinde (90') defined a sequence of second order differential operators L_k on $\mathbb{C}[[t_0, t_1, \dots, t_n, \dots]]$ by

$$L_k = T_k + R_k,$$

where

$$T_k = \frac{\lambda^2}{2} \sum_{m=1}^k \frac{\partial^2}{\partial t_{m-1} \partial t_{k-m}}, \quad \text{for } k \geq 1,$$
$$R_k = \sum_{m=0}^{\infty} \left(m + \frac{1}{2}\right) t_m \frac{\partial}{\partial t_{m+k}}, \quad \text{for } k \geq -1$$

and $T_0 = \frac{1}{16}$, $T_{-1} = \frac{1}{8} \lambda^{-2} t_0^2$.

Virasoro constraints for GW

1. Dijkgraaf, Verlinde, and Verlinde (90') defined a sequence of second order differential operators L_k on $\mathbb{C}[[t_0, t_1, \dots, t_n, \dots]]$ by

$$L_k = T_k + R_k,$$

where

$$T_k = \frac{\lambda^2}{2} \sum_{m=1}^k \frac{\partial^2}{\partial t_{m-1} \partial t_{k-m}}, \quad \text{for } k \geq 1,$$
$$R_k = \sum_{m=0}^{\infty} \left(m + \frac{1}{2}\right) t_m \frac{\partial}{\partial t_{m+k}}, \quad \text{for } k \geq -1$$

and $T_0 = \frac{1}{16}$, $T_{-1} = \frac{1}{8} \lambda^{-2} t_0^2$.

2. These satisfy the **Virasoro commutation relations**

$$[L_m, L_n] = (m - n) L_{m+n}.$$

Virasoro constraints for GW

1. Dijkgraaf, Verlinde, and Verlinde (90') defined a sequence of second order differential operators L_k on $\mathbb{C}[[t_0, t_1, \dots, t_n, \dots]]$ by

$$L_k = T_k + R_k,$$

where

$$T_k = \frac{\lambda^2}{2} \sum_{m=1}^k \frac{\partial^2}{\partial t_{m-1} \partial t_{k-m}}, \quad \text{for } k \geq 1,$$
$$R_k = \sum_{m=0}^{\infty} \left(m + \frac{1}{2}\right) t_m \frac{\partial}{\partial t_{m+k}}, \quad \text{for } k \geq -1$$

and $T_0 = \frac{1}{16}$, $T_{-1} = \frac{1}{8} \lambda^{-2} t_0^2$.

2. These satisfy the **Virasoro commutation relations**

$$[L_m, L_n] = (m - n) L_{m+n}.$$

3. DVV showed that **Witten's conjecture** is equivalent to the **Virasoro constraints**

$$L_k \tau = 0, \quad \text{for } k \geq -1$$

Proof of Witten's conjecture

1. Witten's conjecture was proved famously by Kontsevich (92').

Proof of Witten's conjecture

1. Witten's conjecture was proved famously by Kontsevich (92').
2. Later it was proved differently by Okounkov–Pandharipande(01') and another approach was given by Mirzakhani (07').

Proof of Witten's conjecture

1. **Witten's conjecture** was proved famously by Kontsevich (92').
2. Later it was proved differently by Okounkov–Pandharipande(01') and another approach was given by Mirzakhani (07').



Figure: Kontsevich, Okounkov and Mirzakhani have received Fields medals in part for their work on Witten's conjecture.

Virasoro constraints for $X \neq \text{pt}$

1. Let us now move on to $\overline{\mathcal{M}}_{g,n}(X, \beta)$ where X is a smooth projective variety and $\beta \in H_2(X)$ an algebraic curve class. It parametrizes **stable maps** (C, f, x_1, \dots, x_n) with $f : C \rightarrow X$ such that $f_*([C]) = \beta \in H_2(X)$.

²Here ev_j is the evaluation map for the j 'th marked point.

Virasoro constraints for $X \neq \text{pt}$

1. Let us now move on to $\overline{\mathcal{M}}_{g,n}(X, \beta)$ where X is a smooth projective variety and $\beta \in H_2(X)$ an algebraic curve class. It parametrizes **stable maps** (C, f, x_1, \dots, x_n) with $f : C \rightarrow X$ such that $f_*([C]) = \beta \in H_2(X)$.
2. Fix a basis $B = \{v\} \subset H^*(X)$ with $1 \in B$ for the generator of $H^0(X)$ and define the classes ²

$$\tau_k(v) = (\psi_j)^k \text{ev}_j^*(v), \quad \text{for all } k \geq 0, v \in B$$

²Here ev_j is the evaluation map for the j 'th marked point.

Virasoro constraints for $X \neq \text{pt}$

1. Let us now move on to $\overline{\mathcal{M}}_{g,n}(X, \beta)$ where X is a smooth projective variety and $\beta \in H_2(X)$ an algebraic curve class. It parametrizes **stable maps** (C, f, x_1, \dots, x_n) with $f : C \rightarrow X$ such that $f_*([C]) = \beta \in H_2(X)$.
2. Fix a basis $B = \{\nu\} \subset H^*(X)$ with $1 \in B$ for the generator of $H^0(X)$ and define the classes ²

$$\tau_k(\nu) = (\psi_j)^k \text{ev}_j^*(\nu), \quad \text{for all } k \geq 0, \nu \in B$$

3. Consider the **Gromov–Witten potential**

$$F^X(\vec{t}) = \sum_{\substack{g \geq 0 \\ \beta \in H_2(X)}} \left\langle \exp \left[\sum_{\substack{k \geq 0 \\ \nu \in B}} \tau_k(\nu) t_{k,\nu} \right] \right\rangle_{\beta, g}^X q^\beta \lambda^{2g-2}$$

²Here ev_j is the evaluation map for the j 'th marked point.

Virasoro constraints for $X \neq \text{pt}$

1. Let us now move on to $\overline{\mathcal{M}}_{g,n}(X, \beta)$ where X is a smooth projective variety and $\beta \in H_2(X)$ an algebraic curve class. It parametrizes **stable maps** (C, f, x_1, \dots, x_n) with $f : C \rightarrow X$ such that $f_*([C]) = \beta \in H_2(X)$.
2. Fix a basis $B = \{v\} \subset H^*(X)$ with $1 \in B$ for the generator of $H^0(X)$ and define the classes ²

$$\tau_k(v) = (\psi_j)^k \text{ev}_j^*(v), \quad \text{for all } k \geq 0, v \in B$$

3. Consider the **Gromov–Witten potential**

$$F^X(\vec{t}) = \sum_{\substack{g \geq 0 \\ \beta \in H_2(X)}} \left\langle \exp \left[\sum_{\substack{k \geq 0 \\ v \in B}} \tau_k(v) t_{k,v} \right] \right\rangle_{\beta, g}^X q^\beta \lambda^{2g-2}$$

which collects the **Gromov–Witten invariants**

$$\langle \tau_{k_1}(v_1)^{a_1} \tau_{k_2}(v_2)^{a_2} \cdots \tau_{k_l}(v_l)^{a_l} \rangle_{\beta, g}^X = \int \tau_{k_1}(v_1)^{a_1} \tau_{k_2}(v_2)^{a_2} \cdots \tau_{k_l}(v_l)^{a_l} \left[\overline{\mathcal{M}}_{g,n}(X, \beta) \right]^{\text{vir}}$$

under the condition that $\sum_i a_i = n$.

²Here ev_j is the evaluation map for the j 'th marked point.

Virasoro constraints II

1. T. Eguchi, K. Hori, and C.-S. Xiong (97') and Katz (presented by Eguchi, Jinzenji, and Xiong) extended the operators L_k to the case of $\overline{\mathcal{M}}_{g,n}(X, \beta)$ as second order differential operators in $t_{k,v}$. If one requires that the basis B preserves the Hodge grading – that is every $v \in H^{p,q}(X)$ for some p, q , then L_k depend explicitly on the holomorphic grading p .

Virasoro constraints II

1. T. Eguchi, K. Hori, and C.-S. Xiong (97') and Katz (presented by Eguchi, Jinzenji, and Xiong) extended the operators L_k to the case of $\overline{\mathcal{M}}_{g,n}(X, \beta)$ as second order differential operators in $t_{k,v}$. If one requires that the basis B preserves the Hodge grading – that is every $v \in H^{p,q}(X)$ for some p, q , then L_k depend explicitly on the holomorphic grading p .
2. These operators are rather complicated, so I won't write them.

Virasoro constraints II

1. T. Eguchi, K. Hori, and C.-S. Xiong (97') and Katz (presented by Eguchi, Jinzenji, and Xiong) extended the operators L_k to the case of $\overline{\mathcal{M}}_{g,n}(X, \beta)$ as second order differential operators in $t_{k,v}$. If one requires that the basis B preserves the Hodge grading – that is every $v \in H^{p,q}(X)$ for some p, q , then L_k depend explicitly on the holomorphic grading p .
2. These operators are rather complicated, so I won't write them.
3. EHX conjectured that

$$L_k \exp \left[-\frac{1}{2} F^X(\vec{t}) \right] = 0, \quad k \geq -1.$$

Virasoro constraints II

1. T. Eguchi, K. Hori, and C.-S. Xiong (97') and Katz (presented by Eguchi, Jinzenji, and Xiong) extended the operators L_k to the case of $\overline{\mathcal{M}}_{g,n}(X, \beta)$ as second order differential operators in $t_{k,v}$. If one requires that the basis B preserves the Hodge grading – that is every $v \in H^{p,q}(X)$ for some p, q , then L_k depend explicitly on the holomorphic grading p .
2. These operators are rather complicated, so I won't write them.
3. EHX conjectured that

$$L_k \exp \left[-\frac{1}{2} F^X(\vec{t}) \right] = 0, \quad k \geq -1.$$

4. This was proved by Okounkov–Pandharipande (03') for curves $X = C$ and by Givental (01') and Teleman (12') for toric X (more generally X with semisimple quantum cohomology). Givental's formalism uses Kontsevich's result for $\overline{\mathcal{M}}_{g,n}$, equivariant localization, and cohomological field theories.

Virasoro constraints II

1. T. Eguchi, K. Hori, and C.-S. Xiong (97') and Katz (presented by Eguchi, Jinzenji, and Xiong) extended the operators L_k to the case of $\overline{\mathcal{M}}_{g,n}(X, \beta)$ as second order differential operators in $t_{k,v}$. If one requires that the basis B preserves the Hodge grading – that is every $v \in H^{p,q}(X)$ for some p, q , then L_k depend explicitly on the holomorphic grading p .
2. These operators are rather complicated, so I won't write them.
3. EHX conjectured that

$$L_k \exp \left[-\frac{1}{2} F^X(\vec{t}) \right] = 0, \quad k \geq -1.$$

4. This was proved by Okounkov–Pandharipande (03') for curves $X = C$ and by Givental (01') and Teleman (12') for toric X (more generally X with semisimple quantum cohomology). Givental's formalism uses Kontsevich's result for $\overline{\mathcal{M}}_{g,n}$, equivariant localization, and cohomological field theories.
5. Virasoro constraints for toric 3-folds X were transported to stationary descendants of PT stable pairs using the GW–PT correspondence by Moreira–Oblomkov–Okounkov–Pandharipande (20'). Dimensional reduction was used to prove these constraints for $\text{Hilb}^n(S)$ – a rather long round-about proof, which hides how natural the Virasoro constraints on the sheaf side are.

Virasoro constraints for sheaves and pairs

1. One thing that separates GW theory from sheaves is the **linearity** of the latter. It is manifested by working with **additive categories**. This additivity is in fact sufficient for formulating Virasoro constraints as long as we have **virtual fundamental classes** to integrate against.

³This case is not written anywhere, but it follows the same argument once my proof of wall-crossing is out.


Virasoro constraints for sheaves and pairs

1. One thing that separates GW theory from sheaves is the **linearity** of the latter. It is manifested by working with **additive categories**. This additivity is in fact sufficient for formulating Virasoro constraints as long as we have **virtual fundamental classes** to integrate against.
2. Today, I will only focus on categories of **quiver representations** and **sheaves**. Afterwards, everyone in the room should be able to conjecture Virasoro constraints for any additive category with reasonable virtual fundamental classes, and perhaps even prove them.

³This case is not written anywhere, but it follows the same argument once my proof of wall-crossing is out.

Virasoro constraints for sheaves and pairs

1. One thing that separates GW theory from sheaves is the **linearity** of the latter. It is manifested by working with **additive categories**. This additivity is in fact sufficient for formulating Virasoro constraints as long as we have **virtual fundamental classes** to integrate against.
2. Today, I will only focus on categories of **quiver representations** and **sheaves**. Afterwards, everyone in the room should be able to conjecture Virasoro constraints for any additive category with reasonable virtual fundamental classes, and perhaps even prove them.
3. Fixing a **stability condition** σ and a **class** α , e.g. a dimension vector or a Chern character, Virasoro constraints are stated for each virtual fundamental class $[M_\alpha^\sigma]^{\text{vir}}$, where M_α^σ parametrizes σ -stable objects in class α , separately. I will just write M when σ, α are not important. I will also assume that **M is fine** for now.

³This case is not written anywhere, but it follows the same argument once my proof of wall-crossing is out. 

Virasoro constraints for sheaves and pairs


1. One thing that separates GW theory from sheaves is the **linearity** of the latter. It is manifested by working with **additive categories**. This additivity is in fact sufficient for formulating Virasoro constraints as long as we have **virtual fundamental classes** to integrate against.
2. Today, I will only focus on categories of **quiver representations** and **sheaves**. Afterwards, everyone in the room should be able to conjecture Virasoro constraints for any additive category with reasonable virtual fundamental classes, and perhaps even prove them.
3. Fixing a **stability condition** σ and a **class** α , e.g. a dimension vector or a Chern character, Virasoro constraints are stated for each virtual fundamental class $[M_\alpha^\sigma]^{\text{vir}}$, where M_α^σ parametrizes σ -stable objects in class α , separately. I will just write M when σ, α are not important. I will also assume that **M is fine** for now.
4. Some **examples** to see how much variety the theory offers:

	Without framing	With framing
Sheaves	Gieseker stable torsion-free sheaves on curves or surface, dimension 1 sheaves on surfaces, Fano 3-folds, CY fourfolds ³	Bradlow pairs on curves or surfaces, DT/PT pairs on ≤ 4 -folds, Quot schemes
Quivers with relations (quasi-smooth, CY4)	Bridgeland stable quiver representations	Framed quiver representations: e.g., Grassmanians and Flag varieties

³This case is not written anywhere, but it follows the same argument once my proof of wall-crossing is out.


Quivers with relations and frozen vertices

1. The main focus will be quivers $Q = (V, E)$ with frozen vertices F and an ideal of relations \mathcal{R} generated by (minimal) R



Quivers with relations and frozen vertices


1. The main focus will be quivers $Q = (V, E)$ with frozen vertices F and an ideal of relations \mathcal{R} generated by (minimal) R



2. Quasi-smooth is equivalent to having virtual fundamental classes for the corresponding quiver moduli.

Quivers with relations and frozen vertices


1. The main focus will be quivers $Q = (V, E)$ with frozen vertices F and an ideal of relations \mathcal{R} generated by (minimal) R




2. Quasi-smooth is equivalent to having virtual fundamental classes for the corresponding quiver moduli.
3. Fix a dimension vector \bar{d} and a stability condition σ leading to the moduli space $M_{\bar{d}}^{\sigma}$ of semistable objects.

Quivers with relations and frozen vertices

1. The main focus will be quivers $Q = (V, E)$ with frozen vertices F and an ideal of relations \mathcal{R} generated by (minimal) R




2. Quasi-smooth is equivalent to having virtual fundamental classes for the corresponding quiver moduli.
3. Fix a dimension vector \vec{d} and a stability condition σ leading to the moduli space $M_{\vec{d}}^{\sigma}$ of semistable objects.
4. Examples:



Flag

$$(x_i y_k) \circ x_j = (x_j y_k) \circ x_i, \quad (x_k y_i) \circ y_j = (x_k y_j) \circ y_i$$



Virasoro constraints for quivers and sheaves

Next, I will describe the equivalent ingredients for defining Virasoro constraints on quiver representations and sheaves from left to right.

Virasoro constraints for quivers and sheaves

Next, I will describe the equivalent ingredients for defining Virasoro constraints on quiver representations and sheaves from left to right.

1. The descendent algebras

$$\mathbb{D}^Q = \text{Sym}[\tau_i(v), i > 0, v \in V]$$

$$\mathbb{D}^X = \text{SSym}[\tau_i^H(v), i > 0, v \in B]$$

$B \subset H^*(X)$ is a basis that respects the Hodge grading

The Virasoro operators $L_k = T_k + R_k$ for $k \geq -1$ will now be differential operators on $\mathbb{D}^{Q/X}$.

Virasoro constraints for quivers and sheaves

Next, I will describe the equivalent ingredients for defining Virasoro constraints on quiver representations and sheaves from left to right.

1. The descendent algebras

$$\mathbb{D}^Q = \text{Sym}[\tau_i(v), i > 0, v \in V]$$

$$\mathbb{D}^X = \text{SSym}[\tau_i^H(v), i > 0, v \in B]$$

$B \subset H^*(X)$ is a basis that respects the Hodge grading

The Virasoro operators $L_k = T_k + R_k$ for $k \geq -1$ will now be differential operators on $\mathbb{D}^{Q/X}$.

2. The realization map: depends on a choice of a universal object. We use it when integrating against $[M]^{\text{vir}}$

$$\tau_i(v) \mapsto \text{ch}_i(\mathbb{U}_v)$$

\mathbb{U}_v is a universal vector space at v

$$\tau_i^H(v) = \pi_{2,*}(\pi_1^*(\bar{v})\text{ch}_{i+p}(\mathbb{G}))$$

\mathbb{G} is a universal sheaf on $X \times M$
 $v \in H^{p,q}(X)$, \bar{v} its Poincaré dual

Virasoro constraints for quivers and sheaves

Next, I will describe the equivalent ingredients for defining Virasoro constraints on quiver representations and sheaves from left to right.

1. The descendent algebras

$$\mathbb{D}^Q = \text{Sym}[\tau_i(v), i > 0, v \in V]$$

$$\mathbb{D}^X = \text{SSym}[\tau_i^H(v), i > 0, v \in B]$$

$B \subset H^*(X)$ is a basis that respects the **Hodge grading**

The **Virasoro operators** $L_k = T_k + R_k$ for $k \geq -1$ will now be differential operators on $\mathbb{D}^{Q/X}$.

2. **The realization map:** depends on a choice of a **universal object**. We use it when integrating against $[M]^{\text{vir}}$

$$\tau_i(v) \mapsto \text{ch}_i(\mathbb{U}_v)$$

\mathbb{U}_v is a **universal vector space** at v

$$\tau_i^H(v) = \pi_{2,*}(\pi_1^*(\bar{v}) \text{ch}_{i+p}(\mathbb{G}))$$

\mathbb{G} is a **universal sheaf** on $X \times M$
 $v \in H^{p,q}(X)$, \bar{v} its Poincaré dual

3. Euler pairing and Todd classes

$$\text{td}(Q) = \text{id} - A^Q + S^Q : \mathbb{Z}^V \rightarrow \mathbb{Z}^V = K^0(Q)$$

$$A_{v,w}^Q = \#\{e \in E : t(e) = v, h(e) = w\}$$

$$S_{v,w}^Q = \#\{r \in R : t(r) = v, h(r) = w\}$$

Euler pairing $\chi(\bar{e}, \bar{d}) = \langle \bar{e}, \text{td}(Q)\bar{d} \rangle$

Todd class $\text{td}(X) : K^0(X, \mathbb{Q}) \cong H^*(X) \rightarrow H^*(X)$

Euler pairing $\chi(v, w) = \int_X v^\vee \cdot w \cdot \text{td}(X)$

Virasoro operators

1. Quadratic terms from the diagonal pushforward

$$\begin{aligned}\Delta_* \mathrm{td}(Q) &= \sum_{v \in V} \mathrm{td}(Q) \cdot v \boxtimes v \\ &=: \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V\end{aligned}$$

$$\begin{aligned}\Delta_* \mathrm{td}(X) &= \sum_{v \in B} \mathrm{td}(X) \cdot \bar{v} \boxtimes v \\ &=: \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X), \\ \bar{v} \in \bar{B} \text{ satisfy } \int_X w \cdot \bar{v} &= \delta_{w,v}\end{aligned}$$

Virasoro operators

1. Quadratic terms from the diagonal pushforward

$$\Delta_* \text{td}(Q) = \sum_{v \in V} \text{td}(Q) \cdot v \boxtimes v \\ =: \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\Delta_* \text{td}(X) = \sum_{v \in B} \text{td}(X) \cdot \bar{v} \boxtimes v \\ =: \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X), \\ \bar{v} \in \bar{B} \text{ satisfy } \int_X w \cdot \bar{v} = \delta_{w,v}$$

$$\text{write } \tau_i \tau_j(\gamma) = \sum_v \tau_i(v^L) \tau_j(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\tau_i \tau_j(\gamma) = \sum_v \tau_i^H(v^L) \tau_j^H(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X)$$

2. Virasoro operators $L_k = T_k + R_k$

The operators R_k are the **same**

$$R_k(\tau_i^{(H)}(v)) = i^{(k+1)} \tau_{i+k}^{(H)}(v)$$

where $i^{(k+1)} = i(i+1) \cdots (i+k)$ is the rising factorial.

The operators T_k are expressed **in terms of** $\Delta_* \text{td}(-)$.

Virasoro operators

1. Quadratic terms from the diagonal pushforward

$$\Delta_* \text{td}(Q) = \sum_{v \in V} \text{td}(Q) \cdot v \boxtimes v \\ =: \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\Delta_* \text{td}(X) = \sum_{v \in B} \text{td}(X) \cdot \bar{v} \boxtimes v \\ =: \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X), \\ \bar{v} \in \bar{B} \text{ satisfy } \int_X w \cdot \bar{v} = \delta_{w,v}$$

$$\text{write } \tau_i \tau_j(\gamma) = \sum_v \tau_i(v^L) \tau_j(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\tau_i \tau_j(\gamma) = \sum_v \tau_i^H(v^L) \tau_j^H(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X)$$

2. Virasoro operators $L_k = T_k + R_k$

The operators R_k are the **same**

$$R_k(\tau_i^{(H)}(v)) = i^{(k+1)} \tau_{i+k}^{(H)}(v)$$

where $i^{(k+1)} = i(i+1) \cdots (i+k)$ is the rising factorial.

The operators T_k are expressed **in terms of** $\Delta_* \text{td}(-)$.

3. We can **predict** the form of T_k from the **virtual tangent bundle** T^{vir} .

$$\text{ch}(T^{\text{vir}}) = - \sum_{i,j} (-1)^i \tau_i \tau_j(\text{td}(Q))(+1) \quad - \sum_{i,j} (-1)^{i+p_L} \tau_i^H \tau_j^H(\text{td}(X))(+1),$$

Virasoro operators

1. Quadratic terms from the diagonal pushforward

$$\Delta_* \text{td}(Q) = \sum_{v \in V} \text{td}(Q) \cdot v \boxtimes v \\ =: \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\Delta_* \text{td}(X) = \sum_{v \in B} \text{td}(X) \cdot \bar{v} \boxtimes v \\ =: \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X), \\ \bar{v} \in \bar{B} \text{ satisfy } \int_X w \cdot \bar{v} = \delta_{w,v}$$

$$\text{write } \tau_i \tau_j(\gamma) = \sum_v \tau_i(v^L) \tau_j(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\tau_i \tau_j(\gamma) = \sum_v \tau_i^H(v^L) \tau_j^H(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X)$$

2. Virasoro operators $L_k = T_k + R_k$

The operators R_k are the **same**

$$R_k(\tau_i^{(H)}(v)) = i^{(k+1)} \tau_{i+k}^{(H)}(v)$$

where $i^{(k+1)} = i(i+1) \cdots (i+k)$ is the rising factorial.

The operators T_k are expressed **in terms of** $\Delta_* \text{td}(-)$.

3. We can **predict** the form of T_k from the **virtual tangent bundle** T^{vir} .

$$\text{ch}(T^{\text{vir}}) = - \sum_{i,j} (-1)^i \tau_i \tau_j (\text{td}(Q)) (+1) \quad - \sum_{i,j} (-1)^{i+p_L} \tau_i^H \tau_j^H (\text{td}(X)) (+1),$$

$$T_k = \sum_{i+j=k} i! j! \tau_i \tau_j (\Delta_* \text{td}(Q)) (+1) \quad \sum_{i+j=k} (-1)^{p_L} i! j! \tau_i^H \tau_j^H (\Delta_* \text{td}(X)) (+1).$$

Virasoro operators

1. Quadratic terms from the diagonal pushforward

$$\Delta_* \text{td}(Q) = \sum_{v \in V} \text{td}(Q) \cdot v \boxtimes v \\ =: \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\Delta_* \text{td}(X) = \sum_{v \in B} \text{td}(X) \cdot \bar{v} \boxtimes v \\ =: \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X), \\ \bar{v} \in \bar{B} \text{ satisfy } \int_X w \cdot \bar{v} = \delta_{w,v}$$

$$\text{write } \tau_i \tau_j(\gamma) = \sum_v \tau_i(v^L) \tau_j(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in \mathbb{Z}^V \otimes \mathbb{Z}^V$$

$$\tau_i \tau_j(\gamma) = \sum_v \tau_i^H(v^L) \tau_j^H(v^R) \\ \text{for } \gamma = \sum_v v^L \otimes v^R \in H^*(X) \otimes H^*(X)$$

2. Virasoro operators $L_k = T_k + R_k$

The operators R_k are the **same**

$$R_k(\tau_i^{(H)}(v)) = i^{(k+1)} \tau_{i+k}^{(H)}(v)$$

where $i^{(k+1)} = i(i+1) \cdots (i+k)$ is the rising factorial.

The operators T_k are expressed **in terms of** $\Delta_* \text{td}(-)$.

3. We can **predict** the form of T_k from the **virtual tangent bundle** T^{vir} .

$$\text{ch}(T^{\text{vir}}) = - \sum_{i,j} (-1)^i \tau_i \tau_j(\text{td}(Q))(+1) \quad - \sum_{i,j} (-1)^{i+p_L} \tau_i^H \tau_j^H(\text{td}(X))(+1),$$

$$T_k = \sum_{i+j=k} i!j! \tau_i \tau_j(\Delta_* \text{td}(Q))(+1) \quad \sum_{i+j=k} (-1)^{p_L} i!j! \tau_i^H \tau_j^H(\Delta_* \text{td}(X))(+1).$$

4. All of this can be done for **any nice additive category**! This will be clear from the proof.

Defining Virasoro constraints

1. The operators L_k satisfy the **dual** Virasoro commutation relations

$$[L_m, L_n] = (n - m)L_{m+n}.$$

Defining Virasoro constraints

1. The operators L_k satisfy the **dual** Virasoro commutation relations

$$[L_m, L_n] = (n - m)L_{m+n}.$$

2. **Naïve guess (not quite correct):** If M is fine and carries a virtual fundamental class $[M]^{\text{vir}}$, then

$$\int_{[M]^{\text{vir}}} L_k(D) = 0 \quad \text{for } k \geq -1, D \in \mathbb{D}^{Q/X}.$$

Defining Virasoro constraints

1. The operators L_k satisfy the **dual** Virasoro commutation relations

$$[L_m, L_n] = (n - m)L_{m+n}.$$

2. **Naïve guess (not quite correct):** If M is fine and carries a virtual fundamental class $[M]^{\text{vir}}$, then

$$\int_{[M]^{\text{vir}}} L_k(D) = 0 \quad \text{for } k \geq -1, D \in \mathbb{D}^{Q/X}.$$

3. Instead, we need to make up for the **non-uniqueness of the choice of a universal object**. Use another operator S^k compatible with the universal object. It can also absorb **fixing determinants** of sheaves.

Claim (B.–Lim–Moreira(22'), B.(23'))

Let M be a fine moduli space of stable objects with a virtual fundamental class, then it often satisfies **Virasoro constraints**

$$\int_{[M]^{\text{vir}}} (L_k + S_k)(D) = 0 \quad \text{for } k \geq 0, D \in \mathbb{D}^{Q/X}.$$

Weight zero Virasoro constraints

1. To avoid talking about S_k , we introduced the **weight-zero operator**

$$L_{\text{wt}=0} = \sum_{n \geq -1} \frac{(-1)^n}{(n+1)!} L_n \circ R_{-1}^{n+1}.$$

Weight zero Virasoro constraints

1. To avoid talking about S_k , we introduced the **weight-zero operator**

$$L_{\text{wt}=0} = \sum_{n \geq -1} \frac{(-1)^n}{(n+1)!} L_n \circ R_{-1}^{n+1}.$$

Lemma (BLM)

The previous claim is equivalent to

$$\int_{[M]^{\text{vir}}} L_{\text{wt}=0}(D) = 0 \quad \text{for } D \in \mathbb{D}^{Q/X}.$$

Weight zero Virasoro constraints

1. To avoid talking about S_k , we introduced the **weight-zero operator**

$$L_{\text{wt}=0} = \sum_{n \geq -1} \frac{(-1)^n}{(n+1)!} L_n \circ R_{-1}^{n+1}.$$

Lemma (BLM)

The previous claim is equivalent to

$$\int_{[M]^{\text{vir}}} L_{\text{wt}=0}(D) = 0 \quad \text{for } D \in \mathbb{D}^{Q/X}.$$

2. This formulation a) is **independent of the choice** of the universal object when it exists, b) can be defined **formally without the universal object**.

Flag varieties

1. The **partial flag variety** $\text{Flag}(d_1, d_2, \dots, d_l)$ for $d_1 > d_2 \cdots d_{l-1} > d_l$ parametrizes sequences of quotients

$$\mathbb{C}^{d_1} \twoheadrightarrow \mathbb{C}^{d_2} \twoheadrightarrow \cdots \twoheadrightarrow \mathbb{C}^{d_{l-1}} \twoheadrightarrow \mathbb{C}^{d_l}.$$

and it carries the **universal quotients** \mathcal{Q}_v for $v = 2, \dots, l$.


Flag varieties

1. The **partial flag variety** $\text{Flag}(d_1, d_2, \dots, d_l)$ for $d_1 > d_2 > \dots > d_{l-1} > d_l$ parametrizes sequences of quotients

$$\mathbb{C}^{d_1} \twoheadrightarrow \mathbb{C}^{d_2} \twoheadrightarrow \dots \twoheadrightarrow \mathbb{C}^{d_{l-1}} \twoheadrightarrow \mathbb{C}^{d_l}.$$

and it carries the **universal quotients** Q_v for $v = 2, \dots, l$.

2. Setting $n_1 = d_1$ and $n_v = 0$ for $v = 2, \dots, l$ in




Flag varieties

1. The **partial flag variety** $\text{Flag}(d_1, d_2, \dots, d_l)$ for $d_1 > d_2 > \dots > d_{l-1} > d_l$ parametrizes sequences of quotients

$$\mathbb{C}^{d_1} \twoheadrightarrow \mathbb{C}^{d_2} \twoheadrightarrow \dots \twoheadrightarrow \mathbb{C}^{d_{l-1}} \twoheadrightarrow \mathbb{C}^{d_l}.$$

and it carries the **universal quotients** \mathcal{Q}_v for $v = 2, \dots, l$.

2. Setting $n_1 = d_1$ and $n_v = 0$ for $v = 2, \dots, l$ in



and choosing the right stability condition, we obtain the flag variety as a moduli space of representations of this quiver. We also identify $\mathcal{Q}_v = \mathbb{U}_v$.

3. The only non-constant descendents are $\tau_i(v) = \text{ch}_i(\mathcal{Q}_v)$ for $v \neq \infty, 1$. We see then that

$$T_k = \sum_{\substack{i+j=k \\ v \geq 2}} \tau_i(v) (\tau_j(v) - \tau_j(v+1)) - d_1 \tau_k(2),$$

and that R_k only acts on these descendents.


Flag varieties

1. The **partial flag variety** $\text{Flag}(d_1, d_2, \dots, d_l)$ for $d_1 > d_2 > \dots > d_{l-1} > d_l$ parametrizes sequences of quotients

$$\mathbb{C}^{d_1} \twoheadrightarrow \mathbb{C}^{d_2} \twoheadrightarrow \dots \twoheadrightarrow \mathbb{C}^{d_{l-1}} \twoheadrightarrow \mathbb{C}^{d_l}.$$

and it carries the **universal quotients** \mathcal{Q}_v for $v = 2, \dots, l$.

2. Setting $n_1 = d_1$ and $n_v = 0$ for $v = 2, \dots, l$ in



and choosing the right stability condition, we obtain the flag variety as a moduli space of representations of this quiver. We also identify $\mathcal{Q}_v = \mathbb{U}_v$.

3. The only non-constant descendents are $\tau_i(v) = \text{ch}_i(\mathcal{Q}_v)$ for $v \neq \infty, 1$. We see then that

$$T_k = \sum_{\substack{i+j=k \\ v \geq 2}} \tau_i(v) (\tau_j(v) - \tau_j(v+1)) - d_1 \tau_k(2),$$

and that R_k only acts on these descendents.

4. Choosing a polynomial D in $\text{ch}_i(\mathcal{Q}_v)$ for $v \geq 2$, Virasoro constraints tell us that

$$\int_{\text{Flag}} (T_k + R_k)(D) = 0 \quad \text{for } k \geq 0.$$

Dual algebra to \mathbb{D}^Q

1. From now on, I will focus on **quivers** as everything works in a similar way for varieties X satisfying $H^{p,q}(X) = 0$ whenever $|p - q| > 1$.

Dual algebra to \mathbb{D}^Q

1. From now on, I will focus on **quivers** as everything works in a similar way for varieties X satisfying $H^{p,q}(X) = 0$ whenever $|p - q| > 1$.
2. In general, $H^*(M)$ is not freely generated by $\tau_i(v)$. So we take the larger stack \mathcal{M}_Q of **all perfect complexes** of representations. It has a universal representation $\{\mathcal{U}_v\}$ giving cohomology classes $\tau_i(v) \in H^*(\mathcal{M}_Q)$.

Dual algebra to \mathbb{D}^Q

1. From now on, I will focus on **quivers** as everything works in a similar way for varieties X satisfying $H^{p,q}(X) = 0$ whenever $|p - q| > 1$.
2. In general, $H^*(M)$ is not freely generated by $\tau_i(v)$. So we take the larger stack \mathcal{M}_Q of **all perfect complexes** of representations. It has a universal representation $\{\mathcal{U}_v\}$ giving cohomology classes $\tau_i(v) \in H^*(\mathcal{M}_Q)$.
3. Because we assumed that M is fine, there is a **choice of a map** $\iota : M \rightarrow \mathcal{M}_Q$. The universal representations and thus $\tau_i(v)$ are **compatible** under ι^* .

Dual algebra to \mathbb{D}^Q

1. From now on, I will focus on **quivers** as everything works in a similar way for varieties X satisfying $H^{p,q}(X) = 0$ whenever $|p - q| > 1$.
2. In general, $H^*(M)$ is not freely generated by $\tau_i(v)$. So we take the larger stack \mathcal{M}_Q of **all perfect complexes** of representations. It has a universal representation $\{\mathcal{U}_v\}$ giving cohomology classes $\tau_i(v) \in H^*(\mathcal{M}_Q)$.
3. Because we assumed that M is fine, there is a **choice of a map** $\iota : M \rightarrow \mathcal{M}_Q$. The universal representations and thus $\tau_i(v)$ are **compatible** under ι^* .
4. We know that $H^*(\mathcal{M}_{\vec{d}})$ for the connected component $\mathcal{M}_{\vec{d}} \subset \mathcal{M}_Q$ is (often) **isomorphic** to \mathbb{D}^Q :

$$H^*(\mathcal{M}_Q) = \text{Sym}[\tau_i(v), i > 0, v \in V].$$

Dual algebra to \mathbb{D}^Q

1. From now on, I will focus on **quivers** as everything works in a similar way for varieties X satisfying $H^{p,q}(X) = 0$ whenever $|p - q| > 1$.
2. In general, $H^*(M)$ is not freely generated by $\tau_i(v)$. So we take the larger stack \mathcal{M}_Q of **all perfect complexes** of representations. It has a universal representation $\{\mathcal{U}_v\}$ giving cohomology classes $\tau_i(v) \in H^*(\mathcal{M}_Q)$.
3. Because we assumed that M is fine, there is a **choice of a map** $\iota : M \rightarrow \mathcal{M}_Q$. The universal representations and thus $\tau_i(v)$ are **compatible** under ι^* .
4. We know that $H^*(\mathcal{M}_{\bar{d}})$ for the connected component $\mathcal{M}_{\bar{d}} \subset \mathcal{M}_Q$ is (often) **isomorphic** to \mathbb{D}^Q :

$$H^*(\mathcal{M}_Q) = \text{Sym}[\tau_i(v), i > 0, v \in V].$$

5. Define homology classes $t_{k,v}$ as duals of $\tau_k(v)$:

$$\tau_k(v) \cap (-) = \frac{\partial}{\partial t_{k,v}}.$$

6. The **homology** of $\mathcal{M}_{\bar{d}}$ is the polynomial algebra

$$H_*(\mathcal{M}_{\bar{d}}) = e^{\bar{d}} \otimes \text{Sym}[t_{i,v}, i > 0, v \in V].$$

Recovering a GW-like formulation of Virasoro constraints

1. Let us introduce the notation $\langle \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n) \rangle_M = \int_{[M]^{\text{vir}}} \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n)$

Recovering a GW-like formulation of Virasoro constraints

1. Let us introduce the notation $\langle \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n) \rangle_M = \int_{[M]^{\text{vir}}} \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n)$
2. It becomes clear that

$$\begin{aligned} \iota_*[M]^{\text{vir}} &= \left\langle \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right] \right\rangle_M \\ &= \int_{[M]^{\text{vir}}} \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right], \end{aligned}$$

which replaces F^X from GW theory. Under the integral, one should view $t_{k,v}$ as formal variables which turn into homology classes after evaluation.

Recovering a GW-like formulation of Virasoro constraints

1. Let us introduce the notation $\langle \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n) \rangle_M = \int_{[M]^{\text{vir}}} \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n)$
2. It becomes clear that

$$\begin{aligned} \iota_*[M]^{\text{vir}} &= \left\langle \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right] \right\rangle_M \\ &= \int_{[M]^{\text{vir}}} \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right], \end{aligned}$$

which replaces F^X from GW theory. Under the integral, one should view $t_{k,v}$ as formal variables which turn into homology classes after evaluation.

3. The GW formulation of Virasoro constraints is dual to the one given previously. So we define $T_k = T_k^*, R_k = R_k^*, S_k = S_k^*, L_k = L_k^*$ on $H_*(\mathcal{M}_Q)$.

Recovering a GW-like formulation of Virasoro constraints

1. Let us introduce the notation $\langle \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n) \rangle_M = \int_{[M]^{\text{vir}}} \tau_{k_1}(v_1) \cdots \tau_{k_n}(v_n)$
2. It becomes clear that

$$\begin{aligned} \iota_*[M]^{\text{vir}} &= \left\langle \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right] \right\rangle_M \\ &= \int_{[M]^{\text{vir}}} \exp \left[\sum_{k,v} \tau_k(v) t_{k,v} \right], \end{aligned}$$

which replaces F^X from GW theory. Under the integral, one should view $t_{k,v}$ as formal variables which turn into homology classes after evaluation.

3. The GW formulation of Virasoro constraints is dual to the one given previously. So we define $T_k = T_k^*$, $R_k = R_k^*$, $S_k = S_k^*$, $L_k = L_k^*$ on $H_*(\mathcal{M}_Q)$.
4. As before T_k is a second order differential operator in t 's and R_k is a degree changing operator

$$T_k = \sum_{i+j=k} i!j! \tau_i \tau_j (\text{td}(Q)) \cap,$$

$$R_k = \sum_{\substack{j \geq 1 \\ v \in V}} j_{(k+1)} t_{j-k,v} \frac{\partial}{\partial t_{j,v}}.$$

Here $a_{(b)} = a(a-1) \cdots (a-b+1)$ is the falling factorial.

Virasoro constraints for sheaves: the operators

1. The homology version of Virasoro constraints:

$$(L_k + S_k) \iota_* [M]^{\text{vir}} = 0 \quad \text{for } k \geq 0$$

equivalently $L_{\text{wt}=0} \iota_* [M]^{\text{vir}} = 0$.

Virasoro constraints for sheaves: the operators

1. The homology version of Virasoro constraints:

$$(L_k + S_k) \iota_* [M]^{\text{vir}} = 0 \quad \text{for } k \geq 0$$

equivalently $L_{\text{wt}=0} \iota_* [M]^{\text{vir}} = 0$.

Virasoro constraints for sheaves: the operators

1. The homology version of Virasoro constraints:

$$(L_k + S_k) \iota_* [M]^{\text{vir}} = 0 \quad \text{for } k \geq 0$$


equivalently $L_{\text{wt}=0} \iota_* [M]^{\text{vir}} = 0$.

2. The operator on the second line is defined by

$$L_{\text{wt}=0} = \sum_{n \geq -1} \frac{(-1)^n}{(n+1)!} T^{n+1} \circ L_n.$$


Wall-crossing and vertex algebras

1. I mainly use Joyce's geometric vertex algebras and wall-crossing for proving Virasoro constraints



Wall-crossing and vertex algebras


1. I mainly use Joyce's geometric vertex algebras and wall-crossing for proving Virasoro constraints




2. This wall-crossing allows us to compare virtual fundamental classes $[M_{\overline{d}}^{\sigma}]^{\text{vir}}$, $[M_{\overline{d}}^{\sigma'}]^{\text{vir}}$ of σ and σ' -stable representations in terms of some Lie bracket $[-, -]$ on the quotient $K_* = H_*(\mathcal{M}_Q)/T$.

Wall-crossing and vertex algebras

1. I mainly use Joyce's geometric vertex algebras and wall-crossing for proving Virasoro constraints




2. This wall-crossing allows us to compare virtual fundamental classes $[M_d^\sigma]^\text{vir}$, $[M_d^{\sigma'}]^\text{vir}$ of σ and σ' -stable representations in terms of some Lie bracket $[-, -]$ on the quotient $K_* = H_*(\mathcal{M}_Q)/T$.
3. The Lie bracket comes from a vertex algebra $H_*(\mathcal{M}_Q)$ leading to the following schematic of the proof




Wall-crossing and vertex algebras

1. I mainly use Joyce's geometric vertex algebras and wall-crossing for proving Virasoro constraints



2. This wall-crossing allows us to compare virtual fundamental classes $[M_d^\sigma]^\text{vir}$, $[M_d^{\sigma'}]^\text{vir}$ of σ and σ' -stable representations in terms of some Lie bracket $[-, -]$ on the quotient $K_* = H_*(\mathcal{M}_Q)/T$.
3. The Lie bracket comes from a vertex algebra $H_*(\mathcal{M}_Q)$ leading to the following schematic of the proof



4. The upshot is not only the interpretation that VFC's satisfying Virasoro constraints are physical states, but also that this property is preserved under changing stability conditions.

Example of a wall-crossing formula

1. We never discussed what happens when we have **strictly semistables**. In this case, there is a geometric way of using the **framed quiver** to define classes $[M_d^\sigma]^{\text{in}} \in K_*$ counting semistables.

Example of a wall-crossing formula

1. We never discussed what happens when we have **strictly semistables**. In this case, there is a geometric way of using the **framed quiver** to define classes $[M_d^\sigma]^{\text{in}} \in K_*$ counting semistables.
2. For a quiver Q with relations but **without cycles**, there is a stability condition σ_0 such that

$$[M_d^{\sigma_0}]^{\text{in}} = \delta_{\vec{d}, v}.$$

Example of a wall-crossing formula

1. We never discussed what happens when we have **strictly semistables**. In this case, there is a geometric way of using the **framed quiver** to define classes $[M_{\vec{d}}^{\sigma}]^{\text{in}} \in K_*$ counting semistables.
2. For a quiver Q with relations but **without cycles**, there is a stability condition σ_0 such that

$$[M_{\vec{d}}^{\sigma_0}]^{\text{in}} = \delta_{\vec{d}, \nu}.$$

3. The **wall-crossing formula** becomes

$$[M_{\vec{d}}^{\sigma}] = \sum_{\substack{\vec{d}_i = \delta_v, v \in V: \\ \sum_{i=1}^l \vec{d}_i = \vec{d}}} \tilde{U}(\vec{d}_1, \dots, \vec{d}_l; \sigma_0, \sigma) \\ \left[\cdots \left[[M_{\vec{d}_1}^{\sigma_0}], [M_{\vec{d}_2}^{\sigma_0}] \right], [M_{\vec{d}_3}^{\sigma_0}] \right] \cdots, [M_{\vec{d}_l}^{\sigma_0}] \right]$$

Here I neglected writing $(-)^{\text{in}}$ everywhere.

Example of a wall-crossing formula

1. We never discussed what happens when we have **strictly semistables**. In this case, there is a geometric way of using the **framed quiver** to define classes $[M_{\vec{d}}^{\sigma}]^{\text{in}} \in K_*$ counting semistables.
2. For a quiver Q with relations but **without cycles**, there is a stability condition σ_0 such that

$$[M_{\vec{d}}^{\sigma_0}]^{\text{in}} = \delta_{\vec{d}, \nu}.$$

3. The **wall-crossing formula** becomes

$$[M_{\vec{d}}^{\sigma}] = \sum_{\substack{\vec{d}_i = \delta_v, v \in V: \\ \sum_{i=1}^l \vec{d}_i = \vec{d}}} \tilde{U}(\vec{d}_1, \dots, \vec{d}_l; \sigma_0, \sigma) \\ \left[\cdots \left[[M_{\vec{d}_1}^{\sigma_0}], [M_{\vec{d}_2}^{\sigma_0}] \right], [M_{\vec{d}_3}^{\sigma_0}] \right] \cdots, [M_{\vec{d}_l}^{\sigma_0}] \right]$$

Here I neglected writing $(-)^{\text{in}}$ everywhere.

4. This means that all the information about $[M_{\vec{d}}^{\sigma}]^{\text{in}}$ is already contained in the **Lie algebra structure**.


Vertex algebras

1. A **vertex algebra** is the data of a \mathbb{Z} -graded vector space V_* over \mathbb{C} together with
 - 1.1 a **vacuum vector** $|0\rangle \in V_0$,
 - 1.2 a linear operator $T: V_* \rightarrow V_{*+2}$ called the **translation operator**,
 - 1.3 and a **state-field correspondence** which is a degree 0 linear map

$$Y: V_* \longrightarrow \text{End}(V_*)[[z, z^{-1}]],$$

denoted by $Y(a, z) := \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$, where $\deg(z) = -2$.

2. These need to satisfy some axioms that have enlightening **interpretations in terms of CFTs**



$$\lim_{|z| \rightarrow 0} Y(n, z)|0\rangle = N \quad - \text{state-field correspondence}$$

$$e^{zT}|0\rangle = |0\rangle, \quad Y(10, z) = \text{id}_1$$

- vacuum invariance

$$(z_1 - z_2)^N Y(n, z_1) Y(w_1, z_2) = (z_1 - z_2)^N Y(nw_1, z_2) Y(n, z_1)$$

$$Y(n, z)\omega = e^{zT} Y(w_1 - z)\omega$$



- skew-symmetry



- Locality

Sketch of the geometric construction of vertex algebras

1. Joyce (17') constructed vertex algebras on the homology $V_* = H_{*+\mathrm{vdim}_{\mathbb{C}}}(\mathcal{M}_Q)$, which requires **three ingredients** related to additivity:

Sketch of the geometric construction of vertex algebras

1. Joyce (17') constructed vertex algebras on the homology $V_* = H_{*+\mathrm{vdim}_{\mathbb{C}}}(\mathcal{M}_Q)$, which requires **three ingredients** related to additivity:

1.1 The zero object $p \xrightarrow{0} \mathcal{M}_Q$ gives $|0\rangle = 0_*(p)$.

Sketch of the geometric construction of vertex algebras

1. Joyce (17') constructed vertex algebras on the homology $V_* = H_{*+\mathrm{vdim}_{\mathbb{C}}}(\mathcal{M}_Q)$, which requires **three ingredients** related to additivity:
 - 1.1 The zero object $p \xrightarrow{0} \mathcal{M}_Q$ gives $|0\rangle = 0_*(p)$.
 - 1.2 There is a natural action of $B\mathbb{G}_m$ on \mathcal{M}_Q that rescales the automorphisms of objects. On the level of homology, this action produces the operator T .

Sketch of the geometric construction of vertex algebras

1. Joyce (17') constructed vertex algebras on the homology $V_* = H_{*+\text{vdim}_{\mathbb{C}}}(\mathcal{M}_Q)$, which requires **three ingredients** related to additivity:
 - 1.1 The zero object $p \xrightarrow{0} \mathcal{M}_Q$ gives $|0\rangle = 0_*(p)$.
 - 1.2 There is a natural action of $B\mathbb{G}_m$ on \mathcal{M}_Q that rescales the automorphisms of objects. On the level of homology, this action produces the operator T .
 - 1.3 The complex

$$\text{Ext}_Q = \bigoplus_{v \in V \setminus F} \mathcal{U}_v^\vee \boxtimes \mathcal{U}_v \xrightarrow{\varphi_E} \bigoplus_{e \in E} \mathcal{U}_{t(e)}^\vee \boxtimes \mathcal{U}_{h(e)} \xrightarrow{s_R} \bigoplus_{r \in R} \mathcal{U}_{t(r)}^\vee \boxtimes \mathcal{U}_{h(r)}$$

on $\mathcal{M}_Q \times \mathcal{M}_Q$ is the last piece necessary to write down $Y(v, z)$. Satisfies:

$$T^{\text{vir}} = \Delta^* \text{Ext}[1] + 1$$

This is why Virasoro constraints see T^{vir} .

Sketch of the geometric construction of vertex algebras

1. Joyce (17') constructed vertex algebras on the homology $V_* = H_{*+\mathrm{vdim}_{\mathbb{C}}}(\mathcal{M}_Q)$, which requires **three ingredients** related to additivity:

1.1 The zero object $p \xrightarrow{0} \mathcal{M}_Q$ gives $|0\rangle = 0_*(p)$.

1.2 There is a natural action of $B\mathbb{G}_m$ on \mathcal{M}_Q that rescales the automorphisms of objects. On the level of homology, this action produces the operator T .

1.3 The complex

$$\mathrm{Ext}_Q = \bigoplus_{v \in V \setminus F} \mathcal{U}_v^\vee \boxtimes \mathcal{U}_v \xrightarrow{\varphi_E} \bigoplus_{e \in E} \mathcal{U}_{t(e)}^\vee \boxtimes \mathcal{U}_{h(e)} \xrightarrow{s_R} \bigoplus_{r \in R} \mathcal{U}_{t(r)}^\vee \boxtimes \mathcal{U}_{h(r)}$$

on $\mathcal{M}_Q \times \mathcal{M}_Q$ is the last piece necessary to write down $Y(v, z)$. Satisfies:

$$T^{\mathrm{vir}} = \Delta^* \mathrm{Ext}[1] + 1$$

This is why Virasoro constraints see T^{vir} .

2. The vertex algebra V_* was shown to be a **lattice vertex algebra** by Joyce (17'), which is the most natural vertex algebra with the underlying graded vector space

$$V_* = \mathbb{Q}[\mathbb{Z}^V] \otimes \mathrm{Sym}[[t_{i,v}, i > 0, v \in V]].$$

The geometric construction of a Lie algebra

1. The **stacky quotient** of \mathcal{M}_Q by $B\mathbb{G}_m$ is denoted by $\mathcal{M}_Q^{\text{rig}}$. As one would expect, there is roughly the correspondence

$$K_* = V_{*+2}/TV_* = H_{*+\text{vdim}_{\mathbb{C}}}(\mathcal{M}_Q^{\text{rig}}).$$

The geometric construction of a Lie algebra

1. The **stacky quotient** of \mathcal{M}_Q by $B\mathbb{G}_m$ is denoted by $\mathcal{M}_Q^{\text{rig}}$. As one would expect, there is roughly the correspondence

$$K_* = V_{*+2}/TV_* = H_{*+\text{vdim}_{\mathbb{C}}}(\mathcal{M}_Q^{\text{rig}}).$$

2. Now there are **two roles** that K_* plays:

2.1 The classes $[M]^{\text{vir}}$ live in K_* even if M is not fine.

2.2 K_* has a Lie bracket:

$$[\overline{v}, \overline{w}] = \overline{v_0 w}, \quad \forall v, w \in V_*.$$

Conformal element

1. A conformal element $\omega \in V_4$ leads to a field $Y(\omega, z) = \sum_{k \in \mathbb{Z}} L_k z^{-k-2}$ with

$$[L_n, L_m] = (n - m)L_{m+n} + \frac{n^3 - n}{12} \delta_{n+m,0} \cdot C,$$

which implies that ω generates a Virasoro subalgebra of V_* .

Conformal element

1. A conformal element $\omega \in V_4$ leads to a field $Y(\omega, z) = \sum_{k \in \mathbb{Z}} L_k z^{-k-2}$ with

$$[L_n, L_m] = (n - m)L_{m+n} + \frac{n^3 - n}{12} \delta_{n+m,0} \cdot C,$$

which implies that ω generates a Virasoro subalgebra of V_* .

2. In general, there is no ω in the geometric V_* when the symmetrized pairing $\chi_{\text{sym}}(v, w) = \chi(v, w) + \chi(w, v)$ is degenerate. However, one can define a larger vertex algebra containing ω such that L_k 's restrict to V_* .

Conformal element

1. A conformal element $\omega \in V_4$ leads to a field $Y(\omega, z) = \sum_{k \in \mathbb{Z}} L_k z^{-k-2}$ with

$$[L_n, L_m] = (n - m)L_{m+n} + \frac{n^3 - n}{12} \delta_{n+m,0} \cdot C,$$

which implies that ω generates a Virasoro subalgebra of V_* .

2. In general, there is no ω in the geometric V_* when the symmetrized pairing $\chi_{\text{sym}}(v, w) = \chi(v, w) + \chi(w, v)$ is degenerate. However, one can define a larger vertex algebra containing ω such that L_k 's restrict to V_* .
3. To make everything simpler, I will assume that χ_{sym} is non-degenerate, so that there exists a dual basis $\hat{v} \subset \hat{V}$ to V . Then

$$\omega = \frac{1}{2} \sum_{v \in V} t_{1,v} t_{1,\hat{v}}$$

and $C = |V|$.

Conformal element

1. A conformal element $\omega \in V_4$ leads to a field $Y(\omega, z) = \sum_{k \in \mathbb{Z}} L_k z^{-k-2}$ with

$$[L_n, L_m] = (n - m)L_{m+n} + \frac{n^3 - n}{12} \delta_{n+m,0} \cdot C,$$

which implies that ω generates a Virasoro subalgebra of V_* .

2. In general, there is no ω in the geometric V_* when the symmetrized pairing $\chi_{\text{sym}}(v, w) = \chi(v, w) + \chi(w, v)$ is degenerate. However, one can define a larger vertex algebra containing ω such that L_k 's restrict to V_* .
3. To make everything simpler, I will assume that χ_{sym} is non-degenerate, so that there exists a dual basis $\hat{v} \subset \hat{V}$ to V . Then

$$\omega = \frac{1}{2} \sum_{v \in V} t_{1,v} t_{1,\hat{v}}$$

and $C = |V|$.

Remark

When working with varieties X , we need to include $H^{\text{odd}}(X) \cong K^1(X, \mathbb{Q})$ into the lattice. One can still write $\omega = \frac{1}{2} \sum_{v \in B} t_{1,v}^H t_{1,\hat{v}}^H$ where H denotes some holomorphic grading shift leading to odd degrees. The conformal charge is given by $\chi(X)$.

Physical states and the main claim

1. Before stating the main observation of the works BLM(22') and B.(23'), I will remark that

$$\int_{[M_d^\sigma]^{\text{in}}} L_{\text{wt}=0}(D) = 0 \quad \text{for } D \in H^*(\mathcal{M}_Q)$$

is a well-defined **generalization of Virasoro constraints**.

Physical states and the main claim

1. Before stating the main observation of the works BLM(22') and B.(23'), I will remark that

$$\int_{[M_d^\sigma]^{\text{in}}} L_{\text{wt}=0}(D) = 0 \quad \text{for } D \in H^*(\mathcal{M}_Q)$$

is a well-defined **generalization of Virasoro constraints**.

2. Borchers gave a definition of **physical states** $p \in P \subset V_2$ as satisfying the equations

$$L_k p = \delta_{k,0} p, \quad \text{for } k \geq 0.$$

and showed that they form a Lie subalgebra $\check{P} \subset K_0$.

Physical states and the main claim

1. Before stating the main observation of the works BLM(22') and B.(23'), I will remark that

$$\int_{[M_d^\sigma]^{\text{in}}} L_{\text{wt}=0}(D) = 0 \quad \text{for } D \in H^*(\mathcal{M}_Q)$$

is a well-defined **generalization of Virasoro constraints**.

2. Borchers gave a definition of **physical states** $p \in P \subset V_2$ as satisfying the equations

$$L_k p = \delta_{k,0} p, \quad \text{for } k \geq 0.$$

and showed that they form a Lie subalgebra $\check{P} \subset K_0$.

Theorem (BLM(22'), B. (23'))

The condition that $[M_\alpha^\sigma]^{\text{in}}$ satisfies Virasoro constraints is equivalent to it being a physical state with respect to the ω given above. I.e. $[M_\alpha^\sigma]^{\text{in}} \in \check{P}$.

In particular, wall-crossing, stated in terms of iterated Lie brackets, preserves Virasoro constraints from the RHS to the LHS of the formula.

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*
4. *(BLM) Slope and Gieseker semistable positive rank sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*
4. *(BLM) Slope and Gieseker semistable positive rank sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
5. *(BLM) + B.(24') Slope semistable dimension 1 sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*
4. *(BLM) Slope and Gieseker semistable positive rank sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
5. *(BLM) + B.(24') Slope semistable dimension 1 sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
6. *(BLM) Slope semistable vector bundles on curves.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*
4. *(BLM) Slope and Gieseker semistable positive rank sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
5. *(BLM) + B.(24') Slope semistable dimension 1 sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
6. *(BLM) Slope semistable vector bundles on curves.*
7. *(BLM) + B.(21') Quot schemes parametrizing 0-dimensional quotients of vector bundles on curves and surfaces.*

Main consequences

Theorem

Virasoro constraints hold for the following cases:

1. *B. (23') Bridgeland semistable representations of quasi-smooth quivers with frozen vertices.*
2. *B. (23') Stable framed representations of such quasi-smooth quivers.*
3. *B. (23') Gieseker semistable sheaves on \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$.*
4. *(BLM) Slope and Gieseker semistable positive rank sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
5. *(BLM) + B.(24') Slope semistable dimension 1 sheaves on surfaces with $H^1(\mathcal{O}_S) = H^2(\mathcal{O}_S) = 0$.*
6. *(BLM) Slope semistable vector bundles on curves.*
7. *(BLM) + B.(21') Quot schemes parametrizing 0-dimensional quotients of vector bundles on curves and surfaces.*

Remark

In the second point, I used **derived equivalences** of the surfaces to quivers. This is the first proof of Virasoro constraints for sheaves on surfaces independent of Witten's conjecture. Using a universality of Virasoro constraints for $\text{Hilb}^n(S)$, gives an independent proof of Virasoro constraints for any surface S . In particular, this establishes them as an autonomous phenomenon.