

# Kac polynomials and generalised Kac–Moody algebras

Ben Davison, Victor Kac's 80th birthday conference, Rome 2023



# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations

A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ .  
Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .

# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example



Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations

A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ . Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .


# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example

Kronecker quiver:



(multiple edges allowed)

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations


A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ . Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .

# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example



(loops *sometimes* allowed)

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations


A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ . Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .

# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example



(loops *sometimes* allowed)

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations


A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ . Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .

# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example



(loops *sometimes* allowed)

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations

A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ .


Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .

# Quiver representations

## Quivers

A *quiver*  $Q$  is the data of a set of vertices  $Q_0$ , a set of arrows  $Q_1$ , and a pair of morphisms  $s, t: Q_1 \rightarrow Q_0$  taking an arrow to its source/target.

## Example



(loops *sometimes* allowed)

Fix a field  $K$ . The *path algebra*  $KQ$  is a  $K$ -algebra with basis given by paths (including lazy paths  $e_i$  of length 0 at each vertex  $i$ ), and multiplication given by concatenation of paths.

## Representations

A representation of  $Q$  of dimension  $d \in \mathbb{N}^{Q_0}$  is the data of a set of  $K$ -vector spaces  $\rho_i$  of dimension  $d_i$  along with morphisms  $\rho(a): \rho_{s(a)} \rightarrow \rho_{t(a)}$ . Equivalent to giving a  $KQ$ -module  $N$  with  $\dim(e_i \cdot N) = d_i$ .



## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .

## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .


## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .


## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#P_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#P_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .


## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$

- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .


## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .


## Kac polynomials

A  $KQ$ -module  $N$  is called *absolutely indecomposable* if  $N \otimes_K \overline{K}$  is an indecomposable  $\overline{K}Q$ -module. For  $d \in \mathbb{N}^{Q_0}$  Victor Kac defined

$$a_{Q,d}(q) = \{\text{abs. indecomposable } d\text{-dimensional } \mathbb{F}_q Q\text{-modules}\} / \cong$$

### Example

Let  $Q$  be the Kronecker quiver



Let  $N$  be a  $(1,1)$ -dimensional  $KQ$ -module.  $N(a)$  and  $N(b)$  given by scalars. Iff  $N'(a) = \lambda N(a)$  and  $N'(b) = \lambda N(b)$  for  $\lambda \in K^\times$  then  $N \cong N'$ .

- $a_{Q,(1,1)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$
- $a_{Q,(n,n)}(q) = \#\mathbb{P}_{\mathbb{F}_q}^1 = q + 1$  (e.g. use Beilinson's equivalence)

### Theorem (Kac)

$Q$  any quiver,  $d \in \mathbb{Z}_{\geq 0}^{Q_0}$ . The function  $a_{Q,d}(q)$  is a polynomial in  $\mathbb{Z}[q]$ .

# Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez-Villegas):  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- Constant term conjecture (Kac), Theorem (Hausel):  $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- (Kac):  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- (Kac): Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .



# Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez–Villegas):  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- Constant term conjecture (Kac), Theorem (Hausel):  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- (Kac):  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- (Kac): Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

# Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- **Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez–Villegas):**  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- **Constant term conjecture (Kac), Theorem (Hausel):**  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- **(Kac):**  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- **(Kac):** Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

# Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez–Villegas):  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- Constant term conjecture (Kac), Theorem (Hausel):  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- (Kac):  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- (Kac): Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

# Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez–Villegas):  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- Constant term conjecture (Kac), Theorem (Hausel):  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- (Kac):  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- (Kac): Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

## Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez–Villegas):  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- Constant term conjecture (Kac), Theorem (Hausel):  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- (Kac):  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- (Kac): Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

## Kac polynomials: some results

To start with, we allow  $Q$  to have loops

- **Positivity conjecture (Kac), Theorem (Hausel, Letellier, Rodriguez-Villegas):**  $a_{Q,d}(q) \in \mathbb{Z}_{\geq 0}[q]$ .

Now we (temporarily) forbid loops:

- **Constant term conjecture (Kac), Theorem (Hausel):**  
 $a_{Q,d}(0) = \dim(\mathfrak{g}_{Q,d})$ , the  $d$ th weight space of the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .
- **(Kac):**  $a_{Q,d}(q) = 0$  if  $d$  is not a positive root of  $\mathfrak{g}_Q$ .
- **(Kac):** Let  $d, d'$  be positive roots such that  $w(d) = d'$  for some  $w$  in the Weyl group of  $\mathfrak{g}_Q$ . Then  $a_{Q,d}(q) = a_{Q,d'}(q)$ .

What is this hinting at?:

These facts suggest that *all* the coefficients of  $a_{Q,d}(q)$  are the dimensions of the cohomologically graded pieces of a *generalised* Kac–Moody algebra  $\tilde{\mathfrak{g}}_Q$  such that  $\tilde{\mathfrak{g}}_Q^0 = \mathfrak{g}_Q$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_j) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .



# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “*positive roots*”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of *positive Chevalley generators*.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_j) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of *positive Chevalley generators*.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_j) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of *positive Chevalley generators*.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi^+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i)\alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij}\alpha_j^{\vee}(\alpha_i)h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i) \alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij} \alpha_j^{\vee} (\alpha_i) h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)} (\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .

# GKM algebras

We define a class of generalised Kac–Moody algebras, following Borchers.

## Ingredients

- Let  $\mathfrak{h}$  be a  $\mathbb{Q}$ -vector space with symmetric bilinear form  $(-, -)$ .
- Let  $\{h_i\}_{i \in \Phi_+} \subset \mathfrak{h}$  be a countable set of “positive roots”:  
 $(h_i, h_j) \in \mathbb{Z}_{\leq 0}$  if  $i \neq j$  and  $(h_i, h_i) \in 2 \cdot \mathbb{Z}_{\leq 1}$  for all  $i \in \Phi_+$ .
- Pick  $\mathcal{G} = \bigoplus_{i \in \Phi_+} \mathcal{G}_i$ , a  $\Phi_+$ -graded, cohomologically graded set of positive Chevalley generators.

## Recipe

$\mathfrak{g}_{\mathcal{G}} :=$  the free Lie algebra generated by  $\mathfrak{h} \oplus \mathcal{G} \oplus \mathcal{G}^{\vee}$  modulo the relations:

- $[h, h'] = 0$  for all  $h, h' \in \mathfrak{h}$ .
- $[h, \alpha_i^{(\vee)}] = (-)(h, h_i) \alpha_i^{(\vee)} \quad (\alpha_i^{(\vee)} \in \mathcal{G}_i^{(\vee)})$
- $[\alpha_i, \alpha_j^{\vee}] = \delta_{ij} \alpha_j^{\vee}(\alpha_i) h_i$
- (Serre relations):  $[\alpha_i^{(\vee)}, -]^{1-(h_i, h_j)}(\alpha_j^{(\vee)}) = 0$  if  $(h_i, h_j) = 0$  or  $(h_i, h_i) = 2$ .



# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^{\vee}$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.

# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^\vee$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^+$  is the free Lie algebra generated by  $\mathcal{G}$  modulo the Serre relations.



# Features of GKM algebras

## Trichotomy of roots

Roots  $h_i$  come in three types:

- 1 Real:  $(h_i, h_i) = 2$
- 2 Isotropic:  $(h_i, h_i) = 0$
- 3 Hyperbolic:  $(h_i, h_i) < 0$

The last two types of roots are called *imaginary*.

Many of your favourite theorems about Kac–Moody Lie algebras generalise to GKM algebras with minimal modifications, for example:

## Triangular decomposition

There is a decomposition of vector spaces  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{n}_{\mathcal{G}}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_{\mathcal{G}}^-$  into the Lie algebras generated by  $\mathcal{G}, \mathfrak{h}, \mathcal{G}^{\vee}$  respectively. Moreover  $\mathfrak{n}_{\mathcal{G}}^-$  is the free Lie algebra generated by  $\mathcal{G}^{\vee}$  modulo the Serre relations.

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset Q = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset Q = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset Q = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \mathfrak{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset Q = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset Q = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset \mathbb{Q} = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(nc\text{Hilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .  
 N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset \mathbb{Q} = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .



## Examples of GKM algebras

- Fix a quiver  $Q$  (maybe with loops). Fix  $\mathfrak{h} = \mathbb{Q}^{Q_0}$  and  $(d, e) = \chi_Q(d, e) + \chi_Q(e, d) (= 2 \sum_{i \in Q_0} d_i e_i - \sum_{a \in Q_1} (d_{s(a)} e_{t(a)} + d_{t(a)} e_{s(a)}))$ .
- So  $1_i$  is a real simple root if no loops at  $i$ , an isotropic simple root if exactly one loop, and a hyperbolic root otherwise.

### Example (KM Lie algebras)

Assume  $Q$  has no loops. We set  $\Phi_+ = Q_0$  and  $h_i = 1_i$ . Fix  $\mathcal{G}_i = \mathbb{Q}$ . Then  $\mathfrak{g}_Q := \mathfrak{g}_{\mathcal{G}}$  is the Kac–Moody Lie algebra associated to underlying graph of  $Q$ .

### Example (Nakajima's Heisenberg algebra)

Let  $Q$  have one vertex  $i$ , and one loop. Then  $(-, -) = 0$ . Set  $\Phi_+ = \mathbb{Z}_{\geq 1} \subset \mathbb{Q} = \mathfrak{h}$ . Set  $\mathcal{G}_n = \mathbb{Q}[2]$  for all  $n \geq 1$ . Then  $\mathfrak{g}_{\mathcal{G}} = \text{heis}_{\infty}$  is Nakajima's Lie algebra of operators on  $\bigoplus_{n \in \mathbb{Z}_{\geq 0}} H(\text{ncHilb}_n(\mathbb{A}^2), \mathbb{Q}^{\text{vir}})$ .

N.B.  $\chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G}, n}) := \sum_{i \in \mathbb{Z}} \dim(\mathfrak{g}_{\mathcal{G}, n}^i) q^{i/2} = a_{Q, n}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

Given a quiver  $Q$ , Is there a (non-cohomologically graded) GKM algebra  $\mathfrak{g}^\circ$  with  $\dim(\mathfrak{g}_d^\circ) = a_{Q,d}(1)$  for all  $d \in \mathbb{N}^{Q_0}$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .

## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . *Okounkov conjecture*:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .



## Okounkov's conjecture

On the previous slide, we began to see how nonconstant terms in Kac polynomials *might* be recording graded dimensions of an associated GKM algebra, helping to motivate

### Okounkov's question

More ambitiously, is there a (cohomologically graded) GKM algebra  $\mathfrak{g}$  with  $\chi_{q^{1/2}}(\mathfrak{g}_d) = a_{Q,d}(q^{-1})$ ?

This is closely related to *Okounkov's conjecture*:

- Using stable envelopes, Maulik and Okounkov define  $Y_{Q,MO}$ , a subalgebra of the endomorphism algebra of the cohomology of all Nakajima quiver varieties associated to  $Q$ .
- They prove that there is a PBW isomorphism  $Y_{Q,MO} \cong U(\mathfrak{g}_{MO,Q} \otimes H(BC^*, \mathbb{Q}))$ , where  $\mathfrak{g}_{MO,Q}$  is a Lie algebra containing  $\mathfrak{g}_{Q^{real}}$ .
- MO Conjecture:  $\mathfrak{g}_{Q,MO}$  can be defined as coh. graded vector space over  $\mathbb{Q}$ . Okounkov conjecture:  $\chi_{q^{1/2}}(\mathfrak{g}_{MO,Q,d}) = a_{Q,d}(q^{-1})$ .


# The Kronecker quiver revisited

We plot Kac polynomials  $a_{Q,d}(q^{-1})$  against dimension vectors:




# The Kronecker quiver revisited

We plot Kac polynomials  $a_{Q,d}(q^{-1})$  against dimension vectors: Using Kac's vanishing result




# The Kronecker quiver revisited

We plot Kac polynomials  $a_{Q,d}(q^{-1})$  against dimension vectors: Using constant term theorem



# The Kronecker quiver revisited

We plot Kac polynomials  $a_{Q,d}(q^{-1})$  against dimension vectors: By earlier calculations and Weyl-group invariance



# The Kronecker quiver revisited

The following choice of generating  $\mathcal{G}$  is then *forced*:

3	0	0	0	$\mathbb{Q} \oplus \mathbb{Q}[2]$
2	0	0	$\mathbb{Q} \oplus \mathbb{Q}[2]$	0
1	$\mathbb{Q}$	$\mathbb{Q} \oplus \mathbb{Q}[2]$	0	0
0	*	$\mathbb{Q}$	0	0
	0	1	2	3

# The Kronecker quiver revisited

Yielding the GKM algebra  $\mathfrak{g}_{\mathcal{G}}$  with  $\mathbb{Z}_{\geq 0}^{Q_0}$ -graded pieces:

3	0	0	$\mathbb{Q}$	$\mathbb{Q} \oplus \mathbb{Q}[2]$
2	0	$\mathbb{Q}$	$\mathbb{Q} \oplus \mathbb{Q}[2]$	$\mathbb{Q}$
1	$\mathbb{Q}$	$\mathbb{Q} \oplus \mathbb{Q}[2]$	$\mathbb{Q}$	0
0	*	$\mathbb{Q}$	0	0
	0	1	2	3

# Preprojective algebras

- We denote by  $\overline{Q}$  the *doubled quiver*, obtained by adding an arrow  $a^*$  with the opposite orientation to  $a$  for all  $a \in Q_1$ .
- We define the *preprojective algebra*  $\Pi_Q := \mathbb{C}\overline{Q} / \langle \sum_{a \in Q_1} [a, a^*] \rangle$

## Example

If  $Q$  is the one loop quiver then  $\mathbb{C}\overline{Q} = \mathbb{C}\langle a, a^* \rangle$ , the ring of polynomials in two noncommuting variables, and  $\Pi_Q \cong \mathbb{C}[a, a^*]$ .

## Example

If  $Q$  is the Kronecker quiver then  $\Pi_Q$  is derived equivalent to  $\mathrm{QCoh}(\mathrm{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(-2)))$ .



# Preprojective algebras

- We denote by  $\overline{Q}$  the *doubled quiver*, obtained by adding an arrow  $a^*$  with the opposite orientation to  $a$  for all  $a \in Q_1$ .
- We define the *preprojective algebra*  $\Pi_Q := \mathbb{C}\overline{Q}/\langle \sum_{a \in Q_1} [a, a^*] \rangle$

## Example

If  $Q$  is the one loop quiver then  $\mathbb{C}\overline{Q} = \mathbb{C}\langle a, a^* \rangle$ , the ring of polynomials in two noncommuting variables, and  $\Pi_Q \cong \mathbb{C}[a, a^*]$ .

## Example

If  $Q$  is the Kronecker quiver then  $\Pi_Q$  is derived equivalent to  $\mathrm{QCoh}(\mathrm{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(-2)))$ .

# Preprojective algebras

- We denote by  $\overline{Q}$  the *doubled quiver*, obtained by adding an arrow  $a^*$  with the opposite orientation to  $a$  for all  $a \in Q_1$ .
- We define the *preprojective algebra*  $\Pi_Q := \mathbb{C}\overline{Q}/\langle \sum_{a \in Q_1} [a, a^*] \rangle$

## Example

If  $Q$  is the one loop quiver then  $\mathbb{C}\overline{Q} = \mathbb{C}\langle a, a^* \rangle$ , the ring of polynomials in two noncommuting variables, and  $\Pi_Q \cong \mathbb{C}[a, a^*]$ .

## Example

If  $Q$  is the Kronecker quiver then  $\Pi_Q$  is derived equivalent to  $\mathrm{QCoh}(\mathrm{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(-2)))$ .

# Preprojective algebras

- We denote by  $\overline{Q}$  the *doubled quiver*, obtained by adding an arrow  $a^*$  with the opposite orientation to  $a$  for all  $a \in Q_1$ .
- We define the *preprojective algebra*  $\Pi_Q := \mathbb{C}\overline{Q}/\langle \sum_{a \in Q_1} [a, a^*] \rangle$

## Example

If  $Q$  is the one loop quiver then  $\mathbb{C}\overline{Q} = \mathbb{C}\langle a, a^* \rangle$ , the ring of polynomials in two noncommuting variables, and  $\Pi_Q \cong \mathbb{C}[a, a^*]$ .

## Example

If  $Q$  is the Kronecker quiver then  $\Pi_Q$  is derived equivalent to  $\mathrm{QCoh}(\mathrm{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(-2)))$ .

# Moduli spaces

## The stack of $\Pi_Q$ -modules

- Let  $\mathbb{A}_{\overline{Q},d} := \prod_{a \in \overline{Q}_1} \text{Hom}(\mathbb{C}^{d_{s(a)}}, \mathbb{C}^{d_{t(a)}}) \cong T^* \mathbb{A}_{Q,d}$ .
- Moment map

$$\mu_d: \mathbb{A}_{\overline{Q},d} \rightarrow \prod_{i \in Q_0} \mathfrak{gl}_{d_i}$$
$$(N(a), N(a^*))_{a \in Q_1} \mapsto \sum_{a \in Q_1} [N(a), N(a^*)].$$

- The stack  $\mathfrak{M}_d(\Pi_Q)$  of  $d$ -dimensional  $\Pi_Q$ -modules is equivalent to the quotient  $\mu_d^{-1}(0)/\text{GL}_d$  with  $\text{GL}_d := \prod_{i \in Q_0} \text{GL}_{d_i}(\mathbb{C})$ .

## The coarse moduli space

If we wish to have an algebraic variety instead of a stack, we may consider the coarse quotient:  $X_d(\Pi_Q) := \text{Spec}(\Gamma(\mu_d^{-1}(0))^{\text{GL}_d})$

Points of this space are in bijection with  $d$ -dimensional semisimple  $\Pi_Q$ -modules.

# Moduli spaces

## The stack of $\Pi_Q$ -modules

- Let  $\mathbb{A}_{\overline{Q},d} := \prod_{a \in \overline{Q}_1} \text{Hom}(\mathbb{C}^{d_{s(a)}}, \mathbb{C}^{d_{t(a)}}) \cong T^* \mathbb{A}_{Q,d}$ .
- Moment map

$$\mu_d: \mathbb{A}_{\overline{Q},d} \rightarrow \prod_{i \in Q_0} \mathfrak{gl}_{d_i}$$

$$(N(a), N(a^*))_{a \in Q_1} \mapsto \sum_{a \in Q_1} [N(a), N(a^*)].$$

- The stack  $\mathfrak{M}_d(\Pi_Q)$  of  $d$ -dimensional  $\Pi_Q$ -modules is equivalent to the quotient  $\mu_d^{-1}(0)/GL_d$  with  $GL_d := \prod_{i \in Q_0} GL_{d_i}(\mathbb{C})$ .

## The coarse moduli space

If we wish to have an algebraic variety instead of a stack, we may consider the coarse quotient:  $X_d(\Pi_Q) := \text{Spec}(\Gamma(\mu_d^{-1}(0))^{GL_d})$

Points of this space are in bijection with  $d$ -dimensional semisimple  $\Pi_Q$ -modules.

# Moduli spaces

## The stack of $\Pi_Q$ -modules

- Let  $\mathbb{A}_{\overline{Q},d} := \prod_{a \in \overline{Q}_1} \text{Hom}(\mathbb{C}^{d_{s(a)}}, \mathbb{C}^{d_{t(a)}}) \cong T^* \mathbb{A}_{Q,d}$ .
- Moment map

$$\mu_d: \mathbb{A}_{\overline{Q},d} \rightarrow \prod_{i \in Q_0} \mathfrak{gl}_{d_i}$$

$$(N(a), N(a^*))_{a \in Q_1} \mapsto \sum_{a \in Q_1} [N(a), N(a^*)].$$

- The stack  $\mathfrak{M}_d(\Pi_Q)$  of  $d$ -dimensional  $\Pi_Q$ -modules is equivalent to the quotient  $\mu_d^{-1}(0)/\text{GL}_d$  with  $\text{GL}_d := \prod_{i \in Q_0} \text{GL}_{d_i}(\mathbb{C})$ .

## The coarse moduli space

If we wish to have an algebraic variety instead of a stack, we may consider the coarse quotient:  $X_d(\Pi_Q) := \text{Spec}(\Gamma(\mu_d^{-1}(0))^{\text{GL}_d})$

Points of this space are in bijection with  $d$ -dimensional semisimple  $\Pi_Q$ -modules.

# Moduli spaces

## The stack of $\Pi_Q$ -modules

- Let  $\mathbb{A}_{\overline{Q},d} := \prod_{a \in \overline{Q}_1} \text{Hom}(\mathbb{C}^{d_{s(a)}}, \mathbb{C}^{d_{t(a)}}) \cong T^* \mathbb{A}_{Q,d}$ .
- Moment map

$$\mu_d: \mathbb{A}_{\overline{Q},d} \rightarrow \prod_{i \in Q_0} \mathfrak{gl}_{d_i}$$

$$(N(a), N(a^*))_{a \in Q_1} \mapsto \sum_{a \in Q_1} [N(a), N(a^*)].$$

- The stack  $\mathfrak{M}_d(\Pi_Q)$  of  $d$ -dimensional  $\Pi_Q$ -modules is equivalent to the quotient  $\mu_d^{-1}(0)/\text{GL}_d$  with  $\text{GL}_d := \prod_{i \in Q_0} \text{GL}_{d_i}(\mathbb{C})$ .

## The coarse moduli space

If we wish to have an algebraic variety instead of a stack, we may consider the coarse quotient:  $X_d(\Pi_Q) := \text{Spec}(\Gamma(\mu_d^{-1}(0))^{\text{GL}_d})$

Points of this space are in bijection with  $d$ -dimensional semisimple  $\Pi_Q$ -modules.

# Moduli spaces

## The stack of $\Pi_Q$ -modules

- Let  $\mathbb{A}_{\overline{Q},d} := \prod_{a \in \overline{Q}_1} \text{Hom}(\mathbb{C}^{d_{s(a)}}, \mathbb{C}^{d_{t(a)}}) \cong T^* \mathbb{A}_{Q,d}$ .
- Moment map

$$\mu_d: \mathbb{A}_{\overline{Q},d} \rightarrow \prod_{i \in Q_0} \mathfrak{gl}_{d_i}$$

$$(N(a), N(a^*))_{a \in Q_1} \mapsto \sum_{a \in Q_1} [N(a), N(a^*)].$$

- The stack  $\mathfrak{M}_d(\Pi_Q)$  of  $d$ -dimensional  $\Pi_Q$ -modules is equivalent to the quotient  $\mu_d^{-1}(0)/\text{GL}_d$  with  $\text{GL}_d := \prod_{i \in Q_0} \text{GL}_{d_i}(\mathbb{C})$ .

## The coarse moduli space

If we wish to have an algebraic variety instead of a stack, we may consider the coarse quotient:  $X_d(\Pi_Q) := \text{Spec}(\Gamma(\mu_d^{-1}(0))^{\text{GL}_d})$

Points of this space are in bijection with  $d$ -dimensional semisimple  $\Pi_Q$ -modules.



# From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

## Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

# From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

## Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

# From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

## Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

The catch: if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

The catch: if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!



## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

The catch: if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

The catch: if  $d = l d'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $l \geq 2$  then there are no  $d$ -generic stability conditions!

## From moduli spaces to Kac polynomials

- There is a canonical morphism  $JH: \mathfrak{M}(\Pi_Q) \rightarrow X(\Pi_Q)$ ; geometrically this is the affinization map, at the level of points it takes a module to its semisimplification.
- This map is rather stacky, and the spaces very singular. For  $\zeta \in \mathbb{Q}^{Q_0}$  a stability condition, the analogous morphism  $JH_d^\zeta: \mathfrak{M}_d^{\zeta\text{-sst}}(\Pi_Q) \rightarrow X_d^{\zeta\text{-sst}}(\Pi_Q)$  to the GIT moduli space can be better behaved:

### Theorem (Nakajima, CB+VdB)

Assume that  $\zeta$  is  $d$ -generic (equivalently: no strictly semistables). Then

- 1  $JH_d^\zeta$  is a  $\mathbb{C}^*$ -gerbe
- 2  $X_d^{\zeta\text{-sst}}(\Pi_Q)$  is a smooth quasiprojective variety.
- 3 There is an equality  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(H(X_d^{\zeta\text{-sst}}(\Pi_Q), \mathbb{Q}^{\text{vir}}))$ .

**The catch:** if  $d = Id'$  with  $d' \in \mathbb{N}^{Q_0}$  and  $I \geq 2$  then there are *no*  $d$ -generic stability conditions!

# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)}a - a\omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.

# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)} a - a \omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.

# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)} a - a \omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.

# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)}a - a\omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.

# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)} a - a \omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.



# Kac polynomials from DT theory: a sketch

By definition

$$\begin{aligned} a_{Q,\bullet}(q) &= \#^{\text{naive}} \{ \text{abs. indec. } \mathbb{F}_q Q \text{ modules} \} / \cong \\ &\approx \#^{\text{stacky}} \{ \text{abs. indec. } \mathbb{F}_q Q^+ / R_I \text{ modules} \} / \cong \\ &\approx H^{\text{BM}}(\mathfrak{M}(\mathbb{C} Q^+ / R_I), \mathbb{Q}^{\text{vir}}) \end{aligned}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\mathbb{C} \tilde{Q}), \phi_{\text{Tr}(\tilde{W})}) \quad =: \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

$$(\text{by dim. red.}) \cong H(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$$

- Where  $Q^+$  obtained by adding loops  $\omega_i$  at every  $i \in Q_0$  and  $R_I = \langle \omega_{t(a)}a - a\omega_{s(a)} \rangle$
- Where  $\tilde{Q} = \overline{Q}^+$ , and  $\tilde{W} = (\sum_{i \in Q_0} \omega_i)(\sum_{a \in Q_1} [a, a^*])$  and  $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the KS cohomological Hall algebra for the tripled quiver  $\tilde{Q}$  and potential  $\tilde{W}$ .
- $\mathcal{H}_{\tilde{Q}, \tilde{W}}$  is the “critical CoHA” of  $\text{Jac}(\tilde{Q}, \tilde{W}) \cong \Pi_Q[x]$ -modules.

## The Lie algebra $\mathfrak{g}_{\tilde{Q}, \tilde{W}}$

### Theorem (D, Meinhardt)

*(Proved for arbitrary symmetric quivers with potential). There is a subspace  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \subset \mathcal{H}_{\tilde{Q}, \tilde{W}}$ , closed under the commutator Lie bracket, such that the natural map*

$$\text{Sym} \left( \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \otimes H(\mathbb{B}\mathbb{C}^*, \mathbb{Q}) \right) \rightarrow \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

*is an isomorphism.  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$  is called the BPS Lie algebra.*

- 1  $Q$  can have loops.
- 2 Upgrading heuristic argument from previous slide to rigorous mathematics:

### Theorem (D)

*There are equalities  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\tilde{Q}, \tilde{W}, d}^{\text{BPS}})$ .*

## The Lie algebra $\mathfrak{g}_{\tilde{Q}, \tilde{W}}$

### Theorem (D, Meinhardt)

*(Proved for arbitrary symmetric quivers with potential). There is a subspace  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \subset \mathcal{H}_{\tilde{Q}, \tilde{W}}$ , closed under the commutator Lie bracket, such that the natural map*

$$\text{Sym} \left( \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \otimes H(\mathbb{B}\mathbb{C}^*, \mathbb{Q}) \right) \rightarrow \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

*is an isomorphism.  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$  is called the **BPS Lie algebra**.*

- 1  $Q$  can have loops.
- 2 Upgrading heuristic argument from previous slide to rigorous mathematics:

### Theorem (D)

*There are equalities  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\tilde{Q}, \tilde{W}, d}^{\text{BPS}})$ .*

# The Lie algebra $\mathfrak{g}_{\tilde{Q}, \tilde{W}}$

## Theorem (D, Meinhardt)

*(Proved for arbitrary symmetric quivers with potential). There is a subspace  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \subset \mathcal{H}_{\tilde{Q}, \tilde{W}}$ , closed under the commutator Lie bracket, such that the natural map*

$$\text{Sym} \left( \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \otimes H(\mathbb{B}\mathbb{C}^*, \mathbb{Q}) \right) \rightarrow \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

*is an isomorphism.  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$  is called the **BPS Lie algebra**.*

- 1  $Q$  can have loops.
- 2 Upgrading heuristic argument from previous slide to rigorous mathematics:

## Theorem (D)

*There are equalities  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\tilde{Q}, \tilde{W}, d}^{\text{BPS}})$ .*

# The Lie algebra $\mathfrak{g}_{\tilde{Q}, \tilde{W}}$

## Theorem (D, Meinhardt)

*(Proved for arbitrary symmetric quivers with potential). There is a subspace  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \subset \mathcal{H}_{\tilde{Q}, \tilde{W}}$ , closed under the commutator Lie bracket, such that the natural map*

$$\text{Sym} \left( \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \otimes H(\mathbb{BC}^*, \mathbb{Q}) \right) \rightarrow \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

*is an isomorphism.  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$  is called the **BPS Lie algebra**.*

- 1  $Q$  can have loops.
- 2 Upgrading heuristic argument from previous slide to rigorous mathematics:

## Theorem (D)

*There are equalities  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\tilde{Q}, \tilde{W}, d}^{\text{BPS}})$ .*

## The Lie algebra $\mathfrak{g}_{\tilde{Q}, \tilde{W}}$

### Theorem (D, Meinhardt)

*(Proved for arbitrary symmetric quivers with potential). There is a subspace  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \subset \mathcal{H}_{\tilde{Q}, \tilde{W}}$ , closed under the commutator Lie bracket, such that the natural map*

$$\text{Sym} \left( \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}} \otimes H(\mathbb{B}\mathbb{C}^*, \mathbb{Q}) \right) \rightarrow \mathcal{H}_{\tilde{Q}, \tilde{W}}$$

*is an isomorphism.  $\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$  is called the **BPS Lie algebra**.*

- 1  $Q$  can have loops.
- 2 Upgrading heuristic argument from previous slide to rigorous mathematics:

### Theorem (D)

*There are equalities  $a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\tilde{Q}, \tilde{W}, d}^{\text{BPS}})$ .*

# The decomposition theorem

By dimensional reduction  $\mathcal{H}_{\tilde{Q}, \tilde{W}} \cong H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$ .

With respect to the diagram

$$\begin{array}{ccc} \mathfrak{M}(\Pi_Q) & & \\ \downarrow \text{JH} & \searrow p & \\ X(\Pi_Q) & \xrightarrow{r} & \text{pt} \end{array}$$

we write

$$\begin{aligned} H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}}) &:= p_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \\ &= r_* \text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \end{aligned}$$

Theorem (Decomposition theorem for 2CY categories (D))

*There is an isomorphism*

$$\text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \cong \bigoplus_{n \in 2\mathbb{Z}_{\geq 0}} \bigoplus_{s \in S_n} \text{IC}_{\overline{Z_s}}(\mathcal{L}_s)[\dim(Z_s) - n]$$

*with  $\mathcal{L}_s$  simple local systems on smooth locally closed  $Z_s \subset X(\Pi_Q)$ .*

## The decomposition theorem

By dimensional reduction  $\mathcal{H}_{\tilde{Q}, \tilde{W}} \cong H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$ .

With respect to the diagram

$$\begin{array}{ccc} \mathfrak{M}(\Pi_Q) & & \\ \downarrow \text{JH} & \searrow p & \\ X(\Pi_Q) & \xrightarrow{r} & \text{pt} \end{array}$$

we write

$$\begin{aligned} H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}}) &:= p_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \\ &= r_* \text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \end{aligned}$$

Theorem (Decomposition theorem for 2CY categories (D))

*There is an isomorphism*

$$\text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \cong \bigoplus_{n \in 2\mathbb{Z}_{\geq 0}} \bigoplus_{s \in S_n} \text{IC}_{\overline{Z}_s}(\mathcal{L}_s)[\dim(Z_s) - n]$$

*with  $\mathcal{L}_s$  simple local systems on smooth locally closed  $Z_s \subset X(\Pi_Q)$ .*



## The decomposition theorem

By dimensional reduction  $\mathcal{H}_{\tilde{Q}, \tilde{W}} \cong H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$ .

With respect to the diagram

$$\begin{array}{ccc} \mathfrak{M}(\Pi_Q) & & \\ \downarrow \text{JH} & \searrow p & \\ X(\Pi_Q) & \xrightarrow{r} & \text{pt} \end{array}$$

we write

$$\begin{aligned} H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}}) &:= p_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \\ &= r_* \text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \end{aligned}$$

Theorem (Decomposition theorem for 2CY categories (D))

*There is an isomorphism*

$$\text{JH}_* \mathbb{D}Q_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \cong \bigoplus_{n \in 2\mathbb{Z}_{\geq 0}} \bigoplus_{s \in S_n} \text{IC}_{\overline{Z}_s}(\mathcal{L}_s)[\dim(Z_s) - n]$$

*with  $\mathcal{L}_s$  simple local systems on smooth locally closed  $Z_s \subset X(\Pi_Q)$ .*

## The decomposition theorem

By dimensional reduction  $\mathcal{H}_{\tilde{Q}, \tilde{W}} \cong H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}})$ .

With respect to the diagram

$$\begin{array}{ccc} \mathfrak{M}(\Pi_Q) & & \\ \downarrow \text{JH} & \searrow p & \\ X(\Pi_Q) & \xrightarrow{r} & \text{pt} \end{array}$$

we write

$$\begin{aligned} H^{\text{BM}}(\mathfrak{M}(\Pi_Q), \mathbb{Q}^{\text{vir}}) &:= p_* \mathbb{D}\mathbb{Q}_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \\ &= r_* \text{JH}_* \mathbb{D}\mathbb{Q}_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \end{aligned}$$

## Theorem (Decomposition theorem for 2CY categories (D))

*There is an isomorphism*

$$\text{JH}_* \mathbb{D}\mathbb{Q}_{\mathfrak{M}(\Pi_Q)}^{\text{vir}} \cong \bigoplus_{n \in 2\mathbb{Z}_{\geq 0}} \bigoplus_{s \in S_n} \text{IC}_{\overline{Z_s}}(\mathcal{L}_s)[\dim(Z_s) - n]$$

*with  $\mathcal{L}_s$  simple local systems on smooth locally closed  $Z_s \subset X(\Pi_Q)$ .*

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathrm{JH}_* \mathbb{D}Q^{\mathrm{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau_{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\mathrm{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

### Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}})$ .

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathbb{J}H_* \mathbb{D}Q^{\text{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau_{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\text{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

### Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}})$ .

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathbb{H}\mathbb{H}_* \mathbb{D}Q^{\text{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := p_{\tau}^{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\text{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

### Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\text{BPS}})$ .

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathrm{JH}_* \mathbb{D}\mathbb{Q}^{\mathrm{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau_{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\mathrm{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

### Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}})$ .

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathrm{JH}_* \mathbb{D}\mathbb{Q}^{\mathrm{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau^{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\mathrm{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

### Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}})$ .

# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathrm{JH}_* \mathbb{D}\mathbb{Q}^{\mathrm{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau^{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\mathrm{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

## Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}})$ .



# A sheaf-theoretic Hall algebra

## Convolution tensor product for sheaves on the coarse moduli space

The coarse moduli space  $X(\Pi_Q)$  is a commutative monoid: the morphism  $m: X(\Pi_Q)^{\times 2} \rightarrow X(\Pi_Q)$  sends a pair of points representing modules  $N, N'$  to the point representing  $N \oplus N'$ . For  $\mathcal{F}, \mathcal{G} \in \mathcal{D}^+(X(\Pi_Q))$  two complexes of constructible sheaves, we define

$$\mathcal{F} \boxtimes \mathcal{G} := m_*(\mathcal{F} \boxtimes \mathcal{G}).$$

The complex  $\mathcal{A}_{\Pi_Q} := \mathrm{JH}_* \mathbb{D}\mathbb{Q}^{\mathrm{vir}}$  carries an algebra structure in the tensor category  $(\mathcal{D}^+(X(\Pi_Q)), \boxtimes)$ . The perverse sheaf/MHM  $\mathcal{A}_{\Pi_Q}^0 := {}^p\tau^{\leq 0} \mathcal{A}_{\Pi_Q}$  is an algebra object in  $(\mathrm{Perv}(X(\Pi_Q)), \boxtimes)$  with semisimple underlying object.

## Theorem (D)

- There is Lie subobject  $\mathcal{L} \subset \mathcal{A}_{\Pi_Q}^0$  with  $H(X(\Pi_Q), \mathcal{L}) \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}}$
- There is an isomorphism of algebras  $H(X(\Pi_Q), \mathcal{A}_{\Pi_Q}^0) \cong U(\mathfrak{g}_{\tilde{Q}, \tilde{W}}^{\mathrm{BPS}})$ .

# The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- *Real roots:*  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- *Isotropic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- *Hyperbolic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

In words: *all* coefficients of *all* Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.

# The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- *Real roots:*  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- *Isotropic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- *Hyperbolic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

In words: *all* coefficients of *all* Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.

## The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

### Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- *Real roots:*  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- *Isotropic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- *Hyperbolic roots:*  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

*In words: all coefficients of all Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.*

# The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

## Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- **Real roots:**  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- **Isotropic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- **Hyperbolic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

*In words: all coefficients of all Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.*

# The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

## Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- **Real roots:**  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- **Isotropic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- **Hyperbolic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

In words: all coefficients of all Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.

## The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

### Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- **Real roots:**  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- **Isotropic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- **Hyperbolic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

In words: all coefficients of all Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.

# The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

## Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- **Real roots:**  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- **Isotropic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- **Hyperbolic roots:**  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

*In words: all coefficients of all Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.*



## The main result

Analyzing the possible summands appearing in the decomposition of  $\mathcal{L}$ , we prove the following, for general  $Q$  (loops allowed):

### Theorem (D, Hennecart, Schlegel Mejia)

Let  $Q$  be a quiver. Let  $\Phi_+$  be the set of positive roots of  $\mathfrak{g}_Q$ . We define

- *Real roots*:  $\mathcal{G}_d = \mathbb{Q}$  if  $d = 1_i$ , and no loops at  $i$ .
- *Isotropic roots*:  $\mathcal{G}_d = \mathrm{IH}^*(X_{d'}(\Pi_Q), \mathbb{Q})$  if  $d$  is isotropic and  $d = Id'$  with  $d'$  indivisible.
- *Hyperbolic roots*:  $\mathcal{G}_d = \mathrm{IH}^*(X_d(\Pi_Q), \mathbb{Q})$  if  $d$  is hyperbolic.

Let  $\mathfrak{g}_{\mathcal{G}}$  be the GKM algebra generated by  $\bigoplus_{d \in \Phi_+} \mathcal{G}_d$ . Then  $\mathfrak{g}_{\mathcal{G}} \cong \mathfrak{g}_{\tilde{Q}, \tilde{W}}$  and so

$$a_{Q,d}(q^{-1}) = \chi_{q^{1/2}}(\mathfrak{g}_{\mathcal{G},d})$$

In words: **all** coefficients of **all** Kac polynomials are given by the dimensions of graded pieces of a generalised Kac–Moody Lie algebra with Chevalley generators identified with intersection cohomology of singular Nakajima quiver varieties.

Happy birthday Victor!

and grazie mille all!

Happy birthday Victor!  
and grazie mille all!