Confusion matrix 100% baretsshortsegment 12.5% 1 25.0% 12.5% retroflex-stomach -86.1% 99/115 0.9% 0.9% 1.7% 10.4% 2 12 uleerative-colitis-0-1 -33.3% 16.7% 16.7% 2/6 urerative collis grade 3 -65.0% 13/20 30.0% 5.0% rauvesophagitished - 20.5% 1 15.4% 6/39 2.6% 2.6% 1 dyedresection-margins -18.8% 28 - 75% 0.0% 0/1 normalizhine - 3.6% 78.6% 14.3% 3.6% 20 esophaditista - 14.8% 37.7% 47.5% 23 29/61 ukerative colitis 1.2 -0.0% 0/2 28.6% 14.3% True labels bb05.2.3 -4.0% 8.7% - 50& ileum -0.0% 0/2 99.0% 96/97 1.0% impacted stool -100.0% 20/20 94.1% 143/15 0.7% 0.7% 0.7% 3.3% 0.7% 5 urerstive collis grade 2 -44.8% 30/67 1.5% 16.4% 1.5% 6.0% 20.9% 1.5% ukerative colitis 23 -- 20% 0.0% 0/5 20.0% 20.0% 0.7% 98.0% 147/15 retroflex rectum -98.3% 58/59 1.7% ucefative colitis grade 1 -22.6% 41.9% 7 13/31 16.1% 12.9% 6.5% 0.6% 1.9% 0.6% 1.3% 3.2% 1.9% 87.1% 1.9% 1 3 1 2 5 3 135/155 3 0.6% 7.3% 11 ulcerative-colitis-grade-3 dyedresectionmargins ulcerative colitis grade 2 ulea dive colitis quade 1 - 0% ulcerative-colitiss 2.3 retroflex stomach ulcetative colitisco.1 ulcerative-colitise 1.2 dyedlifted polyps hemormoids normal Ziline esophaghtis a Predicted labels