FEDERAL UNIVERSITY OF PARANÁ

HENRIQUE APARECIDO LAUREANO

CURITIBA

2021

HENRIQUE APARECIDO LAUREANO

MODELING THE CUMULATIVE INCIDENCE FUNCTION OF CLUSTERED COMPETING RISKS DATA: A MULTINOMIAL GLMM APPROACH

Thesis presented to the Graduate Program of Numerical Methods in Engineering, Concentration Area in Mathematical Programming: Statistical Methods Applied in Engineering, Federal University of Paraná, as part of the requirements to the obtention of the Master's Degree in Sciences.

Supervisor: Prof. PhD Wagner Hugo Bonat Co-supervisor: Prof. PhD Paulo Justiniano Ribeiro Jr

MODELING THE CUMULATIVE INCIDENCE FUNCTION OF CLUSTERED COMPETING RISKS DATA: A MULTINOMIAL GLMM APPROACH

HENRIQUE APARECIDO LAUREANO

MODELING THE CUMULATIVE INCIDENCE FUNCTION OF CLUSTERED COMPETING RISKS DATA: A MULTINOMIAL GLMM APPROACH

Thesis presented to the Graduate Program of Numerical Methods in Engineering, Concentration Area in Mathematical Programming: Statistical Methods Applied in Engineering, Federal ences. University of Paraná, as part of the requirements to the obtention of the Master's Degree in Sci-

Master thesis approved. XXX XX, 2021.

Prof. PhD Wagner Hugo Bonat Supervisor

Prof. PhD Paulo Justiniano Ribeiro Jr Co-supervisor

Prof. PhD ...

Internal Examinator - PPGMNE

Prof. PhD ...

Internal Examinator - PPGMNE

External Examiner -Prof. PhD ...

CURITIBA

2021

ACKNOWLEDGEMENTS

As Moro once said, I am thankful for everything and everyone.

and trajectory time levels are correlated. In scenarios with high CIF the model exhibits latent-effects; and 4) an automatic differentiation (AII) routine, the state-of-the-art in derivatives computation. To check the estimability of our model a large simulation the beffer results, but still with an excessive variance, showing that improvements are difficult estimation, with our results converging to a latent structure where the risk to verify which one is most adequate to real scenarios. The model presents to be of study is performed, based on different latent structure formulations, with the aim linear algebra libraries; 3) efficient Laplace approximation implementation for the package, we have 1) the log-likelihood function written in C++; 2) access to efficient model builder (TMB) (KRISTENSEN et al., 2016). With this R (R Core Team, 2021) the model parameters estimation the most efficient as possible, we use the template time level function. The latent effects are inserted in those level functions. To make via CIF, modeled here following Cederkvist et al. (2019) specification, based on its by the parameters of its covariance matrix. The probability distributions are connected effects are accommodated via a multivariate Gaussian distribution and are modeled responsible for the occurrence of an event, a failure; and 2) censorship, when the event of interest happens or not for none of the competing causes, in the study period. To handle this type of data, we propose a generalized linear mixed model (GLMM) in latent-effects framework, instead of a usual according to be the one when the occurrence of the competing causes, in the study period. To latent-effects framework, instead of a usual according to be the one when the occurrence of the causes/variables competing to be the one of interest happens or not for none of the competing causes, in the study period. To decomposition as the product of an instantaneous risk level function with a trajectory for the competing causes and censorship, conditioned on the latent effects. The latent competing cause. In our framework, we suppose a multinomial probability distribution in the probability scale, in terms of the cumulative incidence function (CIF) of each sometimes intractable. We, on the other hand, model the clustered competing causes dependence accommodation ends by generating a complicated likelihood function, the modeling is usually done by means of the hazard rate, and the within-cluster Clustered competing risks data is a special case of failure time data. Besides the cluster

In Link Wilne JM 100 oh oh mulou - James lander of CIF alta o modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresenta os melhores resultados, mas ainda com uma excessiva mulos de para modelo apresentados de para modelo a

IENSEN et al., 2016). Com este pacote R (R Core Team, 2021), temos 1) a função d. implementação eficiente da aproximação de Laplace para os efeitos latentes; e 4) uma de derivadas. Para verificar a estimabilidade do rosso modelo e realizada.

o obietiva do possível, usamos o template model builder (TMB) (KRI TENSEN et al., 2016). Com este pacote R (R Core Team, 2021), temos 1) a função d. implementação eficiente da aproximação de Laplace para os efeitos latentes; e 4) uma de derivadas. Para verificar a estimabilidade do rosso modelo e realizada.

o obietiva do possível, usamos o template model builder (TMB) (KRI TENSEN et al., 2016). Com este pacote R (R Core Team, 2021), temos 1) a função d. implementação de ficientes bibliotecas de álgebra linear; 3, o estado da arte em computação de de derivadas. Para verificar a estimabilidade do rosso modelo e realizada. o objetivo de verificar qual delas é a mais adequada a um cenário real. O modelo se latente onde os níveis de risco e de trajetória estão correlacionados. Em cenários de apresenta de difícil estimatição, com nossos resultados convergindo para uma estrutura função de nível de risco instantâneo com uma função de nível de tempo de trajetória. em Cederkvist et al. (2019), com base em sua decomposição como o produto de uma probabilidade são conectadas por meio da CIF, modeladas aqui seguindo a especificação e são modelados via os parâmetros de sua matriz de covariância. As distribuições de efeitos latentes são acomodados por meio de uma distribuição Gaussiana multivariada nomial para as causas competidoras e censura, condicionado aos efeitos latentes. Os competidora. Em nossa modelagem, supomos uma distribuição de probabilidade multilidada, por meio da função de incidência acumulada (CIF, em inglês) de cada causa por outro lado, modelamos as causas competidoras agrupadas na escala da probabiacaba por gerar uma complicada função de verossimilhança, às vezes intratável. Nós, usualmente feita por meio da taxa de risco, e a acomodação da dependência intra-grupo de um modelo de sobrevivência usual. Em análise de sobrevivência, a modelagem é generalizado misto (GLMM), ou seja, um modelo de efeitos latentes/aleatórios, em vez no período de estudo. Para lidar com este tipo de dado, propomos um modelo linear sura, quando o evento de interesse ocorre ou não por nenhuma das causas concorrentes, competindo para ser a responsável pela ocorrência de um evento, uma falha; e 2) centro de interesse ocorre ou não por nenhuma das causas concorrentes, tre seus elementos, esse tipo de dado é caracterizado por 1) múltiplas causas/variáveis falha. Além da estrutura de grupo que implica uma dependência latente intra-grupo en Dados de riscos competitivos agrupados são um caso especial de dados de tempo de

Keywords: Clustered competing risks. Within-cluster dependence. Multinomial gener- alized linear mixed model (GLMM). TMB: Template Model Builder. Laplace approxima- Laplace approxima-

LIST OF FIGURES

46	Equation 4.1
	ATES, LATENT EFFECTS IN ZERO, AND FIXED EFFECTS IN
	TWO COMPETING CAUSES OF FAILURE, WITHOUT COVARI-
	TIVE DERIVATIVES (dCIF) W.R.T. TIME FOR A MODEL WITH
	FIGURE 10 - CUMULATIVE INCIDENCE FUNCTIONS (CIF) AND RESPEC-
43	COMPONENTS, IN B)
	THE VARIANCE COMPONENTS, IN A), AND CORRELATION
	FIGURE 9 - ILLUSTRATION OF THE PARAMETRIZATION BEHAVIOR FOR
39	TENT EFFECTS u_1 AND η_1
	THE VARIATION BETWEEN FRAMES IS GIVEN BY THE LA-
	CONFIGURATION: $\beta_1 = -2$, $\beta_2 = -1$, $\gamma_1 = 1$, $w_1 = 3$ AND $u_2 = 0$.
	OF FAILURE, WITHOUT COVARIATES AND THE FOLLOWING
	et al. (2019), IN A MODEL WITH TWO COMPETING CAUSES
	TIVE INCIDENCE FUNCTION (CIF), PROPOSED BY Cederkvist
	FIGURE 8 - ILLUSTRATION OF A GIVEN CLUSTER-SPECIFIC CUMULA-
38	ZERO, WITH THE EXCEPTION OF $w_1 = 1$
	IN EACH SCENARIO ALL OTHER COEFFICIENTS ARE SET TO
	THE FOLLOWING CONFIGURATION: $\beta_2 = 0$, $u = 0$ AND $\eta = 0$;
	CAUSES OF FAILURE, WITHOUT COVARIATES, AND WITH
	Cederkvist et al. (2019)), IN A MODEL WITH TWO COMPETING
	CUMULATIVE INCIDENCE FUNCTION (CIF) (PROPOSED BY
	FIGURE 7 - ILLUSTRATION OF COEFFICIENT BEHAVIORS FOR A GIVEN
33	FIGURE 6 - TMB PACKAGE DESIGN
33	MODEL WRITTEN IN TMB
	FIGURE 5 $-$ R CODE FOR THE MODEL FITTING OF A LOGISTIC MIXED
32	MIXED MODEL
	FIGURE 4 - R CODE FOR THE TMB IMPLEMENTATION OF A LOGISTIC
28	FIGURE 3 - EXAMPLE OF A SIMPLE COMPUTATIONAL GRAPH
27	FIGURE 2 - A COMPUTATIONAL GRAPH
15	ILNESS-DEATH MODEL, THE SIMPLEST MULTISTATE MODEL
	FIGURE 1 - ILLUSTRATION OF MULTISTATE MODELS FOR A A) FAIL-

92	FIGURE 34 – PARAMETER γ_1 BIAS WITH 2.5% AND 97.5% QUANTILES
91	FIGURE 33 – PARAMETER eta_2 BIAS WITH 2.5% AND 97.5% QUANTILES \dots
91	FIGURE 32 – PARAMETER eta_1 BIAS WITH 2.5% AND 97.5% QUANTILES \dots
67	SIXTY-THOUSAND DATA POINTS
	MAP IN THE SCENARIO OF CLUSTER SIZE 10, HIGH CIF, AND
	FIGURE 31 COMPLETE MODEL'S PARAMETERS CORRELATION HEAT-
66	OF HIGH CIF AND 60 THOUSAND DATA POINTS
	FIGURE 30 – CORRELATION PARAMETERS DENSITIES IN THE SCENARIOS
65	HIGH CIF AND 60 THOUS
	E PARAMETERS DENSITIES IN THE SCENARIOS (
හ	INCIDENCE FONCTION
62	3
	FIGURE 27 – HIGH CUMULATIVE INCIDENCE FUNCTION (CIF) SCENARIO
61	STANDARD DEVIATIONS
	FIGURE 26 – PARAMETERS $\{z(ho_{13}),z(ho_{24}),z(ho_{14}),z(ho_{23})\}$ BIAS WITH \pm 1.96
60	FIGURE 25 – PARAMETER $z(ho_{34})$ BIAS WITH \pm 1.96 STANDARD DEVIATIONS
60	FIGURE 24 – PARAMETER $z(\rho_{12})$ BIAS WITH \pm 1.96 STANDARD DEVIATIONS
59	FIGURE 23 – PARAMETER $\log(\sigma_4^2)$ BIAS WITH \pm 1.96 STANDARD DEVIATIONS
59	FIGURE 22 – PARAMETER $\log(\sigma_3^2)$ BIAS WITH \pm 1.96 STANDARD DEVIATIONS
58	FIGURE 21 – PARAMETER $\log(c_2^2)$ BIAS WITH ± 1.96 STANDARD DEVIATIONS
58	FIGURE 20 – PARAMETER $\log(c_1^2)$ BIAS WITH \pm 1.96 STANDARD DEVIATIONS
57	FIGURE 19 – PARAMETER w_2 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
57	FIGURE 18 – PARAMETER w_1 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
56	FIGURE 17 – PARAMETER γ_2 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
56	FIGURE 16 – PARAMETER γ_1 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
55	FIGURE 15 – PARAMETER eta_2 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
55	FIGURE 14 – PARAMETER eta_1 BIAS WITH \pm 1.96 STANDARD DEVIATIONS .
50	WITH A COMPLETE 4x4 LATENT STRUCTURE
	FIGURE 13 — C++ CODE FOR THE TMB IMPLEMENTATION OF A multiGLMM
48	COVARIATES, AND LATENT EFFECTS IN ZERO
	WITH TWO COMPETING CAUSES OF FAILURE, WITHOUT
	FIGURE 12 – CUMULATIVE INCIDENCE FUNCTIONS (CIF) FOR A MODEL
47	TIONS SPECIFIED IN Equation 4.1 AND Equation 4.2
	GORITHM 1 GUIDELINES WITH PARAMETER CONFIGURA-
	CLUSTERS OF SIZE TWO. THE SIMULATION FOLLOWED AL-
	DEL WITH TWO COMPETIN
	SPECTIVE OUTPUT PCERCENTAGES AND FAILURE TIMES
	FIGURE 11 – HISTOGRAMS FOR SIMULATED PROBABILITIES WITH RE-

97	AND 97.5% QUANTILES
	IGURE 44 – PARAMETERS $\{z(\rho_{13}), z(\rho_{24}), z(\rho_{14}), z(\rho_{23})\}$ BIAS WITH 2.5%
96	IGURE 43 – PARAMETER $z(ho_{34})$ BIAS WITH 2.5% AND 97.5% QUANTILES .
96	IGURE 42 – PARAMETER $z(\rho_{12})$ BIAS WITH 2.5% AND 97.5% QUANTILES .
95	IGURE 41 – PARAMETER $\log(\sigma_4^2)$ BIAS WITH 2.5% AND 97.5% QUANTILES
95	IGURE 40 – PARAMETER $\log(\sigma_3^2)$ BIAS WITH 2.5% AND 97.5% QUANTILES
94	IGURE 39 – PARAMETER $\log(\sigma_2^2)$ BIAS WITH 2.5% AND 97.5% QUANTILES
94	IGURE 38 – PARAMETER $\log(\sigma_1^2)$ BIAS WITH 2.5% AND 97.5% QUANTILES
93	IGURE 37 - PARAMETER w_2 BIAS WITH 2.5% AND 97.5% QUANTILES
93	IGURE 36 – PARAMETER w_1 BIAS WITH 2.5% AND 97.5% QUANTILES
92	FIGURE 35 – PARAMETER γ_2 BIAS WITH 2.5% AND 97.5% QUANTILES

LIST OF ALGORITHMS

COMPETIN	ALGORITHM 1
MPETING RISKS DATA	SIMULATING FROM A multiGLMM FOR CLUSTERED
45	

CONTENTS

72	BIBLIOGRAPHY	
71	FUTURE WORKS	6.1
88	FINAL CONSIDERATIONS	6
53	SIMULATION STUDY	5.1
53	RESULTS	Ch
48	SIMULATION STUDIES DESIGN	4.2
45	SIMULATING FROM THE MODEL	4.1
45	SIMULATION STUDY PALASETS	4
41	Parametrization	3.2.1
39	MODEL SPECIFICATION	3.2
35	CLUSTER-SPECIFIC CUMULATIVE INCIDENCE FUNCTION (CIF)	3.1
35	RISKS DATA	
	multiglmm: A MULTINOMIAL GLMM FOR CLUSTERED COMPETING	ω
30	TMB: TEMPLATE MODEL BUILDER	2.5
29	Reverse Mode	2.4.2
28	Forward Mode	2.4.1
27	AD: AUTOMATIC DIFFERENTIATION	2.4
24	OPTIMIZATION: MARGINAL LIKELIHOOD FUNCTION	2.3
21	MARGINALIZATION: LAPLACE APPROXIMATION AND ALTERNATIVES	2.2
20	FORMULATION: OBTAINING A JOINT LIKELIHOOD FUNCTION	2.1
20	TION, AND IMPLEMENTATION	
	GENERALIZED LINEAR MIXED MODELS: FORMULATION, OPTIMIZA-	Ν
19	THESIS ORGANIZATION	1.4
18	LIMITATION	1.3
18	JUSTIFICATION	1.2
17	Specific goals	1.1.2
17	General goals	1.1.1
17	GOALS	<u>:</u>
14	INTRODUCTION	-

91	QUANTILES	
	APPENDIX E - MODEL PARAMETERS BIAS WITH 2.5% AND 97.5%	
88	R CODE SHOWING HOW TO LOAD AND FIT THE multiGLMM VERSIONS.	D.4
87	C++ CODE FOR THE TMB IMPLEMENTATION OF A multiGLMM WITH A	D.3
85	2×2 LATENT STRUCTURE ON THE TRAJECTORY TIME LEVEL $ \ldots \cdot $	
	C++ CODE FOR THE TMB IMPLEMENTATION OF A multiGLMM WITH A	D.2
84	2×2 LATENT STRUCTURE ON THE RISK LEVEL \dots	
	C++ CODE FOR THE TMB IMPLEMENTATION OF A multiglimm WITH A	D.1
84	APPENDIX D - C++ CODES FOR THE TMB IMPLEMENTATION OF THE	
82	FOR MORE INFORMATION CHECK SECTION 4.1	
}	COMPETING CAUSES AND CLUSTERS OF SIZE TWO.	
	APPENDIX C - R CODE TO SIMULATE FROM A multiGLMM WITH TWO	
78	RISKS DATA	
	THE JOINT LOG-LIKELIHOOD FUNCTION OF THE MULTINOMIAL GLMM FOR CLUSTERED COMPETING	
	CTS	
77	RISKS DATA	
	MULTINOMIAL GLMM FOR CLUSTERED COMPETING	
	THE JOINT LOG-LIKELIHOOD FUNCTION OF THE	
	APPENDIX A – ANALYTIC GRADIENT OF THE LATENT EFFECTS FOR	
77	APPENDIX	
76	APPENDIX	

Consider a cluster of random variables representing the time until the occurrence of some event. These random variables are assumed to be correlated, i.e. for some biological or environmental reason it is not adequate to assume independence between them. Also, we may be interested in the occurrence of not only one specific event, having in practice a competition of events to see which one happens first, if it happens. Such events may also be of low probability albeit severe consequences, this is the moment when the cluster correlation makes its difference: the occurrence of an event in a cluster member should affect the probability of the same happening in the others.

A realistic context that fits perfectly with the framework described above is the study of disease incidence in family members, where each member is indexed by a random variable and each cluster consists of a familiar structure. The inspiration to the study of these kinds of problems came from the work developed in Cederkvist et al. (2019), where they studied breast cancer incidence in mothers and daughters but using a complicated modeling framework. Based on that, the aim of this thesis is to propose a simpler framework taking advantage of several state-of-art computational libraries and see how far we can go in several scenarios. Until now we have just contextualized, we still need to introduce the methodology. To do this, some definitions and theoretical contexts are welcome.


When the object under study is a random variable representing the time until some event occurs, we are in the field of failure time data (KALBFLEISCH; PRENTICE, 2002). The occurrence of an event is generally denoted failure, and major areas of application are biomedical studies and industrial life testing. In this thesis, we maintain our focus on the former As common in science, same methodologies can receive different names depending on the area. In industrial life testing is performed what is called a reliability analysis; in biomedical studies is performed what is called survival analysis. Generally, the term survival is applied when we are interested in the occurrence of only one event, a failure time process. When we are interested in the occurrence of more than one event we enter in the yard of competing risks and multistate models. A visual aid is presented on Figure 1 and a comprehensive reference is Kalbfleisch & Prentice (2002).

Failure time and competing risks processes may be seen as particular cases of a multistate model. Besides the number of events (states) of interest, the main difference between a multistate model and its particular cases is that only in the multistate scenario we may have transient states, using a *stochastic process* language. In the particular cases, all states besides the initial state 0, are absorbents - once you reached it you do not leave.

The simplest multistate model that exemplify this behavior is the illness-death model, Figure 1 C), where a patient (initially in state 0) can get sick (state 1) or die (state 2); if sick it can recover (returns to state 0) or die. We work in this thesis only with competing risks processes, and for each patient we need the time (age) until the occurrence, or not, of the event.

15

FIGURE 1 – ILLUSTRATION OF MULTISTATE MODELS FOR A A) FAILURE TIME PRO-CESS; B) COMPETING RISKS PROCESS; AND C) ILNESS-DEATH MODEL, THE SIMPLEST MULTISTATE MODEL



SOURCE: The author (2021).

When for some known or unknown reason we are not able to see the occurrence the period of follow up the patient may not get sick or die, staying at state 0. This is denoted *right-censorship*; if a patient is in state 1 at the end of the study, we are *censored* to see him reaching the state 2 or returning to state 0. This is the inherent idea to censorship and must be present in the modeling framework, thus arriving in the so-called *survival models* (KALBFLEISCH; PRENTICE, 2002).

A survival model deals with the survival experience. Usually, the survival experience is modeled in the *luzard* (failure rate) scale and it can be expressed for a subject *i* as

$$\lambda(t \mid x_i) = \lambda_0(t) \times c(x_i\beta)$$
 at time t , (

i.e. as the product of an arbitrary baseline hazard function $\lambda_0(\cdot)$, with a specific function form $c(\cdot)$, that will depend on the probability distribution to be chosen for the failure time and on predictors/covariates/explanatory/independent variables $x_i = [x_1 \dots x_p]$, where $\beta^\top = [\beta_1 \dots \beta_p]$ is the parameters vector.

This structure is specified for a failure time process, as in Figure 1 A). Nevertheless, the idea is easy to extend. We basically have the Equation 1.1's model to each cause-specific (in a competing risks process) or transition (in a multistate process). A complete and extensive detailing can be, again, found in Kalbfleisch & Prentice (2002).

an op. NALAINTON; STALLARD, 1979). In its simplest form, a frailty model (CLAYTON, 1978; an unobserved random proportionality factor that modifies the hazard function of an individual, or of related individuals. Frailty models are extensions of Equation 1.1's model.

In the competing risks setting, the hazard scale (focusing on the cause-specific hazard) is not the only possible scale to work on. A more attractive possibility components of the only scale (ANDERSEN et al., 2017). cluster/group/family dependence that needs to be considered, accommodated, and cause-specific structure, we have to deal with the fact that the events are happening modeled. This, possible, dependence is something that we do not actually measure but in related individuals. This configures what is denoted family studies, i.e. we have a In this work we approach the case of clustered competing risks. Besides the

attractive and logical choice. Since the CIF plays a central role in this master thesis, it studies there is often a strong interest in describing age at disease onset, which is directly will be formally defined later in a place with greater emphasis. With the definitions and described by the cause-specific CIF. Therefore, making the probability scale a more the theoretical context being made, let us be more specific

pairwise approach since we need to add model layers to be able to handle with the about a joint distribution specification, which generally translates also into a compucostly or even inviable. In failure time data problems, the composite likelihood function is built from the product of marginal densities. The marginal specification implies a more complicated, besides the number of small details to workaround from the fact of dependence structure. A clear advantage of this approach is that we do not need to care analysis in high-dimensional situations when a full approach is too computational of the cause-specific CIF as the product of a cluster-specific risk level function with a COX; REID, 2004; VARIN; REID; FIRTH, 2011) is a valid alternative to a full likelihood cluster-specific failure time trajectory function. A composite approach (LINDSAY, 1988, being working with not an exact likelihood function. (2019) proposed a pairwise composite likelihood approach based on the factorization ture allowing for within-cluster dependence of both risk and timing, Cederkvist et al tational advantage. A disadvantage is the model specification, which becomes much To work with competing risks data on the probability scale plus a latent struc-

not viable here, so we try to reach the same goal of Cederkvist et al. (2019) albeit with a simpler framework taking advantage of state-of-art software, something still We do not have any guarantees that a full likelihood inference procedure is

> optimize it. A Fisherian approach per se. an appropriate latent effects structure), marginalize (integrate out the latent effects) and generalized linear mixed model (GLMM). Instead of concentrating on failure time multinomial model with its link function based on the cluster-specific CIF, accouting for or using a composite approach, we just build the joint/full likelihood function (a data and consequently having a survival/frailty model based on the hazard scale, not so common in the statistical modeling community. This simpler framework is a

2001), consider a random suject i. In a standard linear model we assume that the (GLM) and a comprehensive reference is McCullagh & Nelder (1989) distributions. This extended framework received the name of generalized linear model (dichotomic data), gamma (continuous but positive) and Gaussian (continuous data) are useful for practical modelling, such as the Poisson (for counting data), binomial this idea is generalized to distributions of the exponential family. Many of its members much well explained in Nelder & Wedderburn (1972), with the aid of a link function $g(\cdot)$. bution and what we do is to model its mean, $\mu_i \equiv \mathbb{E}(Y_i \mid x_i)$, via a linear combination. As response variable Y_i , conditioned on the covariates x_i , follows a normal/Gaussian distri To a better contextualization of our GLMM approach (MCCULLOCH; SEARLE,

tion of a latent effect u (then, mixed) into the mean structure. The mean structure of a What makes a GLM into a GLMM (MCCULLOCH; SEARLE, 2001) is the addi-

$$g(\mu_i) = x_i \beta + z_i u$$
, $u \sim \text{Multivariate Normal}(0, \Sigma)$

are into x_i , the $i^{ ext{th}}$ vector row of a model-matrix X, with $oldsymbol{eta}$ being a vector of unknown mean structure is made through the i^{th} vector row of a design-matrix Z. The covariates zero mean and a parametrized variance-covariance matrix Σ . Its correct linkage to the where the latent effect is assumed to follow a multivariate Gaussian distribution of

es a Nilw of the July of the J (CIF) of clustered competing risks data. Propose and study the estimability of a multinomial generalized linear mixed multinomial generalized linear mixed re estimability of manimum alklihood

1.1.2 Specific goals

1. Simulate from the model, i.e. generate synthetic data to study statistical properties.

- Write the model in the Template Model Builder (TMB) software, developed by Kristensen et al. (2016) and possibly the most efficient likelihood-based way of doing such task.
- Take advantage of TMB's functionalities with special attention to the computation of gradients and Hessians via a state-of-art automatic differentiation (AD) implementation; and a joint likelihood marginalization via an efficient Laplace approximation routine.
- 4. Study the <u>model identifiability</u> through the proposition of different complexity level models in terms of parametric space and latent effect structures.
- 5. Make exact likelihood-based inference to the cluster and cause-specific CIF of clustered competing risks data.

1.2 JUSTIFICATION

In the biomedical statistical modeling literature, the study of disease occurrence in related individuals receives the name of family studies. Key points of interest are the within-family dependence and determining the role of different risk factors. The within-family dependence may reflect both disease heritability and the impact of shared environmental effects. The role of different risk factors arrives in the class of multivariate models, which options are limited in the statistical literature. Thus, the number of statistical models for competing risks data that accommodate the within-cluster/family dependence is even more limited. Some modeling options are briefly commented in Cederkvist et al. (2019), with his pairwise composite approach being proposed as a new and better option to model the cause-specific cumulative incidence function (CIF), describing age at disease onset, of clustered competing risks data on the probability scale. We propose to model the cause-specific CIF and accommodate the within-family dependence in the same fashion (via a latent structure that allows the absolute risk and the failure time distribution to vary between families) but with an easier framework, based on a multinomial generalized linear mixed model approach.

.3 LIMITATION

This work restraint to the proposition and model identifiability study of a multinomial model for the cause-specific cumulative incidence function (CIF) of competing risks data, with a latent effect structure to accommodate within-family dependence with regard to both risk and timing. Given its considerable model complexity, hypothesis tests; residual analysis; and good-of-fit measures are not contemplated.

Inche quel & response are in contraction of grands and an act of the formal strands and charter a grands

1.4 THESIS ORGANIZATION

19

This master thesis contains 6 chapters including this introduction. Chapter 2 presents a systematic review of the main aspects involved in the formulation, optimzation, and implementation of a generalized linear mixed model (GLMM). Given the modeling framework overview, Chapter 3 presents our multinomial GLMM (multiGLMM) to model the cause-specific cumulative incidence function (CIF) of clustered competing risks data. In Chapter 4 we describe the simulation procedure to generate synthetic data and present some model particularities. In Chapter 5 the obtained results are presented, and in Chapter 6 we discuss the contributions of this thesis and present some suggestions for future work.

Internaçãos nãos neusociamente roba de mátodos de mátodos de metrociamente roba de mátodos.

Memos que ve chejave e conclusión que ve não de mátodos de mátodos de mátodos de mátodos de mátodos para emplica que entrados para de mátodos para de mátodos que emplica que ve não de modulos de aprecionados para de mátodos de materiales de mátodos de materiales de mátodos de materiales de materiales