

5512. Count Unhappy Friends

[My Submissions \(/contest/weekly-contest-206/problems/count-unhappy-friends/submissions/\)](/contest/weekly-contest-206/problems/count-unhappy-friends/submissions/)

[Back to Contest \(/contest/weekly-contest-206/\)](/contest/weekly-contest-206/)

You are given a list of preferences for n friends, where n is always even.

For each person i , `preferences[i]` contains a list of friends **sorted** in the **order of preference**. In other words, a friend earlier in the list is more preferred than a friend later in the list. Friends in each list are denoted by integers from 0 to $n-1$.

All the friends are divided into pairs. The pairings are given in a list `pairs`, where `pairs[i] = [xi, yi]` denotes x_i is paired with y_i and y_i is paired with x_i .

However, this pairing may cause some of the friends to be unhappy. A friend x is unhappy if x is paired with y and there exists a friend u who is paired with v but:

- x prefers u over y , and
- u prefers x over v .

Return the number of unhappy friends.

Example 1:

Input: $n = 4$, `preferences = [[1, 2, 3], [3, 2, 0], [3, 1, 0], [1, 2, 0]]`, `pairs = [[0, 1],`

Output: 2

Explanation:

Friend 1 is unhappy because:

- 1 is paired with 0 but prefers 3 over 0, and
- 3 prefers 1 over 2.

Friend 3 is unhappy because:

- 3 is paired with 2 but prefers 1 over 2, and
- 1 prefers 3 over 0.

Friends 0 and 2 are happy.

Example 2:

Input: $n = 2$, `preferences = [[1], [0]]`, `pairs = [[1, 0]]`

Output: 0

Explanation: Both friends 0 and 1 are happy.

Example 3:

User Accepted: 0

User Tried: 0

Total Accepted: 0

Total Submissions: 0

Difficulty: Medium

Input: $n = 4$, preferences = $[[1, 3, 2], [2, 3, 0], [1, 3, 0], [0, 2, 1]]$, pairs = $[[1, 3], [2, 0]]$,
Output: 4

Constraints:

- $2 \leq n \leq 500$
- n is even.
- $\text{preferences.length} == n$
- $\text{preferences}[i].\text{length} == n - 1$
- $0 \leq \text{preferences}[i][j] \leq n - 1$
- $\text{preferences}[i]$ does not contain i .
- All values in $\text{preferences}[i]$ are unique.
- $\text{pairs.length} == n/2$
- $\text{pairs}[i].\text{length} == 2$
- $x_i \neq y_i$
- $0 \leq x_i, y_i \leq n - 1$
- Each person is contained in **exactly one** pair.

JavaScript



```
1 /**
2  * @param {number} n
3  * @param {number[][]} preferences
4  * @param {number[][]} pairs
5  * @return {number}
6  */
7 var unhappyFriends = function(n, preferences, pairs) {
8
9 };
```