5815. Maximum Number of Points with Cost

My Submissions (/contest/weekly-contest-250/problems/maximum-number-of-points-with-cost/submissions/)

Back to Contest (/contest/weekly-contest-250/)

User Accepted:	0
User Tried:	0
Total Accepted:	0
Total Submissions:	0
Difficulty:	Medium

You are given an `m x n` integer matrix `points` (**0-indexed**). Starting with `0` points, you want to **maximize** the number of points you can get from the matrix.

To gain points, you must pick one cell in **each row**. Picking the cell at coordinates `(r, c)` will **add** `points[r][c]` to your score.

However, you will lose points if you pick a cell too far from the cell that you picked in the previous row. For every two adjacent rows `r` and `r + 1` (where `0 <= r < m - 1`), picking cells at coordinates `(r, c1)` and `(r + 1, c2)` will **subtract** `abs(c1 - c2)` from your score.

Return *the **maximum** number of points you can achieve*.

`abs(x)` is defined as:

- `x` for `x >= 0`.
- `-x` for `x < 0`.

Example 1:




```
Input: points = [[1,2,3],[1,5,1],[3,1,1]]
Output: 9
Explanation:
The blue cells denote the optimal cells to pick, which have coordinates (0, 2), (1, 1), and (2, 
You add 3 + 5 + 3 = 11 to your score.
However, you must subtract abs(2 - 1) + abs(1 - 0) = 2 from your score.
Your final score is 11 - 2 = 9.
```

Example 2: