

6317. Count the Number of Beautiful Subarrays

[My Submissions \(/contest/weekly-contest-336/problems/count-the-number-of-beautiful-subarrays/submissions/\)](/contest/weekly-contest-336/problems/count-the-number-of-beautiful-subarrays/submissions/)

[Back to Contest \(/contest/weekly-contest-336/\)](/contest/weekly-contest-336/)

You are given a **0-indexed** integer array `nums` . In one operation, you can:

- Choose two different indices i and j such that $0 \leq i, j < \text{nums.length}$.
- Choose a non-negative integer k such that the k^{th} bit (**0-indexed**) in the binary representation of $\text{nums}[i]$ and $\text{nums}[j]$ is 1.
- Subtract 2^k from $\text{nums}[i]$ and $\text{nums}[j]$.

A subarray is **beautiful** if it is possible to make all of its elements equal to 0 after applying the above operation any number of times.

Return the number of **beautiful subarrays** in the array `nums`.

A subarray is a contiguous **non-empty** sequence of elements within an array.

User Accepted:	0
User Tried:	0
Total Accepted:	0
Total Submissions:	0
Difficulty:	Medium

Example 1:

Input: nums = [4,3,1,2,4]

Output: 2

Explanation: There are 2 beautiful subarrays in nums: [4,3,1,2,4] and [4,3,1,2,4].

- We can make all elements in the subarray [3,1,2] equal to 0 in the following way:
 - Choose [3, 1, 2] and $k = 1$. Subtract 2^1 from both numbers. The subarray becomes [1, 1, 0].
 - Choose [1, 1, 0] and $k = 0$. Subtract 2^0 from both numbers. The subarray becomes [0, 0, 0].
- We can make all elements in the subarray [4,3,1,2,4] equal to 0 in the following way:
 - Choose [4, 3, 1, 2, 4] and $k = 2$. Subtract 2^2 from both numbers. The subarray becomes [0, 3, 1, 2, 0].
 - Choose [0, 3, 1, 2, 0] and $k = 0$. Subtract 2^0 from both numbers. The subarray becomes [0, 2, 0, 2, 0].
 - Choose [0, 2, 0, 2, 0] and $k = 1$. Subtract 2^1 from both numbers. The subarray becomes [0, 0, 0, 0, 0].

Example 2:

Input: nums = [1,10,4]

Output: 0

Explanation: There are no beautiful subarrays in nums.

Constraints:

- $1 \leq \text{nums.length} \leq 10^5$
- $0 \leq \text{nums}[i] \leq 10^6$

JavaScript ▾



```
1 const beautifulSubarrays = (a) => {
2   let res = 0, n = a.length, m = new Map([[0, 1]]), xor = 0;
3   for (let i = 0; i < n; i++) {
4     xor ^= a[i];
5     res += m.get(xor) || 0;
6     m.set(xor, m.get(xor) + 1 || 1);
7   }
8   return res;
9 };
```

☐ Custom Testcase

Use Example Testcases

Run

Submit

Submission Result: **Accepted** (/submissions/detail/913592174/) ?

More Details > (/submissions/detail/913592174/)

Share your acceptance!