

2470. Number of Subarrays With LCM Equal to K

[My Submissions \(/contest/weekly-contest-319/problems/number-of-subarrays-with-lcm-equal-to-k/submissions/\)](#)

[Back to Contest \(/contest/weekly-contest-319/\)](#)

Given an integer array `nums` and an integer `k`, return the number of **subarrays** of `nums` where the least common multiple of the subarray's elements is `k`.

A **subarray** is a contiguous non-empty sequence of elements within an array.

The **least common multiple of an array** is the smallest positive integer that is divisible by all the array elements.

Example 1:

**Input:** `nums = [3,6,2,7,1]`, `k = 6`  
**Output:** 4  
**Explanation:** The subarrays of `nums` where 6 is the least common multiple of all the subarray's elements are:  
- `[3,6,2,7,1]`  
- `[3,6,2,7,1]`  
- `[3,6,2,7,1]`  
- `[3,6,2,7,1]`

Example 2:

**Input:** `nums = [3]`, `k = 2`  
**Output:** 0  
**Explanation:** There are no subarrays of `nums` where 2 is the least common multiple of all the subarray's elements.

Constraints:

- 1 <= `nums.length` <= 1000
- 1 <= `nums[i]`, `k` <= 1000

[Discuss \(https://leetcode.com/problems/number-of-subarrays-with-lcm-equal-to-k/discuss\)](#)


JavaScript

📄 ↺ ⚙️

```
1 const gcd = (a, b) => b == 0 ? a : gcd(b, a % b);
2 const lcm = (a, b) => (a / gcd(a, b)) * b;
3
4 const subarrayLCM = (a, K) => {
5   let n = a.length, res = 0;
6   for (let i = 0; i < n; i++) {
7     let l = 1;
8     for (let j = i; j < n; j++) {
9       l = lcm(l, a[j]);
10      if (l > K) break;
11      if (l == K) res++;
12    }
13  }
14  return res;
15 };
```

☐ Custom Testcase [Use Example Testcases](#)

[Run](#) [Submit](#)

Submission Result: **Accepted** (/submissions/detail/842478370/)  [More Details > \(/submissions/detail/842478370/\)](#)

Share your acceptance!