

5468. Kth Missing Positive Number

[My Submissions \(/contest/biweekly-contest-32/problems/kth-missing-positive-number/submissions/\)](/contest/biweekly-contest-32/problems/kth-missing-positive-number/submissions/)

[Back to Contest \(/contest/biweekly-contest-32/\)](/contest/biweekly-contest-32/)

Given an array `arr` of positive integers sorted in a **strictly increasing order**, and an integer `k`.

Find the k^{th} positive integer that is missing from this array.

Example 1:

Input: `arr = [2,3,4,7,11]`, `k = 5`

Output: 9

Explanation: The missing positive integers are [1,5,6,8,

User Accepted: 4332

User Tried: 4788

Total Accepted: 4435

Total Submissions: 8607

Difficulty: Easy

Example 2:

Input: `arr = [1,2,3,4]`, `k = 2`

Output: 6

Explanation: The missing positive integers are [5,6,7,...]. The 2nd missing positive integer is 6.

Constraints:

- $1 \leq \text{arr.length} \leq 1000$
- $1 \leq \text{arr}[i] \leq 1000$
- $1 \leq k \leq 1000$
- $\text{arr}[i] < \text{arr}[j]$ for $1 \leq i < j \leq \text{arr.length}$

JavaScript



```
1 /**
2  * @param {number[]} arr
3  * @param {number} k
4  * @return {number}
5  */
6 var findKthPositive = function(arr, k) {
7
8 };
```

☐ **Custom Testcase**[Use Example Testcases](#)[▶ Run](#)[📁 Submit](#)

Copyright © 2020 LeetCode

[Help Center \(/support/\)](#) | [Terms \(/terms/\)](#) | [Privacy Policy \(/privacy/\)](#)

 [United States \(/region/\)](#)