



中山大學
SUN YAT-SEN UNIVERSITY

Lecture 3. Images Transformations

Pattern Recognition and Computer Vision

Guanbin Li,

School of Computer Science and Engineering, Sun Yat-Sen University

扫码签到



What will we learn today?

- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - Scaling
 - Rotation

Transformation Matrices

- Matrices can be used to transform vectors in useful ways, through multiplication: $p' = A p$
- Simplest is scaling:


$$\begin{array}{c} \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \\ A \end{array} \times \begin{array}{c} \begin{bmatrix} x \\ y \end{bmatrix} \\ p \end{array} = \begin{array}{c} \begin{bmatrix} s_x x \\ s_y y \end{bmatrix} \\ p' \end{array}$$

(Verify to yourself that the matrix multiplication works out this way)


Transformation Matrices

$$\begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} s_x x \\ s_y y \end{bmatrix}$$

$A \qquad p \qquad p'$




Rotation




Rotation

- How can you convert a vector represented in the coordinate frame “0” to a new, rotated coordinate frame “1”?
- Remember what a vector is:
[component in direction of the frame’s x axis, component in direction of y axis]



Rotation

- So to rotate it we must produce this vector:
[component in direction of **new** x axis, component in direction of **new** y axis]
- We can do this easily with dot products!
- New x coordinate is [the new x axis] **dot** [original vector]
- New y coordinate is [the new y axis] **dot** [original vector]




Rotation

- Insight: something similar happens in a matrix*vector multiplication!
- The resulting x coordinate, x' , is: [matrix row 1] dot [original vector]


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 0.707 & 0.707 \\ -0.707 & 0.707 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- The matrix multiplication $Rp = p'$ produces the coordinates in the new frame.




Rotation

- Now we have our point in the new coordinate system which is rotated left
- If we plot the result in the **original** coordinate system, we have rotated the point right



Rotation


- Now we have our point in the new coordinate system which is rotated left
- If we plot the result in the **original** coordinate system, we have rotated the point right



– Thus, rotation matrices can be used to rotate vectors. We'll usually think of them in that sense-- as operators to rotate vectors

2D Rotation Matrix Formula

Counter-clockwise rotation by an angle θ



$$x' = \cos \theta \, x - \sin \theta \, y$$

$$y' = \cos \theta \, y + \sin \theta \, x$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$p' = R p$$

Transformation Matrices

- Multiple transformation matrices can be used to transform a point:

$$p' = R_2 R_1 S p$$

- The effect of this is to apply their transformations one after the other, from **right to left**.
- In the example above, the result is equivalent to

$$p' = R_2(R_1(Sp))$$

- The result is exactly the same if we multiply the matrices first, to form a single transformation matrix:

$$p' = (R_2 R_1 S) p$$

What will we learn today?

- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - Scaling
 - Rotation

Homogeneous coordinates

- In general, a matrix multiplication lets us linearly combine components of a vector

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} ax + by \\ cx + dy \end{bmatrix}$$

- This is sufficient for scaling, rotating, and skewing transformations.
- But notice, we can't add a constant! ☹️
- That means, we cannot produce a new (translated) vector $\begin{bmatrix} x + k \\ y + k \end{bmatrix}$.

Homogeneous coordinates

- The (somewhat hacky) solution? Stick a “1” at the end of every vector:

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by + c \\ dx + ey + f \\ 1 \end{bmatrix}$$

- Now we can rotate, scale, and skew like before, **AND translate** (note how the multiplication works out, above)
- This is called “homogeneous coordinates”

Homogeneous coordinates

- In homogeneous coordinates, the multiplication works out so the rightmost column of the matrix is a vector that gets added.

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by + c \\ dx + ey + f \\ 1 \end{bmatrix}$$

- Generally, a homogeneous transformation matrix will have a bottom row of $[0 \ 0 \ 1]$, so that the result has a “1” at the bottom too.

Homogeneous coordinates


- One more thing we might want: to divide the result by something
 - For example, we may want to divide by a coordinate, to make things scale down as they get farther away in an image
 - Matrix multiplication can't actually divide
 - So, **by convention**, in homogeneous coordinates, we'll divide the result by its last coordinate after doing a matrix multiplication

$$\begin{bmatrix} x \\ y \\ 7 \end{bmatrix} \Rightarrow \begin{bmatrix} x/7 \\ y/7 \\ 1 \end{bmatrix}$$


What will we learn today?

- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - Scaling
 - Rotation

2D Translation



2D Translation using Homogeneous Coordinates



$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$


$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} + \begin{bmatrix} t_x \\ t_y \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates



$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$


$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates



$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$


$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates




$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates




$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates




$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$

$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D Translation using Homogeneous Coordinates



$$p = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$t = \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} t_x \\ t_y \\ 1 \end{bmatrix}$$


$$p' = Tp$$

$$p' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} I & t \\ 0 & 1 \end{bmatrix} p = Tp$$


What will we learn today?

- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - **Scaling**
 - Rotation

Scaling



Scaling Equation




$$p = \begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$p' = \begin{bmatrix} s_x x \\ s_y y \end{bmatrix} \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix}$$

$$p' = Sp$$

$$p' \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Scaling Equation




$$p = \begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$p' = \begin{bmatrix} s_x x \\ s_y y \end{bmatrix} \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix}$$

$$p' = Sp$$

$$p' \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s' & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} p = Sp$$

Scaling & Translating



Scaling & Translating

$$p'' = TSp = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & t_x \\ 0 & s_y & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} S' & \mathbf{t} \\ \mathbf{0} & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + t_x \\ s_y y + t_y \\ 1 \end{bmatrix}$$

Scaling & Translating \neq Translating & Scaling


$$p'' = TSp = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & t_x \\ 0 & s_y & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + t_x \\ s_y y + t_y \\ 1 \end{bmatrix}$$

$$p''' = STp = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & s_x t_x \\ 0 & s_y & s_y t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + s_x t_x \\ s_y y + s_y t_y \\ 1 \end{bmatrix}$$

What will we learn today?


- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - Scaling
 - Rotation

Rotation



2D Rotation Matrix Formula

Counter-clockwise rotation by an angle θ



$$x' = \cos \theta x - \sin \theta y$$

$$y' = \cos \theta y + \sin \theta x$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$p' = Rp$$

Rotation Matrix Properties

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

A 2D rotation matrix is 2x2

Note: \mathbf{R} belongs to the category of *normal* matrices and satisfies many interesting properties:

$$\mathbf{R} \cdot \mathbf{R}^T = \mathbf{R}^T \cdot \mathbf{R} = \mathbf{I}$$

$$\det(\mathbf{R}) = 1$$

Rotation Matrix Properties

- Transpose of a rotation matrix produces a rotation in the opposite direction

$$\mathbf{R} \cdot \mathbf{R}^T = \mathbf{R}^T \cdot \mathbf{R} = \mathbf{I}$$

$$\det(\mathbf{R}) = 1$$

- The rows of a rotation matrix are always mutually perpendicular (a.k.a. orthogonal) unit vectors
– (and so are its columns)

Scaling + Rotation + Translation

$$p' = (T R S) p$$

$$p' = TRSp = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} \cos \theta & -\sin \theta & t_x \\ \sin \theta & \cos \theta & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} R & t \\ 0 & 1 \end{bmatrix} \begin{bmatrix} S & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \boxed{\begin{bmatrix} RS & t \\ 0 & 1 \end{bmatrix}} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

This is the form of the general-purpose transformation matrix

Summary

- 2D transformations
 - Transformation Matrices
 - Homogeneous coordinates
 - Translation
 - Scaling
 - Rotation