

MATH2621 — Higher Complex Analysis. IX

Harmonic functions

This lecture?

In this lecture, we introduce harmonic functions, and we discuss their significance.

This lecture?

In this lecture, we introduce harmonic functions, and we discuss their significance.

We see how to find them using holomorphic functions, and conversely, how to find holomorphic functions using harmonic functions.

Harmonic functions

Definition

Suppose that $u : \Omega \rightarrow \mathbb{R}$ is a function, where Ω is an open subset of \mathbb{R}^2 , and that u is twice continuously differentiable, that is, all the partial derivatives $\partial u / \partial x$, $\partial u / \partial y$, $\partial^2 u / \partial x^2$, $\partial^2 u / \partial x \partial y$, $\partial^2 u / \partial y \partial x$ and $\partial^2 u / \partial y^2$ exist and are continuous. Then we say that u is *harmonic* in Ω if

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

Finding harmonic functions

Theorem

Suppose that $f \in H(\Omega)$, where Ω is an open subset of \mathbb{C} , that f is twice continuously differentiable in Ω , and that

$$f(x + iy) = u(x, y) + iv(x, y)$$

in Ω , where u and v are real-valued. Then u and v are harmonic functions.

Finding harmonic functions

Theorem

Suppose that $f \in H(\Omega)$, where Ω is an open subset of \mathbb{C} , that f is twice continuously differentiable in Ω , and that

$$f(x + iy) = u(x, y) + iv(x, y)$$

in Ω , where u and v are real-valued. Then u and v are harmonic functions.

Later we will see that the hypothesis that f is twice differentiable is not needed.

Proof of theorem

Proof. Suppose that $f \in H(\Omega)$, where Ω is an open subset of \mathbb{C} , and that

$$f(x + iy) = u(x, y) + iv(x, y)$$

in Ω , where u and v are real-valued, and that all the partial derivatives $\partial u/\partial x$, $\partial u/\partial y$, $\partial^2 u/\partial x^2$, $\partial^2 u/\partial x \partial y$, $\partial^2 u/\partial y \partial x$ and $\partial^2 u/\partial y^2$ and $\partial v/\partial x$, $\partial v/\partial y$, $\partial^2 v/\partial x^2$, $\partial^2 v/\partial x \partial y$, $\partial^2 v/\partial y \partial x$ and $\partial^2 v/\partial y^2$ exist and are continuous.

Proof of theorem

Proof. Suppose that $f \in H(\Omega)$, where Ω is an open subset of \mathbb{C} , and that

$$f(x + iy) = u(x, y) + iv(x, y)$$

in Ω , where u and v are real-valued, and that all the partial derivatives $\partial u/\partial x$, $\partial u/\partial y$, $\partial^2 u/\partial x^2$, $\partial^2 u/\partial x \partial y$, $\partial^2 u/\partial y \partial x$ and $\partial^2 u/\partial y^2$ and $\partial v/\partial x$, $\partial v/\partial y$, $\partial^2 v/\partial x^2$, $\partial^2 v/\partial x \partial y$, $\partial^2 v/\partial y \partial x$ and $\partial^2 v/\partial y^2$ exist and are continuous.

From vector calculus, this implies that

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x} \quad \text{and} \quad \frac{\partial^2 v}{\partial x \partial y} = \frac{\partial^2 v}{\partial y \partial x}.$$

Proof of theorem (continued)

Now, from the Cauchy–Riemann equations,

$$\begin{aligned}\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} &= \frac{\partial}{\partial x} \frac{\partial u}{\partial x} + \frac{\partial}{\partial y} \frac{\partial u}{\partial y} \\ &= \frac{\partial}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial}{\partial y} \frac{\partial v}{\partial x} \\ &= \frac{\partial^2 v}{\partial x \partial y} - \frac{\partial^2 v}{\partial y \partial x} = 0,\end{aligned}$$

so u is harmonic. Similarly, v is harmonic. □

Exercise 1

Suppose that $u(x, y) = x^3 - 3xy^2$. Show that u is harmonic in \mathbb{C} , and find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer.

Exercise 1

Suppose that $u(x, y) = x^3 - 3xy^2$. Show that u is harmonic in \mathbb{C} , and find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer. The partial derivatives of u are given by:

$$\frac{\partial u}{\partial x} = 3x^2 - 3y^2$$

$$\frac{\partial^2 u}{\partial x^2} = 6x$$

$$\frac{\partial u}{\partial y} = -6xy$$

$$\frac{\partial^2 u}{\partial y^2} = -6x.$$

Exercise 1

Suppose that $u(x, y) = x^3 - 3xy^2$. Show that u is harmonic in \mathbb{C} , and find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer. The partial derivatives of u are given by:

$$\frac{\partial u}{\partial x} = 3x^2 - 3y^2$$

$$\frac{\partial^2 u}{\partial x^2} = 6x$$

$$\frac{\partial u}{\partial y} = -6xy$$

$$\frac{\partial^2 u}{\partial y^2} = -6x.$$

Hence

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 6x - 6x = 0,$$

and u is harmonic.

Answer to Exercise 1

If v exists such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic, then the Cauchy–Riemann equations hold, so

$$\frac{\partial v}{\partial y} = 3x^2 - 3y^2 \quad (1)$$

$$\frac{\partial v}{\partial x} = 6xy. \quad (2)$$

Answer to Exercise 1

If v exists such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic, then the Cauchy–Riemann equations hold, so

$$\frac{\partial v}{\partial y} = 3x^2 - 3y^2 \quad (1)$$

$$\frac{\partial v}{\partial x} = 6xy. \quad (2)$$

Fix $y \in \mathbb{R}$. Integrating (2) with respect to x shows that

$$v(x, y) = 3x^2y + C_1,$$

where C_1 does not depend on x . However, C_1 may depend on y . Thus the general solution to (2), when y can vary, is

$$v(x, y) = 3x^2y + c(y),$$

where c is an unknown function.

Answer to Exercise 1 (continued)

The function v must also satisfy (1), and this happens as long as

$$3x^2 - 3y^2 = \frac{\partial v}{\partial y} = 3x^2 + c'(y), \quad (3)$$

i.e., $c'(y) = -3y^2$, so $c(y) = -y^3 + C$, where C is a constant, and

$$v(x, y) = 3x^2y - y^3 + C.$$

Answer to Exercise 1 (continued)

The function v must also satisfy (1), and this happens as long as

$$3x^2 - 3y^2 = \frac{\partial v}{\partial y} = 3x^2 + c'(y), \quad (3)$$

i.e., $c'(y) = -3y^2$, so $c(y) = -y^3 + C$, where C is a constant, and

$$v(x, y) = 3x^2y - y^3 + C.$$

Set $f(x + iy) = u(x, y) + iv(x, y)$. Then f is holomorphic, and

$$f(z) = f(x + iy) = x^3 - 3xy^2 + i(3x^2y - y^3) + iC = z^3 + iC. \quad \triangle$$

Answer to Exercise 1 (continued)

The function v must also satisfy (1), and this happens as long as

$$3x^2 - 3y^2 = \frac{\partial v}{\partial y} = 3x^2 + c'(y), \quad (3)$$

i.e., $c'(y) = -3y^2$, so $c(y) = -y^3 + C$, where C is a constant, and

$$v(x, y) = 3x^2y - y^3 + C.$$

Set $f(x + iy) = u(x, y) + iv(x, y)$. Then f is holomorphic, and

$$f(z) = f(x + iy) = x^3 - 3xy^2 + i(3x^2y - y^3) + iC = z^3 + iC. \quad \triangle$$

In the solution above, if we just used the constant C_1 rather than the function $c(y)$, then we wouldn't be able to satisfy (3).

Exercise 2

Suppose that $u(x, y) = x^2 - y^2$. Show that u is harmonic in \mathbb{C} . Find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer.

Exercise 2

Suppose that $u(x, y) = x^2 - y^2$. Show that u is harmonic in \mathbb{C} . Find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer. The partial derivatives of u are given by:

$$\frac{\partial u}{\partial x} = 2x$$

$$\frac{\partial^2 u}{\partial x^2} = 2$$

$$\frac{\partial u}{\partial y} = -2y$$

$$\frac{\partial^2 u}{\partial y^2} = -2.$$

Exercise 2

Suppose that $u(x, y) = x^2 - y^2$. Show that u is harmonic in \mathbb{C} . Find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer. The partial derivatives of u are given by:

$$\frac{\partial u}{\partial x} = 2x$$

$$\frac{\partial^2 u}{\partial x^2} = 2$$

$$\frac{\partial u}{\partial y} = -2y$$

$$\frac{\partial^2 u}{\partial y^2} = -2.$$

Hence

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 2 - 2 = 0,$$

and u is harmonic.

Answer to Exercise 2

If v exists such that the function f given by

$$f(x + iy) = u(x, y) + iv(x, y)$$

is holomorphic, then the Cauchy–Riemann equations must hold, so we must have

$$\frac{\partial v}{\partial y} = \frac{\partial u}{\partial x} = 2x \tag{1}$$

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = 2y. \tag{2}$$

Answer to Exercise 2 (continued)

The general solution to equation (1) is

$$v(x, y) = 2xy + c(x),$$

where c is an unknown function.

Answer to Exercise 2 (continued)

The general solution to equation (1) is

$$v(x, y) = 2xy + c(x),$$

where c is an unknown function.

For (2) to hold as well, we require that

$$2y = \frac{\partial v}{\partial x} = 2y + c'(x)$$

for all x . This tells us that $c'(x) = 0$ for all x and c has to be a constant, C say.

Answer to Exercise 2 (continued)

The general solution to equation (1) is

$$v(x, y) = 2xy + c(x),$$

where c is an unknown function.

For (2) to hold as well, we require that

$$2y = \frac{\partial v}{\partial x} = 2y + c'(x)$$

for all x . This tells us that $c'(x) = 0$ for all x and c has to be a constant, C say.

Set $f(x + iy) = u(x, y) + iv(x, y)$. Then f is holomorphic and

$$f(z) = f(x + iy) = x^2 - y^2 + 2ixy + iC = z^2 + iC.$$



Existence of harmonic functions

Recall that a domain is a polygonally path-connected open set.

Existence of harmonic functions

Recall that a domain is a polygonally path-connected open set.

Theorem

If Ω is a simply polygonally connected domain, and $u : \Omega \rightarrow \mathbb{R}$ is harmonic, then there exists a harmonic function $v : \Omega \rightarrow \mathbb{R}$ such that f , given by

$$f(x + iy) = u(x, y) + iv(x, y)$$

in Ω , is holomorphic. Any two such functions v differ by an additive constant.

Remarks

The function v is called a harmonic conjugate of u .

Remarks

The function v is called a harmonic conjugate of u .

The function f may often be determined using the fact that

$$f'(x + iy) = u_x(x, y) + iv_x(x, y) = u_x(x, y) - iu_y(x, y).$$

Remarks

The function v is called a harmonic conjugate of u .

The function f may often be determined using the fact that

$$f'(x + iy) = u_x(x, y) + iv_x(x, y) = u_x(x, y) - iu_y(x, y).$$

We give a partial proof of this theorem later.

Remarks

The function v is called a harmonic conjugate of u .

The function f may often be determined using the fact that

$$f'(x + iy) = u_x(x, y) + iv_x(x, y) = u_x(x, y) - iu_y(x, y).$$

We give a partial proof of this theorem later.

To see why it might be necessary to require that Ω be simply connected in the theorem, consider the following.

Exercise 3

Suppose that $u(x, y) = \ln(\sqrt{x^2 + y^2})$ in $\mathbb{R}^2 \setminus \{(0, 0)\}$. Show that u is harmonic. Find a function v such that the function f in $\mathbb{C} \setminus \{0\}$, defined by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic.

Answer.

Exercise 3

Suppose that $u(x, y) = \ln(\sqrt{x^2 + y^2})$ in $\mathbb{R}^2 \setminus \{(0, 0)\}$. Show that u is harmonic. Find a function v such that the function f in $\mathbb{C} \setminus \{0\}$, defined by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic.

Answer. You do this!



Check that having a harmonic conjugate is a local property. The problem here is that we cannot find a global harmonic conjugate to u .

Exercise 4

Suppose that $u(x, y) = x^2 - y^2$. Show that u is harmonic in \mathbb{C} . Find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer.

Exercise 4

Suppose that $u(x, y) = x^2 - y^2$. Show that u is harmonic in \mathbb{C} . Find a function v such that the function f , given by $f(x + iy) = u(x, y) + iv(x, y)$, is holomorphic in \mathbb{C} .

Answer. The partial derivatives of u are given by:

$$\frac{\partial u}{\partial x} = 2x \quad \text{and} \quad \frac{\partial u}{\partial y} = -2y.$$

Hence

$$\frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = 2(x + iy) = 2z.$$

Answer to Exercise 4

Clearly, if $f(z) = z^2$, then $f'(z) = 2z$, so we just observe that $\operatorname{Re} f(x + iy) = u(x, y)$, so u is the real part of a holomorphic function and hence is harmonic. Further, $f = u + iv$, where $v(x, y) = 2xy$. △

Answer to Exercise 4

Clearly, if $f(z) = z^2$, then $f'(z) = 2z$, so we just observe that $\operatorname{Re} f(x + iy) = u(x, y)$, so u is the real part of a holomorphic function and hence is harmonic. Further, $f = u + iv$, where $v(x, y) = 2xy$. △

The trouble with this method is that it may be hard to express $u_x - iu_y$ as a function of z , or hard to find the integral of this.