

Rudin, Principles of Mathematical Analysis

Chapter 1 Answer Key

1. If r is rational ($r \neq 0$) and x is irrational, prove that $r + x$ and rx are irrational.

Answer: Assume otherwise that $r + x$ is rational with sum α . We write $r + x = \alpha \implies x = \alpha - r$. This implies that x can be expressed as the difference of two rational numbers, $\alpha - r$, which is rational. Thus, $r + x$ must be irrational. $\rightarrow \leftarrow$

We follow a similar argument for the second part. Assume otherwise that rx is rational with product $\frac{\beta}{\gamma}$. We write $rx = \frac{\beta}{\gamma} \implies x = \frac{\beta}{r\gamma}$. This implies that x can be expressed as a (compound) fraction, which violates the irrational property. Thus, rx must be irrational. $\rightarrow \leftarrow$

2. Prove that there is no rational number whose square is 12.

Answer: Assume otherwise that there exists a rational $p = \frac{m}{n}$ for m, n not both even, such that:

$$p^2 = 12$$

We can re-arrange as follows:

$$\left(\frac{m}{n}\right)^2 = 12$$

$$m^2 = 12n^2$$

Since 12 is even, and the product of two terms involving an even number is always even, the RHS must be even. So, by the equality, m^2 is also even, which implies that m is even and n is odd. We can write $m = 2k$ for some integer k and substitute:

$$4k^2 = 12n^2$$

$$k^2 = 3n^2$$

We know that n is odd, hence $3n^2$ is odd, which implies k^2 and k are also odd. We can now express $k = 2j + 1$ and $n = 2l + 1$ for some integers j, l and substitute:

$$(2j + 1)^2 = 3(2l + 1)^2$$

$$4j^2 + 4j + 1 = 3(4l^2 + 4l + 1)$$

$$4j^2 + 4j + 1 = 12l^2 + 12l + 3$$

$$4j^2 + 4j - 12l^2 - 12l = 2$$

$$4(j^2 + j - l^2 - l) = 2$$

This equality cannot hold because 4 is not a multiple of 2. Thus, p must be irrational.
 $\rightarrow\leftarrow$

3. Prove Proposition 1.15.
4. Let E be a nonempty subset of an ordered set; suppose x is a lower bound of E and β is an upper bound of E . Prove that $x \leq \beta$.
5. Let A be a nonempty set of real numbers which is bounded below. Let $-A$ be the set of all numbers $-x$, where $x \in A$. Prove that

$$\inf A = -\sup(-A).$$