



diamond: 10922 orthologs on 219 Pmax scaffolds and 25 Ssol scaffolds.



C Pmax vs Pmax scaffolds

	Pmax scaf	Ssol scaf	p (FET)	Count
0	NC_047015.1	Chr01	0.00e+00	831
1	NC_047017.1	Chr03	0.00e+00	818
2	NC_047018.1	Chr06	0.00e+00	730
3	NC_047019.1	Chr04	0.00e+00	622
4	NC_047025.1	Chr05	0.00e+00	586
5	NC_047021.1	Chr08	0.00e+00	568
6	NC_047020.1	Chr02	0.00e+00	530
7	NC_047022.1	Chr07	0.00e+00	529
8	NC_047024.1	Chr14	0.00e+00	509
9	NC_047023.1	Chr13	0.00e+00	484
10	NC_047026.1	Chr10	0.00e+00	477
11	NC_047016.1	Chr11	0.00e+00	390
12	NC_047028.1	Chr15	0.00e+00	363
13	NC_047029.1	Chr16	0.00e+00	360
14	NC_047030.1	Chr12	0.00e+00	357
15	NC_047032.1	Chr17	0.00e+00	289
16	NC_047016.1	Chr09	6.45e-167	270
17	NC_047033.1	Chr19	1.56e-305	203
18	NC_047027.1	Chr09	1.61e-202	197
19	NC_047031.1	Chr18	3.65e-202	140

e Pmax scaffolds

	Pmax scaf	Ssol scaf	p (FET)	Count
0	NC_047015.1 : 0-end	Chr01 : 0-end	0.00e+00	831
1	NC_047017.1 : 0-end	Chr03 : 0-end	0.00e+00	818
2	NC_047018.1 : 0-end	Chr06 : 0-end	0.00e+00	730
3	NC_047019.1 : 0-end	Chr04 : 0-end	0.00e+00	622
4	NC_047025.1 : 0-end	Chr05 : 0-end	0.00e+00	586
5	NC_047021.1 : 0-end	Chr08 : 0-end	0.00e+00	568
6	NC_047020.1 : 0-end	Chr02 : 0-end	0.00e+00	530
7	NC_047022.1 : 0-end	Chr07 : 0-end	0.00e+00	529
8	NC_047024.1 : 0-end	Chr14 : 0-end	0.00e+00	509
9	NC_047023.1 : 0-end	Chr13 : 0-end	0.00e+00	484
10	NC_047026.1 : 0-end	Chr10 : 0-end	0.00e+00	477
11	NC_047016.1 : 0-end	Chr11 : 0-end	0.00e+00	390
12	NC_047028.1 : 0-end	Chr15 : 0-end	0.00e+00	363
13	NC_047029.1 : 0-end	Chr16 : 0-end	0.00e+00	360
14	NC_047030.1 : 0-end	Chr12 : 0-end	0.00e+00	357
15	NC_047032.1 : 0-end	Chr17 : 0-end	0.00e+00	289
16	NC_047016.1 : 0-end	Chr09 : 0-end	6.45e-167	270
17	NC_047033.1 : 0-end	Chr19 : 0-end	1.56e-305	203
18	NC_047027.1 : 0-end	Chr09 : 0-end	1.61e-202	197
19	NC_047031.1 : 0-end	Chr18 : 0-end	3.65e-202	140

(a) depicts the chromosomes/scaffolds of Pmax (x-axis) plotted against the chromosomes/scaffolds of Ssol (y_axis). Each dot in the plot represents an ortholog, specifically a reciprocal best diamond blastp match between two species. The unit of the x- and y-axes are the number of orthologous proteins between these two species: 10922 orthologs found between 219 Pmax scaffolds and 25 Ssol scaffolds. If there are chromosome breaks, Fisher's exact test is used to calculate the significance of the interactions between the sub-chromosomal pieces. Otherwise Fisher's exact test is calculated on whole chromosomes. The opacity of the dots depict the significance from Fisher's exact test. Dots that are a solid color are in cells with a FET p-value less than or equal to 0.05. Dots that are translucent are in cells with a FET p-value greater than 0.05. The Dx and Dy values depict places where there may be sudden breaks in synteny. See the supplementary information the following paper for more information on Dx and Dy: Simakov, Oleg, et al. Deeply conserved synteny resolves early events in vertebrate evolution. *Nature Ecology & Evolution* 4.6 (2020): 820-830. (b) depicts the same information as panel a, but it is plotted in the organisms chromosome basepair coordinates rather than gene index. This is useful for visualizing gene-poor regions of the chromosomes (c) shows which gene groups are most prevalent in each chromosome pair. The FET p-value in this table corresponds to the whole-chromosome FET p-value, and is not a FET value of the correlation between the chromosome pair and the gene group. (d) shows chromosome-scale significance values. This table shows the same information as c, but does not factor in gene group information. (e) shows the FET p-values of the sub-chromosomal compartments. Useful for showing if single arms of chromosomes are correlated with other regions.