



Logistic Regression Model

Data: $\{(X_{1,1}, X_{1,2}, \cdots, X_{1,p}), Y_1\}, \{(X_{2,1}, X_{2,2}, \cdots, X_{2,p}), Y_2\}, \cdots, \{(X_{n,1}, X_{n,2}, \cdots, X_{n,p}), Y_n\}$ where Y_1, \cdots, Y_n are binary responses

Model: We model the probability of success given the predictor(s)

$$p = p(X_1, \dots, X_p) = Pr(Y = 1 \mid X_1, \dots, X_p)$$

by linking p to the predicting variables through the *logit* link function g:

$$g(p) = \ln\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 X_1 + \dots + \beta_p X_p$$

OR

$$p(X_1, \dots, X_p) = \frac{e^{\beta_0 + \beta_1 X_1 + \dots + \beta_p X_p}}{1 + e^{\beta_0 + \beta_1 X_1 + \dots + \beta_p X_p}}$$

Georgia Tech

Model Interpretation

- The probability of success given one predicting variable X = x is p = p(x) = Pr(Y = 1 | x)
- The logit function $\ln\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 x$ is the **log odds** function.
- The exponential of the logit function $\frac{p(X)}{1-p(X)} = e^{\beta_0 + \beta_1 x}$ is the **odds** of Y = 1 at X = x
- The odds at X = a versus X = b is equal to the **odds ratio**:

$$\frac{\mathrm{e}^{\beta_0 + \beta_1 a}}{\mathrm{e}^{\beta_0 + \beta_1 b}} = \mathrm{e}^{\beta_1 (a - b)}$$

Georgia Tech

Model Interpretation

If we calculate the odds ratio of the odds at X = b + 1 versus X = b, we have

$$\frac{\mathrm{e}^{\beta_0+\beta_1(b+1)}}{\mathrm{e}^{\beta_0+\beta_1b}}=\mathrm{e}^{\beta_1}$$

- The regression coefficient β_1 can be interpreted as the log of the odds ratio for an increase of one unit in the predicting variable.
- If X a dummy variable of a categorical factor, interpret as the log of odds ratio of one category versus baseline.
- Interpret β with respect to the odds of success, not directly with respect to the response variable.

Georgia Tech

Model Estimation

Model the probability of success given predictor(s):

$$\operatorname{Logit}\!\left(\operatorname{Pr}\!\left(Y=1\mid X_1,\cdots,X_p\right)\right)=\operatorname{Logit}\!\left(\operatorname{p}\!\left(X_1,\cdots,X_p\right)\right)=\beta_0+\beta_1X_1+\cdots+\beta_pX_p$$

Parameters: $\beta_0, \beta_1, \cdots, \beta_n$

Approach: Maximum Likelihood Estimation

$$\max_{\beta_0,\beta_1,\cdots,\beta_p} \mathcal{L}(\beta_0,\beta_1,\cdots,\beta_p) = \prod_{i=1}^n p(X_{i,1},X_{i,2},\cdots,X_{i,p})^{Y_i} \left(1 - p(X_{i,1},X_{i,2},\cdots,X_{i,p})\right)^{1-Y_i}$$

or

$$\begin{aligned} &\max_{\beta_0,\beta_1,\cdots,\beta_p} \ell(\beta_0,\beta_1,\cdots,\beta_p) = \max_{\beta_0,\beta_1,\cdots,\beta_p} \log(\mathcal{L}(\beta_0,\beta_1,\cdots,\beta_p)) \\ &= \max_{\beta_0,\beta_1,\cdots,\beta_p} \sum_{i=1}^n \left(Y_i \log\left(\frac{\mathrm{e}^{\beta_0+\beta_1 X_1+\cdots+\beta_p X_p}}{1+\mathrm{e}^{\beta_0+\beta_1 X_1+\cdots+\beta_p X_p}}\right) + (1-Y_i) \log\left(\frac{1}{1+\mathrm{e}^{\beta_0+\beta_1 X_1+\cdots+\beta_p X_p}}\right) \right) \end{aligned}$$

Georgia Tech

Model Estimation (cont'd)

Approach: Maximum Likelihood Estimation

$$\max_{\beta_0,\beta_1,\cdots,\beta_p} \sum_{i=1}^{n} \left(Y_i \log \left(\frac{e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}}{1 + e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}} \right) + (1 - Y_i) \log \left(\frac{1}{1 + e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}} \right) \right)$$

- Maximizing the (log-)likelihood function with respect to $\beta_0, \beta_1, \cdots, \beta_p$ in closed form expression is not possible because the (log-)likelihood function is a non-linear function in the model parameters.
- Use numerical algorithm to estimate $\beta_0, \beta_1, \dots, \beta_p \Rightarrow \hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_p$

Upshot: The estimated parameters and their standard errors are approximate estimates. Do not attempt to do it yourself! Use statistical software to derive the estimated regression coefficients.

Georgia Tech

Summary



Georgia Tech