

# The POWHEG BOX user manual: Higgs boson production through gluon fusion

---

## Simone Alioli

*Deutsches Elektronen-Synchrotron DESY  
Platanenallee 6, D-15738 Zeuthen, Germany  
E-mail: [simone.alioli@desy.de](mailto:simone.alioli@desy.de)*

## Paolo Nason

*INFN, Sezione di Milano-Bicocca, Piazza della Scienza 3, 20126 Milan, Italy  
E-mail: [Paolo.Nason@mib.infn.it](mailto:Paolo.Nason@mib.infn.it)*

## Carlo Oleari

*Università di Milano-Bicocca and INFN, Sezione di Milano-Bicocca  
Piazza della Scienza 3, 20126 Milan, Italy  
E-mail: [Carlo.Oleari@mib.infn.it](mailto:Carlo.Oleari@mib.infn.it)*

## Emanuele Re

*Institute for Particle Physics Phenomenology, Department of Physics  
University of Durham, Durham, DH1 3LE, UK  
E-mail: [emanuele.re@durham.ac.uk](mailto:emanuele.re@durham.ac.uk)*

**ABSTRACT:** This note documents the use of the package POWHEG BOX for Higgs boson production through gluon fusion. Results can be easily interfaced to shower Monte Carlo programs, in such a way that both NLO and shower accuracy are maintained.

**KEYWORDS:** POWHEG, Shower Monte Carlo, NLO.

---

## Contents

<b>1. Introduction</b>	<b>1</b>
<b>2. Generation of events</b>	<b>2</b>
<b>3. Process specific input parameters</b>	<b>2</b>

---

## 1. Introduction

The POWHEG BOX program is a framework for implementing NLO calculations in Shower Monte Carlo programs according to the POWHEG method. An explanation of the method and a discussion of how the code is organized can be found in refs. [1, 2, 3]. The code is distributed according to the “MCNET GUIDELINES for Event Generator Authors and Users” and can be found at the web page

<http://powhegbox.mib.infn.it>.

This program is an implementation of the NLO cross section for Higgs boson production via gluon fusion process, first evaluated in refs. [6, 7, 8], in the POWHEG formalism of refs. [1, 2]. A detailed description of the implementation can be found on ref. [9]. We allow either to retain the full top-mass dependence in the leading order contribution, either to perform the calculation of NLO terms in the large top-mass limit. Spin correlations of Higgs boson decay products are not included, being it a scalar. This issue can be safely left to the subsequent Shower Monte Carlo program. Finite Higgs boson width effects are accounted for. The code, that can be found in the POWHEG-BOX/gg\_H subdirectory, is based on the subtraction scheme by Frixione, Kunszt and Signer implemented in the POWHEG BOX, rather than on the scheme discussed in the paper [9]. Please cite it anyhow if you use the program.

In order to run the POWHEG BOX program, we recommend the reader to start from the POWHEG BOX user manual, which contains all the information and settings that are common between all subprocesses. In this note we focus on the settings and parameters specific to  $gg \rightarrow H$  implementation.

## 2. Generation of events

Build the executable

```
$ cd POWHEG-BOX/gg_H
```

```
$ make pwhg_main
```

Then do (for example)

```
$ cd testrun-lhc
```

```
$ ../pwhg_main
```

At the end of the run, the file `pwgevents.lhe` will contain 100000 events for  $gg \rightarrow H$  in the Les Houches format. In order to shower them with PYTHIA do

```
$ cd POWHEG-BOX/gg_H
```

```
$ make main-PYTHIA-lhef
```

```
$ cd testrun-lhc
```

```
$ ../main-PYTHIA-lhef
```

## 3. Process specific input parameters

The mandatory parameters are

```
gfermi 0.116639D-04 ! Fermi constant
hmass 120 ! mass of Higgs boson in GeV
hwidth 0.003605 ! width of Higgs boson in GeV
topmass 171.3 ! top quark mass in GeV
```

The running of  $\alpha_s$  is evaluated at two loop order, correctly matching, at flavour thresholds, different definitions that depends on the number of flavours that can be considered light at the renormalization scale. Examples of `powheg.input` files are given in the subdirectories `gg_H/testrun-tev` and `gg_H/testrun-lhc`. In all examples, the choice of the parameters that control the grid generation is such that a reasonably small fraction of negative weights is generated, so they can be run as they are. We remind the reader that these negative weights are only due to our choice of generating  $\tilde{B}$  instead of  $\bar{B}$ . They indeed correspond to phase space points where NLO corrections are bigger than LO contributions. Had we performed the integration over the full radiation phase space these negative weights would have disappeared completely.

In case one is interfacing to HERWIG or PYTHIA SMC programs, we provide a facility to select the Higgs boson decay products in these programs :

```
hdecaymode 12      ! code for selection of Higgs boson decay products:
                   ! -1 the Higgs boson is left undecayed by the SMC
                   ! 0 all decay channels are open
                   ! 1-6 d dbar, u ubar, ..., t tbar (as in HERWIG)
                   ! 7-9 e+ e-, mu+ mu-, tau+ tau-
                   ! 10, 11, 12 W+W-, ZZ, gamma gamma
```

Together with the mandatory parameters, the POWHEG BOX input facility allows for an easy setting of run parameters, by explicitly adding the relevant lines to the input card. In case one of the following entries is not present in the input card the reported default value is assumed. In any case, these parameters are printed in the output of the program, so their values can be easily tracked down.

```

masswindow_low 10 ! M_H > Hmass - masswindow_low * Hwidth
masswindow_high 10 ! M_H < Hmass + masswindow_high * Hwidth
runningscale 0 ! choice for ren and fac scales in Bbar integration
                    0: fixed scale M_H
                    1: running scale inv mass H
                    2: running scale transverse mass H

```

Of particular importance are the following parameters:

- **largentlim 0** (default 0) controls how the large top-mass approximation is enforced. The default behaviour is to evaluate all the NLO contributions in the large top mass limit and then reweight them by a top-mass correction factor given by the ratio of LO results ( $B_{(m_t)}/B_{(m_t \rightarrow \infty)}$ ). Setting this parameter to 1 the correction factor is instead omitted and the contributions are all evaluated in the large top mass limit. Setting it to 2 the correction factor is again omitted but the LO contributions are evaluated with the full top-mass dependence (old default)
- **hhfact 100d0 !** (default no dumping factor) dump factor for high-pt radiation:  $> 0$  **dumpfac** =  $h^2/(p_T^2 + h^2)$  controls how much of real contribution enters in the POWHEG Sudakov form factor. By default all real contributions are included, but this may lead to a NNLO mismatch in the higher Higgs boson  $p_T$  distribution tail, with respect to fixed order NLO results. This actually brings POWHEG BOX results closer to NNLO ones, but if one want to switch-off this feature it's possible to use a reduced real contribution  $R^{\text{red}} = R \times \text{dumpfact}$  in the Sudakov and to generate the remaining  $R \times (1 - \text{dumpfact})$  part without suppression, as documented in Sec. 4.3 of ref. [9].
- **zerowidth 1** (default 0 = false) enforce the calculation in the Higgs zero width approximation.
- **bwshape 2** (default 1) choose the functional form of the Breit-Wigner along which the Higgs virtuality is distributed, in case of the zero width approximation has not been chosen. Allowed values are 1 for a BW with a running width, or 2 for a fixed width.

## References

- [1] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms,” JHEP **0411** (2004) 040 [arXiv:hep-ph/0409146].

- [2] S. Frixione, P. Nason and C. Oleari, “Matching NLO QCD computations with Parton Shower simulations: the POWHEG method,” *JHEP* **0711** (2007) 070 [arXiv:0709.2092 [hep-ph]].
- [3] S. Alioli, P. Nason, C. Oleari and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX,” [arXiv:1002.2581 [hep-ph]].
- [4] S. Frixione and B. R. Webber, “Matching NLO QCD computations and parton shower simulations,” *JHEP* **0206** (2002) 029 [arXiv:hep-ph/0204244].
- [5] S. Frixione and B. R. Webber, “The MC@NLO 3.3 event generator,” arXiv:hep-ph/0612272.
- [6] S. Dawson, *Radiative corrections to Higgs boson production*, *Nucl. Phys.* **B359** (1991) 283–300.
- [7] A. Djouadi, M. Spira, and P. M. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, *Phys. Lett.* **B264** (1991) 440–446.
- [8] M. Spira, A. Djouadi, D. Graudenz, and P. M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys.* **B453** (1995) 17–82, [hep-ph/9504378].
- [9] S. Alioli, P. Nason, C. Oleari and E. Re, “NLO Higgs boson production via gluon fusion matched with shower in POWHEG,” arXiv:0812.0578 [hep-ph].
- [10] <http://mcfm.fnal.gov/>
- [11] M. Cacciari and G. P. Salam, *Dispelling the  $N^3$  myth for the  $k_T$  jet-finder*, *Phys. Lett.* **B641** (2006) 57–61, [hep-ph/0512210].
- [12] E. Boos *et al.*, “Generic user process interface for event generators,” arXiv:hep-ph/0109068.
- [13] J. Alwall *et al.*, “A standard format for Les Houches event files,” *Comput. Phys. Commun.* **176** (2007) 300 [arXiv:hep-ph/0609017].
- [14] T. Sjöstrand *et al.*, in “Z physics at LEP1: Event generators and software,” eds. G. Altarelli, R. Kleiss and C. Verzegnassi, Vol 3, pg. 327.
- [15] M. R. Whalley, D. Bourilkov and R. C. Group, “The Les Houches accord PDFs (LHAPDF) and LHAGLUE,” arXiv:hep-ph/0508110.
- [16] S. Alioli, P. Nason, C. Oleari and E. Re, “NLO vector-boson production matched with shower in POWHEG,” *JHEP* **0807**, 060 (2008) [arXiv:0805.4802 [hep-ph]].