

# HERAFitter

## Open Source QCD Fit Project

Version 0.8 (svn 1420)

S. Alekhin<sup>16,17</sup>, O. Behnke<sup>1</sup>, P. Belov<sup>1,12</sup>, M. Botje<sup>18</sup>, D. Britzger<sup>1</sup>, S. Camarda<sup>1</sup>,  
A.M. Cooper-Sarkar<sup>2</sup>, K. Daum<sup>31,32</sup>, C. Diaconu<sup>3</sup>, J. Feltesse<sup>19</sup>, A. Gizhko<sup>1</sup>,  
A. Glazov<sup>1</sup>, A. Guffanti<sup>20</sup>, M. Guzzi<sup>1</sup>, F. Hautmann<sup>13,14,15</sup>, H. Jung<sup>1</sup>, V. Kolesnikov<sup>4</sup>,  
H. Kowalski<sup>1</sup>, O. Kuprash<sup>1</sup>, A. Kusina<sup>21</sup>, S. Levonian<sup>1</sup>, K. Lipka<sup>1</sup>, B. Lobodzinski<sup>30</sup>,  
K. Lohwasser<sup>16</sup>, A. Luszczak<sup>5</sup>, B. Malaescu<sup>26</sup>, R. McNulty<sup>29</sup>, V. Myronenko<sup>1</sup>,  
S. Naumann-Emme<sup>1</sup>, K. Nowak<sup>1,24</sup>, F. Olness<sup>21</sup>, E. Perez<sup>23</sup>, H. Pirumov<sup>1</sup>, R. Plačákytė<sup>1</sup>,  
K. Rabbertz<sup>6</sup>, V. Radescu<sup>1</sup>, R. Sadykov<sup>25</sup>, G. Salam<sup>27,28</sup>, A. Sapronov<sup>4</sup>, A. Schöning<sup>10</sup>,  
T. Schörner-Sadenius<sup>1</sup>, S. Shushkevich<sup>1</sup>, W. Slominski<sup>7</sup>, H. Spiesberger<sup>22</sup>,  
P. Starovoitov<sup>1</sup>, M. Sutton<sup>8</sup>, J. Tomaszewska<sup>9</sup>, O. Turkot<sup>1</sup>, A. Vargas<sup>1</sup>, G. Watt<sup>11</sup>,  
K. Wichmann<sup>1</sup>

<sup>1</sup>Deutsches Elektronen-Synchrotron (DESY), Hamburg, Germany

<sup>2</sup>Department of Physics, University of Oxford, Oxford, United Kingdom

<sup>3</sup>CPPM, IN2P3-CNRS, Univ. Mediterranee, Marseille, France

<sup>4</sup>Joint Institute for Nuclear Research (JINR), Joliot-Curie 6, 141980, Dubna, Moscow Region, Russia

<sup>5</sup>T. Kosciuszko Cracow University of Technology

<sup>6</sup>Institut für Experimentelle Kernphysik, Karlsruhe, Germany

<sup>7</sup>Jagiellonian University, Institute of Physics, Ul. Reymonta 4, PL-30-059 Cracow, Poland

<sup>8</sup>University of Sussex, Department of Physics and Astronomy, Sussex House, Brighton BN1 9RH, United Kingdom

<sup>9</sup>Warsaw University of Technology, Faculty of Physics, Koszykowa 75, 00-662 Warsaw, Poland

<sup>10</sup>Physikalisches Institut, Universität Heidelberg, Heidelberg, Germany

<sup>11</sup>Institute for Particle Physics Phenomenology, Durham University, Durham, DH1 3LE, United Kingdom

<sup>12</sup>Current address: Department of Physics, St. Petersburg State University, Ulyanovskaya 1, 198504 St. Petersburg, Russia

<sup>13</sup>Dept. of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, United Kingdom

<sup>14</sup>Rutherford Appleton Laboratory, Chilton OX11 0QX, United Kingdom

<sup>15</sup>Dept. of Theoretical Physics, University of Oxford, Oxford OX1 3NP, United Kingdom

<sup>16</sup>Deutsches Elektronen-Synchrotron (DESY), Platanenallee 6, D15738 Zeuthen, Germany

<sup>17</sup>Institute for High Energy Physics, 142281 Protvino, Moscow region, Russia

<sup>18</sup>Nikhef, Science Park, Amsterdam, the Netherlands

<sup>19</sup>CEA, DSM/Irfu, CE-Saclay, Gif-sur-Yvette, France

<sup>20</sup>Niels Bohr Institute, University of Copenhagen, Denmark

<sup>21</sup>Southern Methodist University, Dallas, Texas

<sup>22</sup>WA ThEP, Johannes-Gutenberg-Universität Mainz, D-55099 Mainz, Germany

<sup>23</sup>CERN, European Organization for Nuclear Research, Geneva, Switzerland

<sup>24</sup>left DESY

<sup>25</sup>Joint Institute for Nuclear Research, Joliot-Curie str. 6, Dubna, 141980, Russia

<sup>26</sup>Laboratoire de Physique Nucléaire et de Hautes Energies, UPMC and Université, Paris-Diderot and CNRS/IN2P3, Paris, France

<sup>27</sup>CERN, PH-TH, CH-1211 Geneva 23, Switzerland

<sup>28</sup>LPTHE; CNRS UMR 7589; UPMC Univ. Paris 6; Paris 75252, France

<sup>29</sup>University College Dublin, Dublin 4, Ireland

<sup>30</sup>Max Planck Institut Für Physik, Werner Heisenberg Institut, Föhringer Ring 6, Muenchen

<sup>31</sup>Fachbereich C, Universität Wuppertal, Wuppertal, Germany

<sup>32</sup>Rechenzentrum, Universität Wuppertal, Wuppertal, Germany

Received: date / Accepted: date

**Abstract** The paper presents the HERAFitter project which provides a framework for Quantum Chromodynamics (QCD) analyses related to the proton structure. The main processes sensitive to the Parton Distribution Functions (PDFs) of the proton are Deep-Inelastic-Scattering in  $ep$  collisions at HERA and Drell Yan, jet and top quark production in  $pp$  ( $p\bar{p}$ ) collisions at the LHC (Tevatron). Data of recent measurements are included into HERAFitter and can be used for PDF determination based on the concept of the factorisable nature of the cross sections of hard scattering measurements into

process dependent partonic scattering and universal PDFs. HERAFitter provides a comprehensive choice of options in the treatment of the experimental data uncertainties, a large number of theoretical and methodological options through interfaces to external software packages which are described here.

**Keywords** PDFs · QCD · Fit

## Contents

1	Introduction	2
2	HERAFitter Structure	2
3	Theoretical Input	4
3.1	Deep Inelastic Scattering Formalism and Schemes	4
3.2	Drell Yan processes in $pp$ or $p\bar{p}$ collisions	5
3.3	Jet production in $ep$ and $pp$ or $p\bar{p}$ collisions	6
3.4	Top-quark production in $pp$ and $p\bar{p}$ collisions	6
4	Computational Techniques	6
4.1	$k$ -factor Technique	6
4.2	Fast Grid Techniques	7
4.3	Performance Optimisation	8
5	Fit Methodology	8
5.1	Functional Forms for PDF parametrisation	8
5.2	$\chi^2$ representation	9
5.3	Treatment of the Experimental Uncertainties	10
5.4	Treatment of the Theoretical Input Parameters	11
5.5	Bayesian Reweighting Techniques	11
6	Alternatives to DGLAP formalism	12
6.1	DIPOLE models	12
6.2	Transverse Momentum Dependent PDFs with CCFM	12
6.3	Diffractive PDFs	13
7	Application of HERAFitter	13
8	Summary	14

## 1 Introduction

The discovery of the Higgs boson [1, 2] and extensive searches for signals of new physics at the LHC demands accurate precision of the Standard Model (SM) predictions for hard scattering processes in hadron-hadron collisions. The most common approach to calculate the SM cross sections for such reactions is to use collinear factorisation in perturbative QCD (pQCD) [3]:

$$\sigma(\alpha_s, \mu_R, \mu_F) = \sum_{a,b} \int_0^1 dx_1 dx_2 f_a(x_1, \alpha_s, \mu_F) f_b(x_2, \alpha_s, \mu_F) \times \hat{\sigma}^{ab}(x_1, x_2; \alpha_s, \mu_R, \mu_F). \quad (1)$$

Here the cross section  $\sigma$  for any hard-scattering inclusive process  $ab \rightarrow X + \text{all}$  is expressed as a convolution of Parton Distribution Functions (PDFs)  $f_a$  and  $f_b$  with the partonic cross section  $\hat{\sigma}^{ab}$ . The PDFs represent the probability of finding a specific parton  $a$  ( $b$ ) in the first (second) proton carrying a fraction  $x_1$  ( $x_2$ ) of its momentum. Indices  $a$  and  $b$  in the Eq. 1 indicates the various kinds of partons, i.e. gluons, quarks and antiquarks of different flavours, that are considered as the constituents of the proton. Both the PDFs

and the partonic cross section depend on the strong coupling  $\alpha_s$ , and the factorisation and renormalisation scales,  $\mu_F$  and  $\mu_R$ , respectively. The partonic cross sections are calculable in pQCD whereas PDFs cannot be computed analytically in QCD, they must rather be determined from measurement. PDFs are assumed to be universal such that different scattering reactions can be used to constrain them [4, 5].

Measurements of the inclusive Neutral Current (NC) and Charged Current (CC) Deep-Inelastic-Scattering (DIS) at the  $ep$  collider HERA provide crucial information for determining the PDFs. For instance, the gluon density relevant for calculating the dominant gluon-gluon fusion contribution to Higgs production at the LHC can be accurately determined at low and medium  $x$  solely from the HERA data. Many processes in  $pp$  and  $p\bar{p}$  collisions at LHC and Tevatron, respectively, probe PDFs in the kinematic ranges, complementary to the DIS measurements. Therefore inclusion of the LHC and Tevatron data in the QCD analysis of the proton structure provide additional constraints on the PDFs, improving either their precision, or providing important information of the correlations of PDF with the fundamental QCD parameters like strong coupling or quark masses. In this context, the processes of interest at hadron colliders are Drell Yan (DY) production,  $W$  asymmetries, associated production of  $W$  or  $Z$  bosons and heavy quarks, top quark, jet and prompt photon production.

The open-source QCD platform HERAFitter encloses the set of tools necessary for a comprehensive global QCD analysis of hadron-induced processes even at the early stage of the experimental measurement. It has been developed for determination of PDFs and extraction of fundamental QCD parameters such as the heavy quark masses or the strong coupling constant. This platform also provides the basis for comparisons of different theoretical approaches and can be used for direct tests of the impact of new experimental data in the QCD analyses.

The outline of this paper is as follows. The structure and overview of HERAFitter is presented in section 2. Section 3 discusses the various processes and corresponding theoretical calculations performed in the DGLAP [11–15] formalism that are available in HERAFitter. Section 4 presents various techniques employed by the theory calculations used in HERAFitter. Section 5 elucidates the methodology of determining PDFs through fits based on various  $\chi^2$  definitions used in the minimisation procedure. Alternative approaches to the DGLAP formalism are presented in section 6. Specific applications of the package are given in section 7 and the summary is presented in section 8.


## 2 HERAFitter Structure

The processes that are currently available in HERAFitter framework are listed in Tab. 1. The functionality of HERAFitter

Data	Process	Reaction	Theory calculations, schemes	
HERA	DIS NC	$ep \rightarrow eX$	TR', ACOT	113
			ZM (QCDNUM)	114
			FFN (OPENQCDRAD, QCDNUM),	115
			TMD (uPDFevolv)	116
	DIS CC	$ep \rightarrow \nu_e X$	ACOT, ZM (QCDNUM)	117
Fixed Target	DIS NC	$ep \rightarrow eX$	FFN (OPENQCDRAD, QCDNUM)	118
			TMD (uPDFevolv)	119
	DIS heavy quarks	$ep \rightarrow e\bar{c}X$ , $ep \rightarrow e\bar{b}X$	NLOJet++ (fastNLO)	120
			ZM (QCDNUM), TR', ACOT, FFN (OPENQCDRAD, QCDNUM)	121
Tevatron, LHC	DIS NC	$pp(\bar{p}) \rightarrow l\bar{l}X$ , $pp(\bar{p}) \rightarrow l\nu X$	ZM (QCDNUM), TR', ACOT	122
	Drell Yan	$pp(\bar{p}) \rightarrow l\bar{l}X$ , $pp(\bar{p}) \rightarrow l\nu X$	MCfM (APPLGRID)	123
	top pair	$pp(\bar{p}) \rightarrow t\bar{t}X$	MCfM (APPLGRID), HATHOR	124
	single top	$pp(\bar{p}) \rightarrow tlvX$ , $pp(\bar{p}) \rightarrow tX$ , $pp(\bar{p}) \rightarrow tWX$	MCfM (APPLGRID)	125
	jets	$pp(\bar{p}) \rightarrow \text{jets}X$	NLOJet++ (APPLGRID), NLOJet++ (fastNLO)	126
LHC	DY+heavy quarks	$pp \rightarrow VhX$	MCfM (APPLGRID)	127

**Table 1** The list of processes available in the HERAFitter package. The references for the individual calculations and their implementations are given in the text.

is schematically illustrated in Fig. 1 and it can be divided in four main blocks:



**Fig. 1** Schematic structure of the HERAFitter program.

**Input data:** The relevant cross section measurements from the various processes are stored internally in HERAFitter with the full information on their uncorrelated and correlated uncertainties. HERA data sets are the basis of any proton PDF extraction, and they are used by all global PDF groups [6–10]. Additional measurements provide

constraints to the sea flavour decomposition, such as the new results from the LHC, as well as constraints to PDFs in the kinematic phase-space regions where HERA data is not measured precisely, such as the high  $x$  region for the gluon and valence quark distributions from Tevatron and fixed target experiments..

**Theory predictions:** Predictions for cross section of different processes are obtained using the factorisation approach (Eq. 1). The PDFs are parametrised at a starting input scale  $Q_0^2$  by a chosen functional form with a set of free parameters  $\mathbf{p}$ . These PDFs are then evolved from  $Q_0^2$  to the scale of the measurement using the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) [11–15] evolution equations (as implemented in QCDNUM [16]), CCFM [17–20] or dipole models [21–23] and then convoluted with the hard parton cross sections calculated using a relevant theory program (as listed in Tab. 1).

**QCD fit:** The PDFs are extracted from a least square fit by minimising the  $\chi^2$  function with respect to free parameters. The  $\chi^2$  function is formed from the input data and the theory prediction. The  $\chi^2$  is minimised iteratively with respect to the PDF parameters using the MINUIT [24] program. Various choices of accounting for the experimental uncertainties are employed in HERAFitter, either using a nuisance parameter method for the correlated systematic uncertainties, or a covariance matrix method (see details in section 5.2). In addition, HERAFitter allows to study different statistics assumptions for the distributions of the systematic uncertainties (i.e. Gauss or log-normal) [25].

**Results:** The resulting PDFs are provided in a format ready to be used by the LHAPDF library [26, 27] or by TMDlib [28]. HERAFitter drawing tools can be used to display the PDFs with the uncertainty at a chosen scale. A first set of PDFs extracted by HERAFitter is HERAPDF1.0 [29], shown in Fig. 2, which is based on HERA I data. Since then several other PDF sets were produced within the HERA and LHC collaborations. In addition to the PDF display, the visual comparison of data used in the fit to the theory predictions are also produced. In Fig. 3, a comparison of inclusive NC data from the HERA I running period with predictions based on HERAPDF1.0. It also illustrates the comparison to the theory predictions which are adjusted by the systematic uncertainty shifts when using the nuisance parameter method that accounts for correlated systematic uncertainties. As an additional consistency check between data and the theory predictions, pull information, defined as the difference between data and prediction divided by the uncorrelated uncertainty of the data, is displayed in units of sigma shifts for each given data bin.

The HERAFitter project provides a versatile environment for benchmarking studies and a flexible platform for



## Fixed Flavour Number (FFN):

In this scheme [39–41] only the gluon and the light quarks are considered as partons within the proton and massive quarks are produced perturbatively in the final state. The lowest order process is the fusion of a gluon in the proton with a boson from the lepton to produce a heavy quark and an antiquark. In HERAFitter this scheme can be accessed via the QCDNUM implementation or through the interface to the open-source code OPENQCDRAD (as implemented by the ABM group) [42]. Through QCDNUM, the calculation of the heavy quark contributions to DIS structure functions are available at Next-to-Leading-Order (NLO) and only electromagnetic exchange contributions are taken into account. Through the ABM implementation the heavy quark contributions to CC structure functions are available and, for the NC case, the QCD corrections to the massive Wilson coefficients at Next-to-Next-to Leading Order (NNLO) are provided at the best currently known approximation [43]. The ABM implementation also includes the running mass definition of the heavy quark mass [44]. The running mass scheme has the advantage of reducing the sensitivity of the DIS cross sections to higher order corrections, and improving the theoretical precision of the mass definition.

## General-Mass Variable-Flavour Number (GM-VFN):

In this scheme [45], heavy quark production is treated for  $Q^2 \leq m_h^2$  in the FFN scheme and for  $Q^2 \gg m_h^2$  in a fully massive scheme. The recent series of PDF groups that use this scheme are MSTW, CT(CTEQ), NNPDF, and HERAPDF. HERAFitter implements different variants of the GM-VNS scheme and they are presented below:

- **GM-VFN Thorne-Roberts scheme:** The Thorne-Roberts (TR) scheme [46] was designed to provide a smooth transition from the massive FFN scheme at low scales  $Q^2 < m_h^2$  to the massless ZM-VFNS scheme at high scales  $Q^2 \gg m_h^2$ . However, the original version was technically difficult to implement beyond NLO, and was updated to the TR' scheme [47] which is simpler (and closer to the ACOT-scheme, see below). There are two different variants of the TR' schemes: TR' standard (as used in MSTW PDF sets [6, 47]) and TR' optimal [48], with a smoother transition across the heavy quark threshold region. Both of these variants are accessible within the HERAFitter package at LO, NLO and NNLO.
- **GM-VFN ACOT scheme:** The Aivazis-Collins-Olness-Tung scheme belongs to the group of VFN factorisation schemes that use the renormalization method of Collins-Wilczek-Zee (CWZ) [49]. This scheme unifies the low scale  $Q^2 < m_h^2$  and high scale  $Q^2 > m_h^2$  regions; thus, it provides a smooth interpolation across the full energy regime. Within the ACOT package, different variants of the ACOT scheme are avail-

able: ACOT-Full, S-ACOT- $\chi$ , ACOT-ZM,  $\overline{\text{MS}}$  at LO and NLO. For the longitudinal structure function higher order calculations are also available. The ACOT-Full implementation takes into account the quark masses and it reduces to ZM  $\overline{\text{MS}}$  scheme in the limit of masses going to zero, but it has the disadvantage that it is computationally intensive (addressed in section 4).

Calculations of higher-order electroweak corrections to DIS scattering at HERA are available in HERAFitter, performed in the on-shell scheme where the gauge bosons masses  $M_W$  and  $M_Z$  are treated symmetrically as basic parameters together with the top, Higgs and fermion masses. These electroweak corrections are based on the EPRC package [50]. The code provides the running of  $\alpha$  using the most recent parametrisation of the hadronic contribution to  $\Delta_\alpha$  [51], as well as an older version from Burkhard [52].

## 3.2 Drell Yan processes in $pp$ or $p\bar{p}$ collisions

The Drell Yan (DY) process provides further valuable information about PDFs. In  $pp$  and  $p\bar{p}$  scattering, the  $Z/\gamma$  and  $W$  production probe bi-linear combinations of quarks. Complementary information on the different quark densities can be obtained from the  $W$  asymmetry ( $d$ ,  $u$  and their ratio), the ratio of the  $W$  and  $Z$  cross sections (sensitive to the flavor composition of the quark sea, in particular to the  $s$  density), and associated  $W$  and  $Z$  production with heavy quarks (sensitive to  $s$  and  $c$  quark densities).

Presently, the predictions for Drell-Yan and  $W$  and  $Z$  production are known to NNLO and  $W$ ,  $Z$  in association with heavy flavour quarks are known to NLO. There are several possibilities for obtaining the theoretical predictions for DY production in HERAFitter. At LO an analytic calculation is available within the package and described below:

The LO DY triple differential cross section in invariant mass  $M$ , boson rapidity  $y$  and Centre-of-Mass lepton Scattering (CMS) angle  $\cos \theta$ , for NC, can be written as [53, 54]:

$$\frac{d^3\sigma}{dM dy d\cos\theta} = \frac{\pi\alpha^2}{3MS} \sum_q P_q [f_q(x_1, Q^2) f_{\bar{q}}(x_2, Q^2) + (q \leftrightarrow \bar{q})], \quad (4)$$

where  $S$  is the squared CMS beam energy,  $x_{1,2} = \frac{M}{\sqrt{S}} \exp(\pm y)$ ,  $f_q(x_1, Q^2)$  is the parton number density, and  $P_q$  is a partonic cross section.

The expression for CC scattering has a form:

$$\frac{d^3\sigma}{dM dy d\cos\theta} = \frac{\pi\alpha^2}{48S \sin^4 \theta_W} \frac{M^3 (1 - \cos \theta)^2}{(M^2 - M_W^2) + \Gamma_W^2 M_W^2} \sum_{q_1, q_2} V_{q_1 q_2}^2 f_{q_1}(x_1, Q^2) f_{q_2}(x_2, Q^2), \quad (5)$$



where  $V_{q_1 q_2}$  is the CKM quark mixing matrix and  $M_W$  and  $\Gamma_W$  are the  $W$  boson mass and decay width.

The simple form of these expressions allows the calculation of integrated cross sections without the use of Monte-Carlo (MC) techniques which often introduce statistical fluctuations. In both NC and CC expressions PDFs factorise as functions dependent only on boson rapidity  $y$  and invariant mass  $M$ , while the integral in  $\cos \theta$  can be computed analytically.

The NLO and NNLO calculations are highly demanding in terms of the computing power and time, and  $k$ -factor or fast grid techniques must be employed (see section 4 for details), interfaced to programs such as MCFM [55–57], available for NLO calculations, or FEWZ [58] and DNNLO [59] for NLO and NNLO.

### 3.3 Jet production in $ep$ and $pp$ or $p\bar{p}$ collisions

Jet production at high transverse momentum is sensitive to the high- $x$  gluon PDF (see e.g. [6]) and can thus increase the precision of the gluon PDF determination, which is particularly important for the Higgs production and searches for new physics. Jet production cross sections are only currently known to NLO, although NNLO calculations are now quite advanced [60–62]. Within HERAFitter programs such MCFM and NLOJet++ [63, 64] may be used for the calculation of jet production. Similarly to the DY case, the calculation is very demanding in terms of computing power. Therefore, to allow the possibility to include  $ep$ ,  $pp$  or  $p\bar{p}$  jet cross section measurements in QCD fits in order to extract PDFs and  $\alpha_s$ , the fast grid techniques are used (see section 4).

### 3.4 Top-quark production in $pp$ and $p\bar{p}$ collisions

Top-quark pairs ( $t\bar{t}$ ) are produced at hadron colliders dominantly via  $gg$  fusion and  $q\bar{q}$  annihilation. Measured  $t\bar{t}$  cross sections provide additional constraints in particular on the gluon density at medium to high values of  $x$ , on  $\alpha_s$  and on the top-quark mass,  $m_t$ . Single top quarks are produced via electroweak interactions and single-top cross sections can be used, for example, to probe the ratio of the  $u$  and  $d$  densities in the proton as well as the  $b$ -quark PDF. Precise predictions for the total  $t\bar{t}$  cross section have become available to full NNLO recently [65]. They can be used within HERAFitter via an interface to the program HATHOR [66]. Differential  $t\bar{t}$  cross sections and predictions for single-top production can be used with HERAFitter at NLO accuracy from MCFM [57, 67–70] in combination with fast grid techniques.

## 4 Computational Techniques

More precise measurements require theoretical predictions with equally improved accuracy in order to maximize their impact in PDF fits. Perturbative calculations, however, get more and more involved with increasing number of Feynman diagrams at the each higher order. Nowadays even the most advanced perturbative techniques in combination with recent computing hardware do not lead to sufficiently small turn-around times. The direct inclusion of computationally demanding higher-order calculations into iterative fits therefore is not possible. Relying on the fact that a full repetition of the perturbative calculation for arbitrary changes in input parameters is not necessary at each iteration step, two methods have been developed to resolve this problem: the techniques of  $k$ -factors and *fast grids*. Both are available in HERAFitter and described as follows.


### 4.1 $k$ -factor Technique

$k$ -factors are defined as the ratio of the prediction of a higher-order (slow) pQCD calculation to a lower-order (fast) calculation. Because the  $k$ -factors depend on the phase space probed by the measurement they have to be stored into a table in dependence of the relevant kinematic variables. Before the start of a fitting procedure the table of  $k$ -factors has to be computed once for a given PDF with the time consuming higher-order code. In subsequent iteration steps the theory prediction is derived from the fast lower-order calculation multiplied by the pre-tabulated  $k$ -factors.

However, this procedure neglects the fact that the  $k$ -factors are process dependent and, as a consequence, they have to be re-evaluated for the newly determined PDF at the end of the fit in order to check for any changes. Usually, the fit is repeated until input and output  $k$ -factors have converged. In summary, this technique avoids to iterate the higher-order calculation at each step, but still requires a couple of repetitions depending on the analysis.

– In DIS, appropriate treatments of the heavy quarks require computationally slow calculations. For this purpose, “FAST” heavy flavour schemes are implemented in HERAFitter with  $k$ -factors defined as the ratio of calculations at the same perturbative order but for massive vs. massless quarks, e.g. NLO (massive)/NLO (massless). In the HERAFitter implementation, these  $k$ -factors are calculated only for the starting PDF and hence, the “FAST” heavy flavour schemes should only be used for quick checks, i.e. full heavy flavour schemes are recommended (with an exception of ACOT case where, due to long computation time, the  $k$ -factors are used in the default settings).

This “FAST” method was employed in the QCD fits to the HERA data shown in Fig. 4. In this case, the ACOT scheme was used as a cross check of the central results [29].



**Fig. 4** Overview showing the  $u$ – and  $d$ -valence, the total sea (scaled), and gluon (scaled) PDFs of the NLO HERAPDF1.0 set [29] with their total uncertainty at the scale of  $Q^2 = 10 \text{ GeV}^2$  obtained using the TR’ scheme and compared to the PDFs obtained with the ACOT scheme using the  $k$ -factor technique (red).


## 4.2 Fast Grid Techniques

Fast grid techniques exploit the fact that iterative PDF fitting procedures do not impose completely arbitrary changes to the types and shapes of the parameterised functions that represent each PDF. Instead, it can be assumed that a generic PDF can be approximated by a set of interpolating functions with a sufficient number of strategically well-chosen support points. The quality, i.e. the accuracy of this approximation, can be tested and optimised by a number of means, the simplest one being an increase in the number of support points. Ensuring an approximation bias that is negligibly small for all practical purposes this method can be used to perform the time consuming higher-order calculation (see Eq. 1) only once for the set of interpolating functions. The repetition of a cross section evaluation for a particular PDF set then is very fast and implies only sums over the set of interpolators multiplied by factors depending on the respective PDF. The described approach applies equally to processes involving one or two hadrons in the initial state as well as to the renormalisation and factorisation scale dependence in the convolution of the PDFs with the partonic cross section.

This technique was pioneered in the fastNLO project [71] to facilitate the inclusion of notoriously time consuming jet cross sections at NLO into PDF fits. The APPLGRID [72]


package extended first a similar methodology to DY production. While differing in their interpolation and optimisation strategies, both packages construct tables with grids for each bin of an observable in two steps: In the first step the accessible phase space in the parton momentum fractions  $x$  and the renormalisation and factorisation scales  $\mu_R$  and  $\mu_F$  is explored in order to optimize the table size. The second step consists of the actual grid construction and filling for the requested observables. Higher-order cross sections can then be restored very efficiently from the preproduced grids while varying externally provided PDF sets,  $\mu_R$  and  $\mu_F$ , or the strong coupling  $\alpha_s(Q)$ . The approach can in principal be extended to arbitrary processes, but requires to establish an interface between the higher-order theory programs and the fast interpolation frameworks. Work in that direction is ongoing for both packages. They are described in some more detail in the following:

- The fastNLO project [71] has been interfaced to the NLOJet++ program [63] for the calculation of jet production in DIS [73] as well as 2- and 3-jet production in hadron-hadron collisions at NLO [64, 74]. To demonstrate the applicability to higher-orders, threshold corrections at 2-loop order, which approximate the NNLO for the inclusive jet cross section, have been included into the framework as well [75] following Ref. [76]. The latest version of fastNLO [77] allows creation of tables where renormalisation and factorisation scales can be chosen freely as a function of two pre-defined observables, e.g. jet transverse momentum  $p_\perp$  and  $Q$  for DIS. fastNLO can be obtained from [78], where numerous precalculated grid tables for jet cross sections can be downloaded as well. Dedicated fastNLO libraries and tables required for comparison to particular datasets are included in the HERAFitter package. In this case, the evaluation of the strong coupling constant is taken consistently with the PDF evolution from the QCDNUM code. The interface to the fastNLO tables from within HERAFitter was used in a recent CMS analysis, where the impact on the extraction of the PDFs from the inclusive jet cross section is investigated [34]. The influence on the gluon density by the CMS inclusive jet data is illustrated in Fig. 5.
- The APPLGRID package [72], which is also available from [79], in addition to the jet cross sections from NLOJet++ in  $pp(\bar{p})$  and DIS processes, implements the calculations of DY production. The look-up tables (also called grids) can be generated with modified versions of the MCFM parton level generator for DY [55–57]. Alternative values of the strong coupling constant as well as a posteriori variation of the renormalisation and factorisation scales can be freely chosen in the calculation of the theory predictions with the APPLGRID tables. For NNLO predictions in HERAFitter  $k$ -factors can be applied.



**Fig. 5** The gluon density as a function of  $x$  as derived from HERA inclusive DIS data alone (cyan) and in combination with CMS inclusive jet data from 2011 (blue hatched) [34], where bands represent the total uncertainty of the PDFs. The PDFs are shown at the starting scale  $Q^2 = 1.9 \text{ GeV}^2$ .

The HERAFitter interface to APPLGRID was used by the ATLAS collaboration to extract the strange quark density of the proton from  $W$  and  $Z$  cross sections [30]. An illustration of ATLAS PDFs extracted using the  $k$ -factors is shown in Fig. 6 together with the comparison to global PDF sets CT10 [7] and NNPDF2.1 [8].



**Fig. 6** The strange anti-quark density versus  $x$  for the ATLAS epWZ free  $\bar{s}$  NNLO fit (magenta band) compared to predictions from NNPDF2.1 (blue hatched) and CT10 (green hatched) at  $Q^2 = 1.9 \text{ GeV}^2$ . The figure is taken from [30].

### 4.3 Performance Optimisation

An important factor for a feasible QCD fit which is performed by iterative  $\chi^2$  minimisation, is performance in terms of how long a calculation takes for each given data point. The performance of the HERAFitter code is greatly improved with several special built-in options including the  $k$ -factor techniques (see section 4) and the grid techniques for the fast calculation of cross sections of particular processes for arbitrary sets of PDFs. There are also cache options, fast evolution kernels, and usage of the OpenMP (Open Multi-Processing) interface which allows parallel applications of some of the heavy flavour scheme theory predictions in DIS.

## 5 Fit Methodology

There is a considerable number of choices available when performing a QCD fit analysis (i.e. functional parametrisation form, choice for heavy quarks mass values, alternative theoretical calculations, method of minimisation, interpretation of uncertainties etc.). It is desirable to be able to discriminate or quantify the effect of the chosen ansatz, ideally within a common framework, and HERAFitter is optimally designed for such tests. The methodology employed by HERAFitter relies on a flexible and modular framework that allows for independent integration of the state-of-the-art techniques, either related to the inclusion of a new theoretical calculation, or to new approaches to treat uncertainties.

In this section we briefly describe the available options in HERAFitter ranging from the functional form used to parametrise PDFs and the choice of the form of the  $\chi^2$  function, to different methods to assess the experimental uncertainties on extracted PDFs.

In addition, as an alternative approach to a complete QCD fit, the Bayesian reweighting method, which is also available in HERAFitter, is described in this section.

### 5.1 Functional Forms for PDF parametrisation

The PDFs are parametrised at a starting scale chosen by the user. In HERAFitter various functional forms to parametrise PDFs can be tested:

**Standard Polynomials:** The term refers to using a simple polynomial to interpolate between the low and high  $x$  regions:

$$xf(x) = Ax^B(1-x)^C P_i(x), \quad (6)$$

The standard polynomial form is most commonly used by PDF groups. The parametrised PDFs at HERA are the valence distributions  $xu_v$  and  $xd_v$ , the gluon distribution  $xg$ , and the  $u$ -type and  $d$ -type sea  $x\bar{U}$ ,  $x\bar{D}$ , where



$x\bar{U} = x\bar{u}$ ,  $x\bar{D} = x\bar{d} + x\bar{s}$  at the starting scale chosen below the charm mass threshold. The  $P_i(x)$  for the HERAPDF [29] style takes the simple Regge-inspired form  $(1 + \varepsilon\sqrt{x} + Dx + Ex^2)$  with additional constraints relating to the flavour decomposition of the light sea. For the CTEQ style,  $P_i(x)$  takes the form  $e^{a_3x}(1 + e^{a_4x} + e^{a_5x^2})$ . QCD number and momentum sum-rules are used to determine the normalisations  $A$  for the valence and gluon distributions. The sum-rules can be evaluated analytically.

**Bi-Log-Normal Distributions:** This parametrisation is motivated by multiparticle statistics [25] and holds the following functional form:

$$xf(x) = x^{p-b\log(x)}(1-x)^{q-\log(1-x)}. \quad (7)$$

This function can be regarded as a generalisation of the standard functional form described above. In order to satisfy the QCD sum rules this parametric form requires numerical integration.


**Chebyshev Polynomials:** A flexible Chebyshev polynomial based parametrisation can be used for the gluon and sea densities. The polynomials use  $\log x$  as an argument to emphasize the low  $x$  behavior. The PDFs are multiplied by a  $(1-x)$  term to ensure that they vanish as  $x \rightarrow 1$ . The resulting parametric form is

$$xg(x) = A_g(1-x) \sum_{i=0}^{N_g-1} A_{gi} T_i \left( -\frac{2\log x - \log x_{\min}}{\log x_{\min}} \right), \quad (8)$$

$$xS(x) = (1-x) \sum_{i=0}^{N_S-1} A_{Si} T_i \left( -\frac{2\log x - \log x_{\min}}{\log x_{\min}} \right). \quad (9)$$


Here the sum runs over  $i$  up to  $N_{g,S} = 15$  order Chebyshev polynomials of the first type  $T_i$  for the gluon,  $g$ , and sea-quark,  $S$ , density, respectively. The normalisation  $A_g$  is given by the momentum sum rule. The advantages of this parametrisation are that the momentum sum rule can be evaluated analytically and that for  $N \geq 5$  the fit quality is already similar to the standard Regge-inspired parametrisation with a similar number of parameters. Such a study of the parametrisation uncertainty at low Bjorken  $x \leq 0.1$  for PDFs was presented in [80]. Figure 7 shows that the accuracy of the HERA data allows the gluon density to be determined in the kinematic range of  $0.0005 \leq x \leq 0.05$  with a reduced parametrisation uncertainty.

**External PDFs:** HERAFitter provides the possibility to access external PDF sets, which can be used to construct theoretical predictions for the various processes of interest as implemented in HERAFitter. This is possible via an interface to LHAPDF [26, 27] which provides access to the global PDF sets available at LO, NLO or NNLO evolved either locally through the HERAFitter or taken as provided by the LHAPDF grids. Figure 8 is produced



**Fig. 7** The gluon density is shown at the starting scale. The black lines correspond to the error band of the gluon distribution using a standard parameterization and it is to be compared to the case of the Chebyshev parameterization [80].

with the drawing tools available in HERAFitter and illustrates the PDFs accessed from LHAPDF.



**Fig. 8** Gluon density as extracted by various PDF groups at the scale of  $Q^2 = 2 \text{ GeV}^2$ , plotted using the drawing tools from HERAFitter.

## 5.2 $\chi^2$ representation

The PDF parameters are extracted from a  $\chi^2$  minimisation process. The construction of the  $\chi^2$  accounts for the experimental uncertainties. There are various forms that can be used to represent the experimental uncertainties, e.g. using covariance matrices or providing nuisance parameters for

dependence of each systematic source on the data point. In addition, there are various methods to deal with correlated systematic (or statistical) uncertainties (e.g. different scaling options, etc.). Here we summarise the options available in HERAFitter.

**Covariance Matrix Representation:** For a data point  $\mu_i$  with a corresponding theory prediction  $m_i$ , the  $\chi^2$  function for the case when experimental uncertainties are given as a covariance matrix  $C_{i,j}$  over data bins  $i$  and  $j$ , can be expressed in the following form:

$$\chi^2(m) = \sum_{i,j} (m_i - \mu_i) C_{i,j}^{-1} (m_j - \mu_j). \quad (10)$$

The covariance matrix can be decomposed into statistical, uncorrelated and correlated systematic contributions:

$$C_{ij} = C_{ij}^{\text{stat}} + C_{ij}^{\text{uncor}} + C_{ij}^{\text{sys}}. \quad (11)$$

With this representation the particular effect of a particular source of the systematic uncertainty can no longer be distinguished from other uncertainties.

**Nuisance Parameters Representation:** The  $\chi^2$  form is expressed as

$$\chi^2(m, b) = \sum_i \frac{[\mu_i - m_i (1 - \sum_j \gamma_j^i b_j)]^2}{\delta_{i,\text{unc}}^2 m_i^2 + \delta_{i,\text{stat}}^2 \mu_i m_i (1 - \sum_j \gamma_j^i b_j)} + \sum_j b_j^2, \quad (12)$$

where  $\mu_i$  is the measured central value at a point  $i$  with relative statistical  $\delta_{i,\text{stat}}$  and relative uncorrelated systematic uncertainty  $\delta_{i,\text{unc}}$ . Further,  $\gamma_j^i$  quantifies the sensitivity of the measurement  $\mu_i$  at the point  $i$  to the correlated systematic source  $j$ . The function  $\chi^2$  depends in addition on the set of systematic nuisance parameters  $b_j$ . This definition of the  $\chi^2$  function assumes that systematic uncertainties are proportional to the central prediction values (multiplicative errors), whereas the statistical uncertainties scale with the square root of the expected number of events. The systematic shift nuisance parameters  $b_j$  as well as the PDF parameters are free parameters of the fit. The fit determines the best PDF parameters to the data taking into account correlated systematic shifts of the data.

**Mixed Form Representation:** It can happen that various parts of the systematic and statistical uncertainties are stored in different forms. A situation can be envisaged when the correlated systematic experimental uncertainties are provided as nuisance parameters, but the statistical bin-to-bin correlations are given in the form of a covariance matrix. HERAFitter offers the possibility to include such information, when provided, as well as any other mixed form of treating statistical, uncorrelated and correlated systematic uncertainties.

## 5.3 Treatment of the Experimental Uncertainties

Three distinct methods for propagating experimental uncertainties to PDFs are implemented in HERAFitter and reviewed here: the Hessian, Offset, and Monte Carlo method.

**Hessian method:** The technique developed by [81] presents an estimate of PDF uncertainties reflecting the experimental precision of data used in the QCD fit by examining the behaviour of  $\chi^2$  in the neighborhood of the minimum. This is known as the Hessian or error matrix method. The Hessian matrix is built by the second derivatives of  $\chi^2$  at the minimum. The Hessian matrix is diagonalised through an iterative procedure and its PDF eigenvectors are obtained, which correspond to the orthogonal sources of uncertainties on the obtained PDF.


**Offset method:** Another method to propagate the correlated systematic experimental uncertainties from the measurements to PDFs [82] is Offset method. It uses also the  $\chi^2$  function for the central fit for which only uncorrelated uncertainties are taken into account to get the best PDF parameters. The goodness of fit can no longer be judged from the  $\chi^2$  since correlated uncertainties are ignored. The correlated systematic uncertainties of the data are then used to estimate the errors on the PDF parameters as follows. The cross section is varied by one sigma shift from the central value for each systematic source and the fit is performed. This is done for both positive and negative one sigma shifts. After this has been done for all sources the resulting deviations of each of these fits from the central PDF parameters are added in quadrature.

In most cases, the uncertainties estimated through the offset method are larger than those from the Hessian method, as the offset method does not use the information on correlated systematic uncertainties optimally.

**Monte Carlo method:** The PDF uncertainties can be estimated using a Monte Carlo technique [83, 84]. The method consists in preparing replicas of data sets by allowing the central values of the cross sections to fluctuate within their systematic and statistical uncertainties taking into account all point-to-point correlations. The preparation of the data is repeated for large  $N$  ( $> 100$  times) and for each of these replicas a QCD fit is performed to extract the PDF set. The PDF central values and uncertainties are estimated using the mean values and standard deviations over the replicas.

The MC method was checked against the standard error estimation of the PDF uncertainties as used by the Hessian method. A good agreement was found between the methods when employing for the MC approach the assumption that uncertainties (statistical and systematic) follow Gaussian distribution [25]. This comparison is il-

illustrated in Fig. 9. Similar findings were observed also in the MSTW global analysis [85].



**Fig. 9** Comparison between the standard error calculations as employed by the Hessian approach (black lines) and the MC approach assuming Gaussian distribution for uncertainty distributions, shown here for each replica (green lines) together with the evaluated standard deviation (red lines) [25].

Usage of the nuisance parameters for the experimental uncertainty treatment in QCD fits is quite common and has an advantage of the flexible assessment of such uncertainties on PDFs. Generally, the experimental uncertainties are symmetrised when QCD fits are performed, however often the provided uncertainties are rather asymmetric. HERAFitter provides the possibility to use asymmetric systematic uncertainties. The technical implementation relies on the assumption that asymmetric uncertainties can be described by a parabolic function, as given below:

$$f_i(b_j) = \omega_j^i b_j^2 + \gamma_j^i b_j, \quad (13)$$

where the coefficients  $\omega_j^i$ ,  $\gamma_j^i$  are defined as up and down shifts of the cross sections to a nuisance parameter,  $S_{ij}^\pm$ ,

$$\omega_j^i = \frac{1}{2} (S_{ij}^- + S_{ij}^+), \quad \gamma_j^i = \frac{1}{2} (S_{ij}^- - S_{ij}^+) \quad (14)$$

For this case the definition of the  $\chi^2$  from Eq. 12 is extended with the parabolic approximation for asymmetric uncertainties, such that the expected cross section is adjusted to be

$$m_i (1 - \sum_j \gamma_j^i b_j) \rightarrow m_i \left( 1 - \sum_j b_j (\omega_j^i b_j + \gamma_j^i) \right). \quad (15)$$

The minimisation is performed using fixed number of iterations (typically ten), with rapid convergence.

## 5.4 Treatment of the Theoretical Input Parameters

The results of a QCD fit depend not only on the input data but also on the input theoretical ansatz, which is also uncertain. Nowadays, modern PDF sets try to address the impact of the choices of theoretical parameters by providing alternative PDFs with different choices of the mass of the charm quarks  $m_c$ , mass of the bottom quarks  $m_b$  and the value of  $\alpha_s(M_Z)$ , etc. Another important input is the choice of the functional form for the PDFs at the starting scale and indeed the value of the starting scale itself. HERAFitter provides a platform in which such choices can readily be varied within a common framework.

## 5.5 Bayesian Reweighting Techniques

As an alternative to a complete QCD fit, the reweighting method (Bayesian Reweighting) is available in HERAFitter. Because no fit is performed, the method provides a fast estimate of the impact of new data on PDFs. The original suggestion [83] was developed by the NNPDF collaboration [86, 87] and later extended [85] to work not only on the NNPDF replicas, but also on the eigenvectors provided by most PDF groups.

The Bayesian Reweighting technique uses the PDF probability distributions which are modified with weights to account for the difference between theory predictions and new data. In the NNPDF method the PDFs are constructed as ensembles of  $N_{\text{rep}}$  parton distribution functions and observables  $\mathcal{O}(\text{PDF})$  are conventionally calculated from the average of the predictions obtained from the ensemble  $\langle \mathcal{O}(\text{PDF}) \rangle = \frac{1}{N_{\text{rep}}} \sum_{k=1}^{N_{\text{rep}}} \mathcal{O}(\text{PDF}_k)$ . In the case of PDF uncertainties provided by standard Hessian eigenvector error sets, this can be achieved by creating the  $k$ -th random replica by introducing random fluctuations around the central PDF set.

As a next step, the initial PDF probability distributions are updated by applying weights  $w_k$ , calculated as:

$$w_k = \frac{(\chi_k^2)^{\frac{1}{2}(N_{\text{data}}-1)} e^{-\frac{1}{2}\chi_k^2}}{\frac{1}{N_{\text{rep}}} \sum_{k=1}^{N_{\text{rep}}} (\chi_k^2)^{\frac{1}{2}(N_{\text{data}}-1)} e^{-\frac{1}{2}\chi_k^2}}, \quad (16)$$

where  $N_{\text{data}}$  is the number of new data points,  $k$  denotes the specific replica for which the weight is calculated and  $\chi_k^2$  is a difference between a given data point  $y_i$  and its theoretical prediction obtained with the  $k$ -th PDF replica:

$$\chi^2(y, \text{PDF}_k) = \sum_{i,j=1}^{N_{\text{data}}} (y_i - y_i(\text{PDF}_k)) \sigma_{ij}^{-1} (y_j - y_j(\text{PDF}_k)) \quad (17)$$

The new, reweighted PDFs commonly are chosen to be based upon a smaller number of PDF sets compared to the input because replicas that are incompatible with the data are discarded in order to create a more stream-lined PDF set.

## 6 Alternatives to DGLAP formalism

Different approaches that are alternatives to the DGLAP formalism can be used to analyse DIS data in HERAFitter. These include several different dipole models and the use of transverse momentum dependent, or unintegrated PDFs, uPDFs. These approaches are discussed below.

### 6.1 DIPOLE models

The dipole picture provides an alternative approach to virtual photon-proton scattering at low  $x$  which allows the description of both inclusive and diffractive processes. In this approach, the virtual photon fluctuates into a  $q\bar{q}$  (or  $q\bar{q}g$ ) dipole which interacts with the proton [88]. The dipoles can be viewed as quasi-stable quantum mechanical states, which have very long life time  $\propto 1/m_p x$  and a size which is not changed by scattering. The dynamics of the interaction are embedded in the dipole scattering amplitude.

Several dipole models which assume different behavior of the dipole-proton cross sections are implemented in HERAFitter: the Golec-Biernat-Wüsthoff (GBW) dipole saturation model [21], the colour glass condensate approach to the high parton density regime called the Iancu-Itakura-Munier (IIM) dipole model [22] and a modified GBW model which takes into account the effects of DGLAP evolution called the Bartels-Golec-Kowalski (BGK) dipole model [23].

**GBW model:** In the GBW model the dipole-proton cross section  $\sigma_{\text{dip}}$  is given by

$$\sigma_{\text{dip}}(x, r^2) = \sigma_0 \left( 1 - \exp \left[ -\frac{r^2}{4R_0^2(x)} \right] \right), \quad (18)$$

where  $r$  corresponds to the transverse separation between the quark and the antiquark, and  $R_0^2$  is an  $x$ -dependent scale parameter which represents the spacing of the gluons in the proton.  $R_0^2(x) = (x/x_0)^\lambda$  is called the saturation radius. The fitted parameters are the cross-section normalisation  $\sigma_0$  and  $x_0$  and  $\lambda$ . This model gives exact Bjorken scaling when the dipole size  $r$  is small.

**IIM model:** The IIM model assumes an improved expression for the dipole cross section which is based on the Balitsky-Kovchegov equation [89]. The explicit formula for  $\sigma_{\text{dip}}$  can be found in [22]. The fitted parameters are an alternative scale parameter  $\tilde{R}$ ,  $x_0$  and  $\lambda$ .

**BGK model:** The BGK model modifies the GBW model by taking into account the DGLAP evolution of the gluon density. The dipole cross section is given by

$$\sigma_{\text{dip}}(x, r^2) = \sigma_0 \left( 1 - \exp \left[ -\frac{\pi^2 r^2 \alpha_s(\mu^2) x g(x, \mu^2)}{3\sigma_0} \right] \right). \quad (19)$$

The factorisation scale  $\mu^2$  has the form  $\mu^2 = C_{bgk}/r^2 + \mu_0^2$ . This model relates to the GBW model using the idea

that the spacing  $R_0$  is inverse to the gluon density. The gluon density parametrized at some starting scale  $Q_0^2$  by Eq. 6 is evolved to larger scales using DGLAP evolution. The fitted parameters for this model are  $\sigma_0$ ,  $\mu_0^2$  and three parameters for the gluon density:  $A_g$ ,  $\lambda_g$ ,  $C_g$ . The parameter  $C_{bgk}$  is fixed:  $C_{bgk} = 4.0$ .

### BGK model with valence quarks:

The dipole models are valid in the low- $x$  region only, where the valence quark contribution is small. The new HERA  $F_2$  data have a precision which is better than 2%. Therefore, in HERAFitter the contribution of the valence quarks can be taken from the PDF fits and added to the original BGK model [90, 91], this is uniquely possible within the HERAFitter framework.

### 6.2 Transverse Momentum Dependent PDFs with CCFM

QCD calculations of multiple-scale processes and complex final-states require in general transverse-momentum dependent (TMD) [92], or unintegrated, parton density and parton decay functions [93–101]. TMD factorisation has been proven recently [92] for inclusive DIS. For special processes in hadron-hadron scattering, like heavy flavor or vector boson (including Higgs) production, TMD factorisation has also been proven in the high-energy limit (small  $x$ ) [102–104]

In the framework of high-energy factorisation [102, 105, 106] the DIS cross section can be written as a convolution in both longitudinal and transverse momenta of the TMD parton density function  $\mathcal{A}(x, k_t, \mu)$  with off-shell partonic matrix elements, as follows

$$\sigma_j(x, Q^2) = \int_x^1 dz \int d^2 k_t \hat{\sigma}_j(x, Q^2, z, k_t) \mathcal{A}(z, k_t, \mu) \quad (20)$$

with the DIS cross sections  $\sigma_j$ , ( $j = 2, L$ ) related to the structure functions  $F_2$  and  $F_L$ . The hard-scattering kernels  $\hat{\sigma}_j$  of Eq. (20), are  $k_t$ -dependent and the evolution of the transverse momentum dependent gluon density  $\mathcal{A}$  is obtained by combining the resummation of small- $x$  logarithmic contributions [107–109] with medium- $x$  and large- $x$  contributions to parton splitting [11, 14, 15] according to the CCFM evolution equation [19, 110, 111].

The factorisation formula (20) allows resummation of logarithmically enhanced  $x \rightarrow 0$  contributions to all orders in perturbation theory, both in the hard scattering coefficients and in the parton evolution, taking fully into account the dependence on the factorisation scale  $\mu$  and on the factorisation scheme [112, 113].

The cross section  $\sigma_j$ , ( $j = 2, L$ ) is calculated in a FFN scheme, where only the boson-gluon fusion process ( $\gamma^* g^* \rightarrow q\bar{q}$ ) is included. The masses of the quarks are explicitly included with the light and heavy quark masses being free parameters. In addition to  $\gamma^* g^* \rightarrow q\bar{q}$ , the contribution from



valence quarks is included via  $\gamma^* q \rightarrow q$  as described later by using a CCFM evolution of valence quarks [114, 115].

### CCFM Grid Techniques:

The CCFM evolution cannot easily be written in an analytic closed form. For this reason a Monte Carlo method is employed, which is however time-consuming, and cannot be used in a straightforward manner in a fit program. Following the convolution method introduced in [115, 116], the kernel  $\tilde{\mathcal{A}}(x'', k_t, p)$  is determined from the Monte Carlo solution of the CCFM evolution equation, and then folded with the non-perturbative starting distribution  $\mathcal{A}_0(x)$ :

$$\begin{aligned} x\mathcal{A}(x, k_t, p) &= x \int dx' \int dx'' \mathcal{A}_0(x') \tilde{\mathcal{A}}(x'', k_t, p) \delta(x'x'' - x) \\ &= \int dx' \mathcal{A}_0(x') \cdot \frac{x}{x'} \tilde{\mathcal{A}}\left(\frac{x}{x'}, k_t, p\right) \end{aligned} \quad (21)$$

with  $k_t$  being the transverse momentum of the propagator gluon and  $p$  being the evolution variable. The kernel  $\tilde{\mathcal{A}}$  incorporates all of the dynamics of the evolution. It is determined on a grid of  $50 \otimes 50 \otimes 50$  bins in  $x, k_t, p$ . The binning in the grid is logarithmic, except for the longitudinal variable  $x$  where 40 bins in logarithmic spacing below 0.1, and 10 bins in linear spacing above 0.1 are used.

The calculation of the cross section according to Eq. (20) involves a multidimensional Monte Carlo integration which is time consuming and suffers from numerical fluctuations. This cannot be employed directly in a fit procedure involving the calculation of numerical derivatives in the search for the minimum. Instead the following procedure is applied:

$$\begin{aligned} \sigma(x, Q^2) &= \int_x^1 dx_g \mathcal{A}(x_g, k_t, p) \hat{\sigma}(x, x_g, Q^2) \\ &= \int_x^1 dx' \mathcal{A}_0(x') \cdot \tilde{\sigma}(x/x', Q^2) \end{aligned} \quad (22)$$

Here, first  $\tilde{\sigma}(x', Q^2)$  is calculated numerically with a Monte Carlo integration on a grid in  $x$  for the values of  $Q^2$  used in the fit. Then the last step in Eq. (22) is performed with a fast numerical gauss integration, which can be used in standard fit procedures.

### Functional Forms for TMD parameterisation:

For the starting distribution  $\mathcal{A}_0$ , at the starting scale  $Q_0$ , the following form is used:

$$x\mathcal{A}_0(x, k_t) = Nx^{-B} \cdot (1-x)^C (1-Dx + E\sqrt{x}) \exp[-k_t^2/\sigma^2], \quad (23)$$

with  $\sigma^2 = Q_0^2/2$  and the free parameters  $N, B, C, D, E$ . Valence quarks are treated using the method of [114] as described in [115] with a starting distribution taken from any collinear PDF. At every scale  $p$  the flavor sum rule is fulfilled.

### 6.3 Diffractive PDFs

Similarly to standard DIS, diffractive parton distributions (DPDFs) can be derived from QCD fits to diffractive cross sections. At HERA about 10% of deep inelastic interactions are diffractive leading to events in which the interacting proton stays intact ( $ep \rightarrow eXp$ ). In the diffractive process the proton appears well separated from the rest of the hadronic final state by a large rapidity gap and this is interpreted as the diffractive dissociation of the exchanged virtual photon to produce a hadronic system  $X$  with mass much smaller than  $W$  and the same net quantum numbers as the exchanged photon. For such processes, the proton vertex factorisation approach is assumed where diffractive DIS is mediated by the exchange of a hard Pomeron or a secondary Reggeon. The factorisable pomeron picture has proved remarkably successful in the description of most of these data.

In addition to the usual variables  $x, Q^2$ , one must consider the squared four-momentum transfer  $t$  (the undetected momentum transfer to the proton system) and the mass  $M_X$  of the diffractively produced final state. In practice, the variable  $M_X$  is often replaced by  $\beta = \frac{Q^2}{M_X^2 + Q^2 - t}$ . In models based on a factorisable Pomeron,  $\beta$  may be viewed as the fraction of the pomeron longitudinal momentum which is carried by the struck parton,  $x = \beta x_{IP}$ .

For the inclusive case, the diffractive cross-section can be expressed as:

$$\frac{d\sigma}{d\beta dQ^2 dx_{IP} dt} = \frac{2\pi\alpha^2}{\beta Q^4} (1 + (1-y)^2) \bar{\sigma}^{D(4)}(\beta, Q^2, x_{IP}, t) \quad (24)$$

where the “reduced cross-section”,  $\bar{\sigma}$ , is defined as

$$\bar{\sigma}^{D(4)} = F_2^{D(4)} - \frac{y^2}{1+(1-y)^2} F_L^{D(4)} = F_T^{D(4)} + \frac{2(1-y)}{1+(1-y)^2} F_L^{D(4)}. \quad (25)$$

With  $x = x_{IP}\beta$  we can relate this to the standard DIS formula. The diffractive structure functions can be expressed as convolutions of the calculable coefficient functions with diffractive quark and gluon distribution functions, which in general depend on all of  $x_{IP}, Q^2, \beta, t$ .

The diffractive PDFs in HERAFitter are implemented following the prescription of ZEUS publication [117] and can be used to reproduce the main results.

### 7 Application of HERAFitter

The HERAFitter project has successfully introduced into a wide variety of tools to facilitate investigations of the HEP experimental data and theoretical calculations. It provides a versatile interface for understanding and interpreting new data and the derived PDFs. The HERAFitter platform not only allows the extraction of PDFs but also of theory parameters such as the strong coupling and heavy quark masses.

The parameters and distributions are output with a quantitative assessment of the fit quality with fully detailed information on experimental and theoretical uncertainties. The results are also output to PDF LHAPDF grids that can be used to study predictions for SM or beyond SM processes, as well as for the study of the impact of future collider measurements (using pseudo-data).

So far the HERAFitter platform has been used to produce grids from the QCD analyses performed at HERA ([29]), and at the LHC, using measurements from ATLAS [30, 31] (the first ever ATLAS PDF sets [118]).

New results that have been based on the HERAFitter platform include the following SM processes studied at the LHC: inclusive Drell-Yan and W and Z production [30, 32, 33]; inclusive jets [31, 34] production. At HERA, the results of QCD analyses using HERAFitter are published for inclusive H1 measurements [35] and the recent combination of charm production measurements in DIS [36]. The HERAFitter framework also provides a unique possibility to make impact studies for future colliders as illustrated by the QCD studies that have been performed to explore the potential of the LHeC data [119].

A determination of the transverse momentum dependent gluon density using precision HERA data obtained with HERAFitter has been reported in [120].

In addition, a recent study based on a set of parton distribution functions determined with the HERAFitter program using HERA data was performed [121]. It addresses the issue of correlations between uncertainties for the LO, NLO and NNLO sets. These sets are then propagated to study uncertainties for ratios of cross sections calculated at different orders in QCD and a reduction of overall theoretical uncertainty is observed.

## 8 Summary

The HERAFitter project is a unique platform for QCD analyses to study the structure of the proton. It incorporates not only the crucial data on Deep Inelastic Scattering from HERA but also data from the hadron colliders which are sensitive to Parton Distribution Functions. A variety of up-to-date theory calculations are available for each process at LO, NLO and NNLO when possible. HERAFitter has flexible modular structure and contains many different useful tools for PDF interpretation. HERAFitter is the first open source platform which is optimal for benchmarking studies.

**Acknowledgements** HERAFitter developers team acknowledges the kind hospitality of DESY and funding by the Helmholtz Alliance "Physics at the Terascale" of the Helmholtz Association. We are grateful to the DESY IT department for their support of the HERAFitter developers. Additional support was received from BMBF-JINR cooperation program, Heisenberg-Landau program and RFBR grant 12-02-91526-

CERN a. We also acknowledge Nathan Hartland with Luigi Del Debbio for contributing to the implementation of the Bayesian Reweighting technique and would like to thank R. Thorne for fruitful discussions.

## References

1. G. Aad *et al.* [ATLAS Collaboration], Phys.Lett. **B716**, 1 (2012), [[1207.7214](#)].
2. S. Chatrchyan *et al.* [CMS Collaboration], Phys.Lett. **B716**, 30 (2012), [[1207.7235](#)].
3. J. C. Collins *et al.* (1989), *Factorization of Hard Processes (in QCD in Perturbative Quantum Chromodynamics)*, ISBN: 9971-50-564-9, 9971-50-565-7.
4. E. Perez and E. Rizvi, Rep.Prog.Phys. **76**, 046201 (2013), [[1208.1178](#)].
5. S. Forte and G. Watt, Ann.Rev.Nucl.Part.Sci. **63**, 291 (2013), [[1301.6754](#)].
6. A. Martin, W. Stirling, R. Thorne, and G. Watt, Eur. Phys. J. C **63**, 189 (2009), [[arXiv:0901.0002](#)], URL <http://mstwpdf.hepforge.org/>.
7. J. Gao, M. Guzzi, J. Huston, H.-L. Lai, Z. Li, *et al.*, Phys.Rev. **D89**, 033009 (2014), [[1302.6246](#)], URL <http://hep.pa.msu.edu/cteq/public/>.
8. R. D. Ball, V. Bertone, S. Carrazza, C. S. Deans, L. Del Debbio, *et al.*, Nucl.Phys. **B867**, 244 (2013), [[1207.1303](#)], URL <https://nnpdf.hepforge.org/>.
9. S. Alekhin, J. Bluemlein, and S. Moch (2013), [[1310.3059](#)].
10. P. Jimenez-Delgado and E. Reya, Phys.Rev. **D80**, 114011 (2009), [[0909.1711](#)], URL <http://www.het.physik.tu-dortmund.de/pdfserver/index.html>.
11. V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**, 438 (1972).
12. V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**, 675 (1972).
13. L. N. Lipatov, Sov. J. Nucl. Phys. **20**, 94 (1975).
14. Y. L. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977).
15. G. Altarelli and G. Parisi, Nucl. Phys. B **126**, 298 (1977).
16. M. Botje (2010), <http://www.nikef.nl/h24/qcdnum/index.html>, [[arXiv:1005.1481](#)].
17. M. Ciafaloni, Nucl. Phys. B **296**, 49 (1988).
18. S. Catani, F. Fiorani, and G. Marchesini, Phys. Lett. B **234**, 339 (1990).
19. S. Catani, F. Fiorani, and G. Marchesini, Nucl. Phys. B **336**, 18 (1990).
20. G. Marchesini, Nucl. Phys. B **445**, 49 (1995).
21. K. Golec-Biernat and M. Wüsthoff, Phys. Rev. D **59**, 014017 (1999), [[hep-ph/9807513](#)].
22. E. Iancu, K. Itakura, and S. Munier, Phys. Lett. **B590**, 199 (2004), [[hep-ph/0310338](#)].

23. J. Bartels, K. Golec-Biernat, and H. Kowalski, Phys. Rev. D **66**, 014001 (2002), [[hep-ph/0203258](#)].
24. F. James and M. Roos, Comput. Phys. Commun. **10**, 343 (1975).
25. M. Dittmar, S. Forte, A. Glazov, and S. Moch (2009), Altarelli, G. and others (contributing authors), [[arXiv:0901.2504](#)].
26. M. R. Whalley, D. Bourilkov, and R. Group (2005), [[hep-ph/0508110](#)].
27. *LHAPDF*, URL <http://lhapdf.hepforge.org>.
28. *TMD Collaboration*, to be published.
29. F. Aaron *et al.* [H1 and ZEUS Collaborations], JHEP **1001**, 109 (2010), [[arXiv:0911.0884](#)].
30. G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. Lett. **109**, 012001 (2012), [[arXiv:1203.4051](#)].
31. G. Aad *et al.* [ATLAS Collaboration], Eur.Phys.J. **73**, 2509 (2013), [[arXiv:1304.4739](#)].
32. G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. **B725**, 223 (2013), [[arXiv:1305.4192](#)].
33. S. Chatrchyan *et al.* [CMS Collaboration], submitted to Phys. Rev. D (2014), [[arXiv:1312.6283](#)].
34. S. Chatrchyan *et al.* [CMS Collaboration], CMS PAS **SMP-12-028** (2014).
35. F. Aaron *et al.* [H1 Collaboration], JHEP **1209**, 061 (2012), [[arXiv:1206.7007](#)].
36. H. Abramowicz *et al.* [H1 and ZEUS Collaborations], Eur. Phys. J. **C73**, 2311 (2013), [[arXiv:1211.1182](#)].
37. R. Devenish and A. Cooper-Sarkar (2011), *Deep Inelastic Scattering*, ISBN: 0199602255, 9780199602254.
38. J. C. Collins and W.-K. Tung, Nucl. Phys. B **278**, 934 (1986).
39. E. Laenen *et al.*, Phys. Lett. **B291**, 325 (1992).
40. E. Laenen *et al.*, Nucl. Phys. **B392**, 162, 229 (1993).
41. S. Riemersma, J. Smith, and van Neerven. W.L., Phys. Lett. **B347**, 143 (1995), [[hep-ph/9411431](#)].
42. S. Alekhin, *OPENQCDRAD*, a program description and the code are available via: <http://www-zeuthen.desy.de/~alekhin/OPENQCDRAD>.
43. H. Kawamura, N. Lo Presti, S. Moch, and A. Vogt, Nucl.Phys. **B864**, 399 (2012).
44. S. Alekhin and S. Moch, Phys. Lett. **B699**, 345 (2011), [[arXiv:1011.5790](#)].
45. R. Demina, S. Keller, M. Kramer, S. Kretzer, R. Martin, *et al.* (1999), [[hep-ph/0005112](#)].
46. R. S. Thorne and R. G. Roberts, Phys. Rev. D **57**, 6871 (1998), [[hep-ph/9709442](#)].
47. R. S. Thorne, Phys. Rev. **D73**, 054019 (2006), [[hep-ph/0601245](#)].
48. R. S. Thorne, Phys. Rev. D **86**, 074017 (2012), [[arXiv:1201.6180](#)].
49. J. C. Collins, Phys.Rev. **D58**, 094002 (1998), [[hep-ph/9806259](#)].
50. H. Spiesberger, Private communication.
51. F. Jegerlehner, Proceedings, LC10 Workshop **DESY 11-117** (2011).
52. H. Burkhard, F. Jegerlehner, G. Penso, and C. Verzegnassi, in CERN Yellow Report on "Polarization at LEP" 1988.
53. S. D. Drell and T.-M. Yan, Phys. Rev. Lett. **25**, 316 (1970).
54. M. Yamada and M. Hayashi, Nuovo Cim. **A70**, 273 (1982).
55. J. M. Campbell and R. K. Ellis, Phys. Rev. **D60**, 113006 (1999), [[arXiv:9905386](#)].
56. J. M. Campbell and R. K. Ellis, Phys. Rev. **D62**, 114012 (2000), [[arXiv:0006304](#)].
57. J. M. Campbell and R. K. Ellis, Nucl. Phys. Proc. Suppl. **205-206**, 10 (2010), [[arXiv:1007.3492](#)].
58. Y. Li and F. Petriello, Phys.Rev. **D86**, 094034 (2012), [[arXiv:1208.5967](#)].
59. G. Bozzi, J. Rojo, and A. Vicini, Phys.Rev. **D83**, 113008 (2011), [[arXiv:1104.2056](#)].
60. A. Gehrmann-De Ridder, T. Gehrmann, E. Glover, and J. Pires, Phys. Rev. Lett. **110**, 162003 (2013), [[arXiv:1301.7310](#)].
61. E. Glover and J. Pires, JHEP **1006**, 096 (2010), [[arXiv:1003.2824](#)].
62. J. Currie, A. Gehrmann-De Ridder, E. Glover, and J. Pires, JHEP **1401**, 110 (2014), [[1310.3993](#)].
63. Z. Nagy and Z. Trocsanyi, Phys.Rev. **D59**, 014020 (1999), [[hep-ph/9806317](#)].
64. Z. Nagy, Phys.Rev.Lett. **88**, 122003 (2002), [[hep-ph/0110315](#)].
65. M. Czakon, P. Fiedler, and A. Mitov, Phys. Rev. Lett. **110**, 252004 (2013), [[1303.6254](#)].
66. M. Aliev, H. Lacker, U. Langenfeld, S. Moch, P. Uwer, *et al.*, Comput.Phys.Commun. **182**, 1034 (2011), [[arXiv:1007.1327](#)].
67. J. M. Campbell, R. Frederix, F. Maltoni, and F. Tramontano, Phys.Rev.Lett. **102**, 182003 (2009), [[0903.0005](#)].
68. J. M. Campbell and F. Tramontano, Nucl.Phys. **B726**, 109 (2005), [[hep-ph/0506289](#)].
69. J. M. Campbell, R. K. Ellis, and F. Tramontano, Phys.Rev. **D70**, 094012 (2004), [[hep-ph/0408158](#)].
70. J. M. Campbell and R. K. Ellis (2012), report FERMILAB-PUB-12-078-T, [[1204.1513](#)].
71. T. Kluge, K. Rabbertz, and M. Wobisch, pp. 483–486 (2006), [[hep-ph/0609285](#)].
72. T. Carli *et al.*, Eur. Phys. J. **C66**, 503 (2010), [[arXiv:0911.2985](#)].
73. Z. Nagy and Z. Trocsanyi, Phys.Rev.Lett. **87**, 082001 (2001), [[hep-ph/0104315](#)].
74. Z. Nagy, Phys.Rev. **D68**, 094002 (2003), [[hep-ph/0307268](#)].

- 1087 75. M. Wobisch, D. Britzger, T. Kluge, K. Rab- 1140  
1088 bertz, and F. Stober [fastNLO Collaboration] (2011), 1141  
1089 [arXiv:1109.1310]. 1142
- 1090 76. N. Kidonakis and J. Owens, Phys.Rev. **D63**, 054019 1143  
1091 (2001), [hep-ph/0007268]. 1144
- 1092 77. D. Britzger, K. Rabbertz, F. Stober, and M. Wobisch 1145  
1093 [fastNLO Collaboration] (2012), [arXiv:1208.3641]. 1146
- 1094 78. <http://fastnlo.hepforge.org>, URL [http://fastnlo.](http://fastnlo.hepforge.org) 1147  
1095 [hepforge.org](http://fastnlo.hepforge.org). 1148
- 1096 79. <http://applgrid.hepforge.org>, URL [http:](http://applgrid.hepforge.org) 1149  
1097 [//applgrid.hepforge.org](http://applgrid.hepforge.org). 1150
- 1098 80. A. Glazov, S. Moch, and V. Radescu, Phys. Lett. B 1151  
1099 **695**, 238 (2011), [arXiv:1009.6170]. 1152
- 1100 81. J. Pumplin, D. Stump, R. Brock, D. Casey, J. Hus- 1153  
1101 ton, *et al.*, Phys.Rev. **D65**, 014013 (2001), [hep- 1154  
1102 ph/0101032]. 1155
- 1103 82. M. Botje, J.Phys. **G28**, 779 (2002), [hep-ph/0110123]. 1156
- 1104 83. W. T. Giele and S. Keller, Phys.Rev. **D58**, 094023 1157  
1105 (1998), [hep-ph/9803393]. 1158
- 1106 84. W. T. Giele, S. Keller, and D. Kosower (2001), [hep- 1159  
1107 ph/0104052]. 1160
- 1108 85. G. Watt and R. Thorne, JHEP **1208**, 052 (2012), 1161  
1109 [arXiv:1205.4024]. 1162
- 1110 86. R. D. Ball, V. Bertone, F. Cerutti, L. Del Deb- 1163  
1111 bio, S. Forte, *et al.*, Nucl.Phys. **B855**, 608 (2012), 1164  
1112 [arXiv:1108.1758]. 1165
- 1113 87. R. D. Ball *et al.* [NNPDF Collaboration], Nucl.Phys. 1166  
1114 **B849**, 112 (2011), [arXiv:1012.0836]. 1167
- 1115 88. N. N. Nikolaev and B. Zakharov, Z.Phys. **C49**, 607 1168  
1116 (1991). 1169
- 1117 89. I. Balitsky, Nucl. Phys. B **463**, 99 (1996), [hep- 1170  
1118 ph/9509348]. 1171
- 1119 90. P. Belov, Doctoral thesis, Universität Hamburg (2013), 1172  
1120 [DESY-THESIS-2013-017]. 1173
- 1121 91. A. Luszczak and H. Kowalski (2013), [1312.4060]. 1174
- 1122 92. J. Collins, *Foundations of perturbative QCD*, vol. 32 1175  
1123 (Cambridge monographs on particle physics, nuclear 1176  
1124 physics and cosmology., 2011). 1177
- 1125 93. S. M. Aybat and T. C. Rogers, Phys.Rev. **D83**, 114042 1178  
1126 (2011), [1101.5057]. 1179
- 1127 94. M. Buffing, P. Mulders, and A. Mukherjee, 1180  
1128 Int.J.Mod.Phys.Conf.Ser. **25**, 1460003 (2014),  
1129 [1309.2472].
- 1130 95. M. Buffing, A. Mukherjee, and P. Mulders, Phys.Rev. 1181  
1131 **D88**, 054027 (2013), [1306.5897].
- 1132 96. M. Buffing, A. Mukherjee, and P. Mulders, Phys.Rev. 1182  
1133 **D86**, 074030 (2012), [1207.3221].
- 1134 97. P. Mulders, Pramana **72**, 83 (2009), [0806.1134].
- 1135 98. S. Jadach and M. Skrzypek, Acta Phys.Polon. **B40**,  
1136 2071 (2009), [0905.1399].
- 1137 99. F. Hautmann, Acta Phys.Polon. **B40**, 2139 (2009).
- 1138 100. F. Hautmann, M. Hentschinski, and H. Jung (2012),  
1139 [1205.6358].
101. F. Hautmann and H. Jung, Nucl.Phys.Proc.Suppl. **184**,  
64 (2008), [0712.0568].
102. S. Catani, M. Ciafaloni, and F. Hautmann, Phys. Lett.  
B **242**, 97 (1990).
103. J. C. Collins and R. K. Ellis, Nucl. Phys. B **360**, 3  
(1991).
104. F. Hautmann, H. Jung, and V. Pandis, AIP Conf.Proc.  
**1350**, 263 (2011), [1011.6157].
105. S. Catani, M. Ciafaloni, and F. Hautmann, Nucl. Phys.  
B **366**, 135 (1991).
106. S. Catani, M. Ciafaloni, and F. Hautmann, Phys. Lett.  
B **307**, 147 (1993).
107. L. Lipatov, Phys.Rept. **286**, 131 (1997), [hep-  
ph/9610276].
108. V. S. Fadin, E. Kuraev, and L. Lipatov, Phys.Lett. **B60**,  
50 (1975).
109. I. I. Balitsky and L. N. Lipatov, Sov. J. Nucl. Phys. **28**,  
822 (1978).
110. M. Ciafaloni, Nucl. Phys. **B296**, 49 (1988).
111. G. Marchesini, Nucl. Phys. B **445**, 49 (1995), [hep-  
ph/9412327].
112. S. Catani and F. Hautmann, Nucl. Phys. B **427**, 475  
(1994), [hep-ph/9405388].
113. S. Catani and F. Hautmann, Phys.Lett. **B315**, 157  
(1993).
114. M. Deak, F. Hautmann, H. Jung, and K. Kutak,  
*Forward-Central Jet Correlations at the Large Hadron  
Collider* (2010), [arXiv:1012.6037].
115. F. Hautmann and H. Jung, Nuclear Physics B **883**, 1  
(2014), [1312.7875].
116. H. Jung and F. Hautmann (2012), [arXiv:1206.1796].
117. S. Chekanov *et al.* [ZEUS Collaboration], Nucl. Phys.  
**B831**, 1 (2010), [hep-ex/09114119].
118. *ATLAS NNLO epWZ12*, available via:  
<http://lhapdf.hepforge.org/pdfsets>.
119. J. L. Abelleira Fernandez *et al.* [LHeC Study  
Group], Journal of Phys. **G**, 075001 (2012),  
[arXiv:1206.2913].
120. F. Hautmann and H. Jung (2013), [1312.7875].
121. HERAFitter Developers Team and M. Lisovyi (2014),  
[arXiv:1404.4234].