

Lenses

<https://github.com/heptagons/lenses>

2024/1/4

Abstract

Lenses are equilateral hexagons resembling concave and convex optical lenses. Lenses consecutive six internal angles are $(\theta_1, \theta_2, \theta_3, \theta_1, \theta_2, \theta_3)$ where $\theta_1 = x\theta_0$, $\theta_2 = y\theta_0$, and $\theta_3 = z\theta_0$ where $\theta_0 = 2\pi/s$ is the base angle of symmetry $s = x + y + z$. Lenses can be formed adding and subtracting rhombi or by intersecting equi-stars with others.

1 Equi-stars




Figure 1: Equi-stars of symmetries 3,5,7 and 9.

Equi-stars are equilateral polygons with an even number of sides and vertices of at most two different angles. These stars can be defined with only two numbers: A symmetry integer s and a minimum angle integer a so the star is defined as $S(s, a)$. Here we are interested only in symmetries of the form $s = 2n + 1$ for $n = 1, 2, 3, \dots$. Every symmetry $s = 2n + 1$ has exactly n different stars: $S(s, 1), S(s, 2), \dots, S(s, n)$. Stars of the form $S(s, n)$ correspond to the regular polygons of $2s$ sides.

Figure 1 show the stars for the smaller symmetries in translucent colors and intersecting with others of the same symmetry. At (i) we have for the symmetry 3 the only star in red $S(3, 1)$ which is the regular hexagon. At (ii) we have for the symmetry 5 the regular decagon in red $S(5, 2)$ and the star $S(5, 1)$ in yellow; the region in orange is the intersection of the two stars. At (iii) for symmetry 7 we have three stars: The equilateral 14-gon $S(7, 3)$ in red, the $S(7, 2)$ in yellow and the $S(7, 1)$ in green. At (iv) we have for the symmetry 9 four stars: The regular 18-gon $S(9, 4)$ in red, the $S(9, 3)$ in yellow, the $S(9, 2)$ in green and the $S(9, 1)$ in blue.

2 Symmetry 5

Symmetry 5 is based in angle $\beta = \frac{2\pi}{5}$ and produces the two rhombi (b, c) and the two lenses (B, C) .

2.1 Rhombi (b, c)




Figure 2: Rhombi (b, c) from dissecting stars S_{10} .

Figure 2 show the rhombi (b, c) . Inspecting the stars we get the areas simply adding their rhombi. At (i) the star $S_{10}(1, 8)$ with area $A = 10b$. At (ii) the star $S_{10}(2, 6) = S_5(1, 3)$ with area $A = 5c$. At (iii) the regular decagon equivalent to stars $S_{10}(4, 4) = S_5(2, 2)$ with area $A = 5b + 5c$. Table 1 show the rhombi (b, c) internal angles in terms of angle $\beta = 2\pi/5$ and areas for side equals to 1. Dividing areas we find $\frac{c}{b} = 2 \cos\left(\frac{\pi}{5}\right) = \frac{\sqrt{5} + 1}{2}$.

Rhombus	θ_1	θ_2	Area
b	$\beta/2$	$4\beta/2$	$\sin(2\beta) = \frac{\sqrt{10 - 2\sqrt{5}}}{4}$
c	$2\beta/2$	$3\beta/2$	$\sin(\beta) = \frac{\sqrt{10 + 2\sqrt{5}}}{4} = b \cos\left(\frac{\pi}{5}\right) = b \left(\frac{\sqrt{5} + 1}{2}\right)$

Table 1: Rhombi (b, c) internal angles and areas. $\theta_1 + \theta_2 = \pi$ and $\beta = 2\pi/5$.

2.2 Regular pentagon and star $|5/2|$




Figure 3: Regular pentagon $|5/1|$ at (i). Star $|5/2|$ at (ii). Double pentagon at (iii).

Figure 3 show regular pentagon and isotoxal star $|5/2|$ dissected with rhombi (\mathbf{b}, \mathbf{c}) plus concave pentagons (in gray). Let \mathbf{x} be the area of such gray piece. By inspection the area of regular pentagon at (i) is $A_1 = \mathbf{c} + \mathbf{x}$ and the area of regular pentagon at (iii) is $P_2 = \mathbf{b} + 5\mathbf{c} + 2\mathbf{x}$. Since the side of A_2 is the double of A_1 its area is four times so we can get the value of \mathbf{x} in terms of (\mathbf{b}, \mathbf{c})

$$\begin{aligned} 4P_1 &= P_4 \\ 4(\mathbf{c} + \mathbf{x}) &= \mathbf{b} + 5\mathbf{c} + 2\mathbf{x} \\ \mathbf{x} &= \frac{\mathbf{b} + \mathbf{c}}{2} \end{aligned} \tag{1}$$

We use the value of \mathbf{x} to get the areas of pentagon (i) and star (ii):

$$\begin{aligned} A|5/1| &= \mathbf{c} + \mathbf{x} \\ &= \frac{\mathbf{b} + 3\mathbf{c}}{2} \end{aligned} \tag{2}$$

$$\begin{aligned} A|5/2| &= \mathbf{b} + 2\mathbf{x} \\ &= 2\mathbf{b} + \mathbf{c} \end{aligned} \tag{3}$$

2.3 Lenses (\mathbf{B}, \mathbf{C})




Figure 4: Lenses (\mathbf{B}, \mathbf{C}) build with rhombi (\mathbf{b}, \mathbf{c}) .

Figure 4 show lenses (\mathbf{B}, \mathbf{C}) construction and two stars formed with them. At (i) we form the lense \mathbf{B} with perimeter $\overline{B_1 \dots B_6}$ adding two rhombi \mathbf{b} ($\overline{B_1 B_2 B_3 O}$ and $\overline{B_1 O B_5 B_6}$) and adding one rhombus \mathbf{c} ($\overline{O B_3 B_4 B_5}$) so its area is $2\mathbf{b} + \mathbf{c}$. Lense \mathbf{B} is equivalent to the hexagon $H_5(1, 2, 2)$. At (ii) we form the lense \mathbf{C} with perimeter $\overline{C_1 \dots C_6}$ adding two rhombi \mathbf{c} ($\overline{O C_1 C_2 C_3}$ and $\overline{O C_3 C_4 C_5}$) and subtracting one rhombus \mathbf{b} ($\overline{O C_1 C_6 C_5}$) so its area is $2\mathbf{c} - \mathbf{b}$. Lense \mathbf{C} is equivalent to the hexagon $H_5(1, 1, 3)$. Table 2 show the lenses (\mathbf{B}, \mathbf{C}) internal angles and areas.

Lense	θ_1	θ_2	θ_3	Area
\mathbf{B}	β	2β	2β	$2\mathbf{b} + \mathbf{c}$
\mathbf{C}	β	β	3β	$-\mathbf{b} + 2\mathbf{c}$

Table 2: Lenses (\mathbf{B}, \mathbf{C}) internal angles and areas in terms of rhombi (\mathbf{b}, \mathbf{c}) . $\theta_1 + \theta_2 + \theta_3 = 2\pi$ where $\beta = 2\pi/5$.




Figure 5: Two stars dissected with lenses (B , C).

Figure 5 show two stars dissected with lenses (B , C). At (i) the star $S_5(1,3)$ dissection implies its area is $A = B + 2C = 5c$. At (ii) the regular decagon or star $S_5(2,2)$ dissection implies its area is $A = 3B + C = 5(b + c)$.

3 Symmetry 7

Symmetry 7 is based in angle $\gamma = \frac{2\pi}{7}$ and produces the three rhombi (d, f, e) and the three lenses (D, E, F).

3.1 Rhombi (d, e, f)




Figure 6: Rhombi (d, e, f) from dissected stars S_{14} .

Figure 6 show rhombi (d, e, f). Inspecting the stars we get the areas simply adding their rhombi. At (i) the star $S_{14}(1,12)$ with area $A = 14d$. At (ii) the star $S_{14}(2,10) = S_7(1,5)$ with area $A = 7e$. At (iii) the star $S_{14}(4,8) = S_7(2,4)$ with area $A = 7(e + f)$. At (iv) the regular 14-gon equivalent to stars $S_{14}(6,6) = S_7(3,3)$ with area $A = 7(d + e + f)$. Table 3 show the symmetry 7 lenses internal angles based in angle $\gamma = 2\pi/7$ and the areas.

Rhombus	θ_1	θ_2	Area
d	$\gamma/2$	$6\gamma/2$	$\sin(3\gamma)$
e	$2\gamma/2$	$5\gamma/2$	$\sin(\gamma)$
f	$3\gamma/2$	$4\gamma/2$	$\sin(2\gamma)$

Table 3: Rhombi (d, e, f) internal angles. $\theta_1 + \theta_2 = \pi$ and $\gamma = 2\pi/7$.

3.2 Regular heptagon and stars $|7/3|$ and $|7/2|$




Figure 7: Heptagon $|7/1|$ at (i). Star $|7/3|$ at (ii). Star $|7/2|$ at (iii). Double heptagon at (iv).

Figure 7 show regular heptagon and heptagrams dissected with rhombi (c, d, e) plus equilateral concave heptagons (in gray). Let y be the area of such gray piece. By inspection the area of regular heptagon at (i) is $A_1 = e + 2f + y$ while the area of regular heptagon at (iv) is $A_2 = d + 7(e + f) + 2y$. Since the side of A_2 is the double of A_1 its area is four times so we can get the value of y in terms of (d, e, f):

$$\begin{aligned} 4A_1 &= A_2 \\ 4(e + 2f + y) &= d + 7(e + f) + 2y \\ y &= \frac{d + 3e - f}{2} \end{aligned} \quad (4)$$

We use the value of y to calculate the areas of heptagon (i) and stars (ii) and (iii) in terms of (d, e, f):

$$\begin{aligned} A|7/1| &= e + 2f + y \\ &= \frac{d + 5e + 3f}{2} \end{aligned} \quad (5)$$

$$\begin{aligned} A|7/3| &= d + 2y \\ &= 2d + 3e - f \end{aligned} \quad (6)$$

$$\begin{aligned} A|7/2| &= A\{7/3\} + 7f \\ &= 2d + 3e + 6f \end{aligned} \quad (7)$$

3.3 Lenses (D, E, F)




Figure 8: Lenses (D, E, F) build from rhombi (d, e, f).

Figure 8 show lenses (\mathbf{B}, \mathbf{C}) construction. At (i) we form the lense \mathbf{D} with perimeter $\overline{D_1 \dots D_6}$ adding two rhombi \mathbf{d} ($\overline{D_1 D_2 D_3 O}$ and $\overline{D_1 O D_5 D_6}$) and adding one rhombus \mathbf{e} ($\overline{O D_3 D_4 D_5}$) so its area is $2\mathbf{d} + \mathbf{e}$. Lense \mathbf{D} is equivalent to the hexagon $H_7(1, 3, 3)$.

At (ii) we form the lense \mathbf{E} with perimeter $\overline{E_1 \dots E_6}$ adding one rhombus \mathbf{e} ($\overline{E_1 E_2 O E_6}$) adding one rhombus \mathbf{f} ($\overline{O E_4 E_5 E_6}$) and subtracting one rhombus \mathbf{d} ($\overline{E_2 O E_4 E_3}$) so its area is $-\mathbf{d} + \mathbf{e} + \mathbf{f}$. Lense \mathbf{E} is equivalent to the hexagon $H_7(1, 2, 4)$.

At (iii) we form the lense \mathbf{F} with perimeter $\overline{F_1 \dots F_6}$ adding two rhombi \mathbf{f} ($\overline{F_1 F_2 F_3 O}$ and $\overline{F_3 F_4 F_5 O}$) and adding one rhombus \mathbf{d} ($\overline{F_1 O F_5 F_6}$) so its area is $\mathbf{d} + 2\mathbf{f}$. Lense \mathbf{F} is equivalent to the hexagon $H_7(2, 2, 3)$. Table 4 show the lenses $(\mathbf{D}, \mathbf{E}, \mathbf{F})$ internal angles and areas.

Lense	θ_1	θ_2	θ_3	Area
\mathbf{D}	γ	3γ	3γ	$2\mathbf{d} + \mathbf{e}$
\mathbf{E}	γ	2γ	4γ	$-\mathbf{d} + \mathbf{e} + \mathbf{f}$
\mathbf{F}	2γ	2γ	3γ	$-\mathbf{d} + 2\mathbf{f}$

Table 4: Lenses $(\mathbf{D}, \mathbf{E}, \mathbf{F})$ internal angles and areas in terms of rhombi $(\mathbf{d}, \mathbf{e}, \mathbf{f})$. $\theta_1 + \theta_2 + \theta_3 = 2\pi$ and $\gamma = 2\pi/7$.




Figure 9: Stars dissected with only lenses $(\mathbf{D}, \mathbf{E}, \mathbf{F})$.

Figure 9 show stars $S_7(2, 4)$ and $S_7(3, 3)$ dissected with lenses $(\mathbf{D}, \mathbf{E}, \mathbf{F})$. At (i) we have star $S_7(2, 4)$ and by inspection we deduce its area is $A = 2\mathbf{D} + 5\mathbf{E} + \mathbf{F}$. At (ii) we have regular 14-gon (or star $S_7(3, 3)$) and by inspection we deduce its area is $4\mathbf{D} + 3\mathbf{E} + 2\mathbf{F}$. Both stars have in common an area in green resembling a tree leaf. The star at (i) also contains two regions in yellow resembling crowns while the star at (ii) contains a region in cyan resembling a moon phase.




Figure 10: Stars dissected with octagons O_7 (in blue) and lenses (D, E, F) .

Figure 10 show Stars $S_7(1, 5)$, $S_7(2, 4)$ and $S_7(3, 3)$ dissected with octagons O_7 (in blue) and lenses (D, E, F) . At (i) we have the star $S_7(1, 5)$ and by inspection we deduce its area is $A = D + 2O_7$. At (ii) we have the star $S_7(2, 4)$ and we can conclude its area is $E + 3F + 2O_7$. Similarly the area of the 14-gon at (iii) is $3D + E + 3F + O_7$. Comparing the areas of the two 14-gons of figures 9 and 10 we can find the area of O_7 in terms of (E, F, G) :

$$\begin{aligned} 4D + 3E + 2F &= 3D + E + 3F + O_7 \\ O_7 &= D + 2E - F \end{aligned} \tag{8}$$

So we can calculate the area of star $S_7(1, 5)$ in terms of (E, F, G) :

$$\begin{aligned} S_7(1, 5) &= D + 2(D + 2E - F) \\ &= 3D + 4E - 2F \end{aligned} \tag{9}$$