

Horns unit

<https://github.com/heptagons/meccano/units/horns>

Abstract

Horns unit is a group of seven meccano ¹ strips intended to build polygons. We found the formula to calculate the internal angles then look for polygons and found hexagons, octagons and dodecagons. We found no pentagons.




Figure 1: The **horn unit** has seven strips: Two of length a , two of length $b + d$, two of length c and one of length e . We expect to build polygons with internal angle $C_1B_1B_2$ and perimeter including segments a, e, a .

1 Algebra

From figure 1 we start with triangle $\triangle A_1B_1C_1$. At vertex A_1 we have angle A and the supplement A' :

$$A \equiv \angle B_1A_1C_1 \tag{1}$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad \text{if and only if } a < b + c \tag{2}$$

$$A' \equiv \angle EA_1B_1 = \pi - A \tag{3}$$

$$\cos A' = \cos(\pi - A) = -\cos A = \frac{-b^2 - c^2 + a^2}{2bc} \tag{4}$$

We define $f \equiv b + d$ and $g \equiv \overline{EB_1}$. With the law of cosines we have:

$$f \equiv \boxed{b + d} \in \mathbb{N} \tag{5}$$

$$g^2 = c^2 + d^2 - 2cd \cos A' \tag{6}$$

$$\begin{aligned} &= c^2 + d^2 - (2cd) \frac{-b^2 - c^2 + a^2}{2bc} \\ &= \frac{bc^2 + bd^2 + b^2d + c^2d - a^2d}{b} \\ &= \frac{(b + d)(bd + c^2) - a^2d}{b} \end{aligned} \tag{7}$$

¹ Meccano mathematics by 't Hooft

Define a new variable $h = (b + d)(bd + c^2) - a^2d$:

$$h \equiv \boxed{(bd + c^2)f - a^2d} \in \mathbb{Z} \quad (8)$$

$$g^2 = \boxed{\frac{h}{b}} \quad \text{if and only if } 0 < h < b \quad (9)$$

We calculate angles $F \equiv \angle C_1B_1E$ and $G \equiv \angle B_2B_1E$. We replace g^2 by h/b :

$$\cos F = \frac{a^2 + g^2 - f^2}{2ag} = \frac{a^2b - bf^2 + h}{2abg} \quad (10)$$

$$\cos G = \boxed{\frac{e}{2g}} \quad (11)$$

Define new variable $j = a^2b - bf^2 + h$ so:

$$j \equiv \boxed{a^2b - bf^2 + h} \in \mathbb{Z} \quad (12)$$

$$\cos F = \frac{a^2b - bf^2 + h}{2abg} = \boxed{\frac{j}{2abg}} \quad (13)$$

We calculate cosines squares and products. Again we replace g^2 by h/b :

$$\cos F \cos G = \frac{ej}{4abg^2} = \frac{bej}{4abh} = \boxed{\frac{ej}{4ah}} \in \mathbb{Q} \quad (14)$$

$$\cos^2 F = \frac{j^2}{4a^2b^2g^2} = \frac{bj^2}{4a^2b^2h} = \boxed{\frac{j^2}{4a^2bh}} \in \mathbb{Q} \quad (15)$$

$$\cos^2 G = \frac{e^2}{4g^2} = \boxed{\frac{be^2}{4h}} \in \mathbb{Q} \quad (16)$$

$$\cos^2 F \cos^2 G = \frac{be^2j^2}{16a^2bh^2} = \boxed{\frac{e^2j^2}{16a^2h^2}} \in \mathbb{Q} \quad (17)$$

$$(18)$$

We calculate the sines part squared and set a common denominator as square $16a^2b^2h^2$:

$$(\sin F \sin G)^2 = (1 - \cos^2 F)(1 - \cos^2 G) \quad (19)$$

$$= 1 - \cos^2 F - \cos^2 G + \cos^2 F \cos^2 G$$

$$= 1 - \frac{j^2}{4a^2bh} - \frac{be^2}{4h} + \frac{e^2j^2}{16a^2h^2}$$

$$= 1 - \frac{j^2}{4a^2bh} - \frac{be^2}{4h} + \frac{e^2j^2}{16a^2h^2}$$

$$= \frac{16a^2b^2h^2 - (4bh)j^2 - (4a^2b^2h)be^2 + (b^2)e^2j^2}{16a^2b^2h^2}$$

$$= \frac{16a^2b^2h^2 - 4bhj^2 - 4a^2b^3e^2h + b^2e^2j^2}{16a^2b^2h^2}$$

$$= \frac{b(be^2 - 4h)(j^2 - 4a^2bh)}{16a^2b^2h^2} \quad (20)$$

Extract square root to get $\sin F \sin G = \sqrt{D}/A$ where $D, A \in \mathbb{Z}$:

$$\sin F \sin G = \boxed{\frac{\sqrt{b(be^2 - 4h)(j^2 - 4a^2bh)}}{4abh}} \in \mathbb{A} \quad (21)$$

We sum the angles F and G to get:

$$F + G \equiv \angle B_2 B_1 C_1 \quad (22)$$

$$\cos(F + G) = \cos F \cos G - \sin F \sin G \quad (23)$$

$$\begin{aligned} &= \frac{ej}{4ah} - \frac{\sqrt{b(be^2 - 4h)(j^2 - 4a^2bh)}}{4abh} \\ &= \frac{bej - \sqrt{b(be^2 - 4h)(j^2 - 4a^2bh)}}{4abh} \in \mathbb{A} \end{aligned} \quad (24)$$

2 Software

3 Examples

3.1 Hexagons examples

```

1 segments min=0 max=20 a,b,c,d,e:
2   1) 3,7,5,0,10
3   2) 3,8,7,0,13
4   3) 10,13,7,0,13
5   4) 8,13,7,0,14
6   5) 3,8,7,1,15
7   6) 5,8,7,2,15
8   7) 11,7,7,7,15
9   8) 7,13,8,0,16
10  9) 3,8,7,2,17
11 10) 5,8,7,3,17
12 11) 16,14,6,7,17
13 12) 8,7,5,7,18
14 13) 3,8,7,3,19
15 14) 5,8,7,4,19
16 15) 7,3,5,11,19
17 16) 7,8,5,6,19
18 --- PASS: TestHornsEHexagons (28.73s)

```




Figure 2: Hexagon size 10.




Figure 3: Hexagons size 13.




Figure 4: Horns() 60° .




Figure 5: Horns() 60° .

3.2 Octagons examples

```

1 segments min=1 max=40 a,b,c,d,e:
2   1) 2,3,3,3,8
3   2) 3,2,3,7,12
4   3) 14,9,9,9,16
5   4) 14,11,11,11,24
6   5) 21,21,14,6,24
7   6) 7,18,17,3,28
8   7) 9,10,11,17,36
9   8) 9,20,19,7,36
10  9) 10,9,11,21,40
11  10) 35,27,22,18,40
12 --- PASS: TestHornsEOctagons (566.86s)

```




Figure 6: Horns(2,3,3,3,8) 45° .




Figure 7: Horns(3,2,3,7,12) 45° .




Figure 8: Horns(14,9,9,9,16) 45° .




Figure 9: Octagons. Horns(14,11,11,11,24) and (21,21,14,6,24).

3.3 Dodecagons examples

```

1 segments min=1 max=40 a,b,c,d,e:
2   1) 6,5,5,5,8
3   2) 15,4,13,21,20
4   3) 15,9,12,16,20
5   4) 15,14,13,11,20
6   5) 15,18,15,7,20

```

```

7 | 6) 10,13,13,13,24
8 | 7) 21,10,17,25,28
9 | 8) 16,17,17,17,30
10 | 9) 30,11,25,39,40
11 | --- PASS: TestHornsEDodecagons (566.19s)

```




Figure 10: Dodecagon: Horns(6,5,5,5,8).




Figure 11: Dodecagons. Horns (15,4,13,21,20). Special case $d > e$.




Figure 12: Dodecagons. Horns (15,9,12,16,20).




Figure 13: Dodecagons. Horns (15,14,13,11,20).




Figure 14: Dodecagons. Horns (15,18,15,7,20).




Figure 15: Dodecagon. Horns(10,13,13,13,24).




Figure 16: Dodecagon. Horns(21,10,17,25,28).




Figure 17: Dodecagon. Horns(16,17,17,17,30).