

# Analysis of Massachusetts' Standardized Testing Multi-Group Structural Equation Modelling

Landon Hurley <sup>1</sup>   Herb Susmann <sup>2</sup>

<sup>1</sup>Psychology, SUNY Geneseo

<sup>2</sup>Mathematics, SUNY Geneseo

April 12, 2014

# Outline

Data

Measurement Invariance

Results

Discussion

Regression Tree

Discussion

Conclusions

References

# Introduction

- ▶ Massachusetts standardized test data

# Introduction

- ▶ Massachusetts standardized test data
- ▶ Is the test psychometrically valid?

# Introduction

- ▶ Massachusetts standardized test data
- ▶ Is the test psychometrically valid?
- ▶ **No! It doesn't measure the same thing in the same way across groups.**

# Data

- ▶ 10,515 10th grade student test results

# Data

- ▶ 10,515 10th grade student test results
- ▶ English, Math, and Science components

# Data

- ▶ 10,515 10th grade student test results
- ▶ English, Math, and Science components
- ▶ Answers from every question for each section



# Data

- ▶ 10,515 10th grade student test results
- ▶ English, Math, and Science components
- ▶ Answers from every question for each section
- ▶ Basic demographic information from questionnaire

# Data

- ▶ 10,515 10th grade student test results
- ▶ English, Math, and Science components
- ▶ Answers from every question for each section
- ▶ Basic demographic information from questionnaire

# Measurement Invariance

- ▶ Testing: set of standardized questions designed to elicit responses for use in measuring the traits, capacities, or achievements of an individual.
- ▶ Tests imply a global solution: the individual should never affect how items on the test perform.
- ▶ In psychometrics, validity is tested in one of two way: predictive invariance & measurement (factorial) invariance.

# Measurement Invariance Process

1. Configural invariance: Tests whether the same factor model is found within each subgroup.
2. Weak invariance: Tests whether in addition to the same factor structure, the same items load equivalently onto the same structure.
3. Strong invariance: Tests that the intercepts, item loadings, and factor structure are equivalent across groups.
4. Strict invariance: Tests the assumption that in addition to the preceding steps, the residual variances are equivalent across groups.

# Assessing model fit

- ▶  $\chi^2$

$$\chi^2_{d.f.} = 2(N - 1) \cdot F_0 \quad (1)$$

Scales with  $N$ , making it difficult to fit

- ▶ CFI:

$$1 - \frac{\chi^2_t - df_t}{\chi^2_n - df_n} \quad (2)$$

- ▶ NNFI:


$$1 - \frac{\frac{\chi^2_t}{df_t}}{\frac{\chi^2_n}{df_n}} \quad (3)$$

- ▶ RMSEA:

$$\sqrt{\frac{\chi^2_{df} - df_t}{df_t(N - 1)}} \quad (4)$$


# Results

## English Oblimin




# Results

## Mathematics Oblimin



# Results

## Biology Oblimin





# Results

English	$\chi^2$	CFI	TLI	RMSEA
Comparison	-1670.297	0.030	0.034	-0.006
Strict fit indices	6007.770	0.918	0.912	0.030

# Results

Mathematics	$\chi^2$	CFI	TLI	RMSEA
Comparison	-998.348	.014	.017	.005
Strict fit indices	4130.561	.949	.946	.033

# Results

Biology	$\chi^2$	CFI	TLI	RMSEA
Comparison	-2132.914	.026	.03	.006
Strict fit indices	7686.782	.926	.922	.031

# Discussion


Strict invariance: what does it mean for these tests?

- ▶ SES: Lack of strict invariance typically indicates that an unmeasured latent variable is differentially affecting the test outcome. In other words, the existence of unmeasured effects is specified by the existence of error variance heterogeneity. This is despite the invariance of both the factor model, the intercepts, and slopes.
- ▶ Meredith (1993): When using measures to make decisions strict factorial invariance is essential.

# Regression Tree

- ▶ CART allows for complex relationships to be modelled within data without parametric assumptions or linearity.
- ▶ Multivariate regression trees are ideal in this instance because it can model both the complex relationships, but also non-response, in relation to a multiple vectored response surface: English, Mathematics, & Science.

# Results



Error : 0.903 CV Error : 0.904 SE : 0.0113

# Discussion

- ▶ Missing data: Question 3 deals with after college goals. These results indicate that students who either intend to go to attain a baccalaureate degree OR skip the question perform better on the three sub-tests.
- ▶ Ethnicity: Amongst students who do plan on a baccalaureate degree, ethnicity efficiently segments the science scores.
- ▶ Quest14: How often do you use scientific instruments?
- ▶ Confirmation: Each of the responses were regressed onto multiply imputed forms of the questionnaire, which had equivalent  $R^2$ , indicating effects were linear only.

# Conclusions

- ▶ Fails strict invariance



# Conclusions

- ▶ Fails strict invariance
- ▶ Regression tree: split on ethnicity

# References I

- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological bulletin*, 107(2), 238.
- Chen, F. F. (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling*, 14(3), 464–504.
- Cronbach, L. J. (1947). Test reliability: Its meaning and determination. *Psychometrika*, 12(1), 1–16.
- De'Ath, G. (2002). Multivariate regression trees: a new technique for modeling species-environment relationships. *Ecology*, 83(4), 1105–1117.
- DeShon, R. P. (2004). Measures are not invariant across groups without error variance homogeneity. *Psychology Science*, 46, 137–149.
- Gaudet, R. D. (2000). Effective school districts in massachusetts. *The Donahue Institute, Boston*.

## References II


- Kim, K. H., & Bentler, P. M. (2002). Tests of homogeneity of means and covariance matrices for multivariate incomplete data. *Psychometrika*, 67(4), 609–623.
- Little, R. J. (1988). A test of missing completely at random for multivariate data with missing values. *Journal of the American Statistical Association*, 83(404), 1198–1202.
- Meredith, W. (1993). Measurement invariance, factor analysis and factorial invariance. *Psychometrika*, 58(4), 525–543.
- Pornprasertmanit, S., Miller, P., Schoemann, A., & Rosseel, Y. (2013). semtools: Useful tools for structural equation modeling. [Computer software manual]. Retrieved from <http://CRAN.R-project.org/package=semTools> (R package version 0.4-0)

## References III

- Revelle, W. (2014). psych: Procedures for psychological, psychometric, and personality research [Computer software manual]. Evanston, Illinois. Retrieved from <http://CRAN.R-project.org/package=psych> (R package version 1.4.2)
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. Retrieved from <http://www.jstatsoft.org/v48/i02/>
- Rubin, D. B. (2009). *Multiple imputation for nonresponse in surveys* (Vol. 307). John Wiley & Sons.
- Spring 2009 mcas tests: Summary of state results.* (2009, September).
- van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by chained equations in r. *Journal of Statistical Software*, 45(3), 1–67. Retrieved from <http://www.jstatsoft.org/v45/i03/>


# Appendix A

## English Oblimin



# Appendix A

## Mathematics Oblimin



# Appendix A

## Biology Oblimin

