

¹ Cont2SAS: A python package for calculating SAS parameters from continuum nanostructures

³ **Arnab Majumdar**  ^{1,3}, **Martin Müller** ^{1,2,3}, and **Sebastian Busch**  ^{1¶}

⁴ 1 German Engineering Materials Science Centre (GEMS) at Heinz Maier-Leibnitz Zentrum (MLZ),
⁵ Helmholtz-Zentrum Hereon GmbH, Lichtenbergstr. 1, 85748 Garching, Germany 2 Institute of Materials
⁶ Physics, Helmholtz-Zentrum Hereon GmbH, Max-Planck-Str. 1, 21502 Geesthacht, Germany 3 Institut
⁷ für Experimentelle und Angewandte Physik (IEAP), Christian-Albrechts-Universität zu Kiel, Leibnizstr.
⁸ 19, 24098 Kiel, Germany ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a

Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

⁹ Summary

¹⁰ Cont2SAS is a software tool built around the existing software solution Sassen - known for calculating scattering patterns from simulated atomic structures. The goal of Cont2SAS is to provide a software similar to Sassen, but for calculating Small Angle Scattering (SAS) data from simulated Continuum (Cont) nanostructures. Cont2SAS can calculate SAS patterns, i.e. SAS intensity (I) vs. scattering vector magnitude (Q) using a novel numerical method. Cont2SAS can also calculate effective cross-section (σ_{eff}), i.e. the count rate of scattered radiation per incident unit flux. σ_{eff} is calculated by integrating instrument-agnostic SAS patterns taking instrument geometry into account. Cont2SAS can be used for different purposes, such as validating simulations, tuning simulation parameters, and analyzing SAS data.

¹⁸ Statement of need

Small Angle Scattering (SAS) experiments with neutrons (SANS) and X-rays (SAXS) are useful techniques for probing material nanostructures (Chen et al., 2012). With the Time Resolved (TR) variant of SAS, the time evolution of nanostructures can also be probed (Hollamby, 2013). However, a direct retrieval of nanostructure from SAS data is not possible due to the so-called phase problem (Billinge & Levin, 2007). Therefore, it is fruitful to combine SAS data with simulations to study nanostructures (Majumdar et al., 2024). This approach is also useful for validating theories underlying simulations (Reich et al., 2022).

Simulated structures in real space can be generated using atomistic and continuum simulation. However, continuum simulation is the only option when the nanostructure evolution is influenced by bigger features or the SAS data is recorded for long time period, as atomistic simulation can not simulate large structures for long time (Ghavanloo et al., 2019). Despite this indispensability, there is no software available for calculating SAS data from continuum simulations. Cont2SAS aims to fill this void by providing a tool that offers fast and error-free comparison of continuum simulations and SAS data. The ultimate goal is to study nanomaterials and validating theories using SAS data, particularly when length-scales and time-scales, inaccessible to atomistic simulations, need to be simulated.

³⁶ State of the field

³⁷ From atomistic simulations, different software such as nMoldyn (Hinsen, 2023; Rög et al., 2003), MDANSE (Goret et al., 2017; Neutron & Facility, 2023), LiquidLib (Walter et al., 2018; Z-laboratory, 2023), and Sassen (Lindner, 2012, 2017; Lindner & Smith, 2012; Majumdar et

al., 2024; Majumdar & Lindner, 2023) can calculate SAS patterns but they can not calculate the same from continuum simulations. An in-house code is available for continuum simulations but it was built for a 2D membrane system lacking generality (Dorrell et al., 2020). Hence, Cont2SAS is built using Sassena as a backend calculator. Sassena was chosen over other software due to its notable computation speed and robustness (Majumdar et al., 2024).

45 Software features




Figure 1: Workflow of SAS pattern calculation: [left] Mesh generation, [middle] SLD assignment, [right] numerical calculation of SAS pattern. The numerical calculation matches well with the known analytical formula for spherical nanoparticles (Guinier et al., 1955).

46 Cont2SAS calculates SAS patterns taking simulated nanostructures as an input. The simulated
47 structure must provide either Scattering Length Density (SLD) (β) values or a set of variables
48 from which SLD values can be calculated. The simulated input is processed to a data taylor-
49 made for Sassena. Sassena calculates the SAS intensity (I) as a function of scattering vector
50 magnitude (Q), i.e. SAS pattern, for different time steps (see Figure 1).




Figure 2: Workflow of effective cross-section (σ_{eff}) calculation: [left] Calculated SAS pattern at different time steps, [right] σ_{eff} calculated from SAS patterns at different time steps.

51 Cont2SAS can also calculate the time evolution of effective cross-section (σ_{eff}), which is defined
52 as the count rate per incident unit flux. This feature was not available in the in-house code
53 (Dorrell et al., 2020) or Sassena (Majumdar & Lindner, 2023). The time evolution of count
54 rate is useful when the chemical composition changes over time instead of the nanostructure,
55 e.g. while storing hydrogen in ball-milled powder sample (Aslan et al., 2019). Figure 2
56 demonstrates the workflow of such calculation from a series of SAS patterns. The calculated
57 σ_{eff} must be multiplied by an empirical factor before comparing with measured count rate.

58 Conclusion

59 Cont2SAS provides a software platform for calculating SAS pattern from continuum simulations
60 of nanostructures. The addition of effective cross-section in the software package further
61 enables the analysis of count rate. Cont2SAS can be used for analyzing SAS data and validating
62 simulations to study nanomaterials.

63 Acknowledgements

64 This publication was written in the context of the work of the consortium DAPHNE4NFDI
65 in association with the German National Research Data Infrastructure (NFDI) e.V. NFDI is
66 financed by the Federal Republic of Germany and the 16 federal states and the consortium
67 is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) -
68 project number 460248799. The authors would like to thank for the funding and support.
69 Furthermore, thanks go to all institutions and actors who are committed to the association
70 and its goals.

71 References

- 72 Aslan, N., Horstmann, C., Metz, O., Kotlyar, O., Dornheim, M., Pistidda, C., Busch, S.,
73 Lohstroh, W., Müller, M., & Pranzas, K. (2019). High-pressure cell for in situ neutron
74 studies of hydrogen storage materials. *Journal of Neutron Research*, 21(3-4), 125–135.
75 <https://doi.org/10.3233/JNR-190116>
- 76 Billinge, S. J., & Levin, I. (2007). The problem with determining atomic structure at the
77 nanoscale. *Science*, 316(5824), 561–565. <https://doi.org/10.1126/science.1135080>
- 78 Chen, Z. H., Kim, C., Zeng, X., Hwang, S. H., Jang, J., & Ungar, G. (2012). Characterizing size
79 and porosity of hollow nanoparticles: SAXS, SANS, TEM, DLS, and adsorption isotherms
80 compared. *Langmuir*, 28(43), 15350–15361. <https://doi.org/10.1021/la302236u>
- 81 Dorrell, M. W., Beaven, A. H., & Sodt, A. J. (2020). A combined molecular/continuum-
82 modeling approach to predict the small-angle neutron scattering of curved membranes.
83 *Chemistry and Physics of Lipids*, 233, 104983. <https://doi.org/10.1016/j.chemphyslip.2020.104983>
- 85 Ghavanloo, E., Rafii-Tabar, H., & Fazelzadeh, S. A. (2019). *Computational continuum
86 mechanics of nanoscopic structures*. Springer. <https://doi.org/10.1007/978-3-030-11650-7>
- 87 Goret, G., Aoun, B., & Pellegrini, E. (2017). MDANSE: An interactive analysis environment
88 for molecular dynamics simulations. *Journal of Chemical Information and Modeling*, 57(1),
89 1–5. <https://doi.org/10.1021/acs.jcim.6b00571>
- 90 Guinier, A., Fournet, G., Walker, C. B., & Yudowitch, K. L. (1955). *Small-angle scattering of
91 x-rays*. Wiley New York.
- 92 Hinsen, K. (2023). *nMOLDYN3*—version 3.0.12. <https://github.com/khinsen/nMOLDYN3/>.
- 93 Hollamby, M. J. (2013). Practical applications of small-angle neutron scattering. *Phys. Chem.
94 Chem. Phys.*, 15, 10566–10579. <https://doi.org/10.1039/C3CP50293G>
- 95 Lindner, B. (2012). *Towards a unification of supercomputing, molecular dynamics simulation
96 and experimental neutron and x-ray scattering techniques* [PhD thesis]. University of
97 Tennessee, United States.
- 98 Lindner, B. (2017). *Sassena*. <https://github.com/camm/sassena>.
- 99 Lindner, B., & Smith, J. C. (2012). Sassena—x-ray and neutron scattering calculated from
100 molecular dynamics trajectories using massively parallel computers. *Computer Physics*

- 101 *Communications*, 183(7), 1491–1501. <https://doi.org/10.1016/j.cpc.2012.02.010>
- 102 Majumdar, A., & Lindner, B. (2023). *Sassena*. <https://codebase.helmholtz.cloud/DAPHNE4NFDI/sassena>. <https://doi.org/10.5281/zenodo.10037485>
- 104 Majumdar, A., Müller, M., & Busch, S. (2024). Computation of x-ray and neutron scattering
105 patterns to benchmark atomistic simulations against experiments. *International Journal of
106 Molecular Sciences*, 25(3), 1547. <https://doi.org/10.3390/ijms25031547>
- 107 Neutron, U. S. I., & Facility, M. (2023). *MDANSE-version 1.5.2*. <https://github.com/ISISNeutronMuon/MDANSE/>.
- 109 Reich, V., Majumdar, A., Müller, M., & Busch, S. (2022). Comparison of molecular dy-
110 namics simulations of water with neutron and x-ray scattering experiments. *EPJ Web of
111 Conferences*, 272, 01015. <https://doi.org/10.1051/epjconf/202227201015>
- 112 Rög, T., Murzyn, K., Hinsen, K., & Kneller, G. R. (2003). nMoldyn: A program package
113 for a neutron scattering oriented analysis of molecular dynamics simulations. *Journal of
114 Computational Chemistry*, 24(5), 657–667. <https://doi.org/10.1002/jcc.10243>
- 115 Walter, N. P., Jaiswal, A., Cai, Z., & Zhang, Y. (2018). LiquidLib: A comprehensive
116 toolbox for analyzing classical and ab initio molecular dynamics simulations of liquids and
117 liquid-like matter with applications to neutron scattering experiments. *Computer Physics
118 Communications*, 228, 209–218. <https://doi.org/10.1016/j.cpc.2018.03.005>
- 119 Z-laboratory. (2023). *Liquidlib-version 1.0*. <https://github.com/Z-Laboratory/LiquidLib/tree/master>.